

Kac's Walk on Rotation Matrices Mixes in $\Theta(n^2)$ Steps: A Proof Discovered with AI

Tianle Liu

Department of Statistics and Operations Research
University of North Carolina at Chapel Hill
tianletl@unc.edu

29 August 2026

Abstract

Let $N = \binom{n}{2} = \dim \text{SO}(n)$. We prove that the coordinate-plane Kac walk on $\text{SO}(n)$ has total-variation mixing time of order N : for every fixed $0 < \varepsilon < 1$,

$$t_{\text{mix}}^{(n)}(\varepsilon) = \Theta_\varepsilon(n^2).$$

The lower bound is the dimensional singularity obstruction before N steps. The upper bound removes the final logarithm from the previously known $O(n^2 \log n)$ estimate.

The proof combines the discrete Malliavin coupling and low-degree pseudo-mixing inputs with a new log-free analysis of the derivative shells. Its static core is a circuit-anchored, arbitrary-spectrum root/pass identity for the physical five-box prime. Keeping one normalization base per original circuit permits simultaneous scalar regluing without paying for artificial cuts. Its temporal core is an exact chronological calculus: passive singleton runs acquire a coboundary/Riesz gain, while root-interrupted components are allocated by vertex-labelled packets before absolute values are taken. The curvature split into pure-Weyl and Ricci parts is kept at its physical tensor type. All-Weyl packets retain a full N^{-1} resource; mixed packets contain a typed $O(n^{-1/2})$ Ricci debit; and the final packetless Ricci cell is closed by a joint invariant-column estimate on its two root-hit circuits and an exact causal restoration of the marked root time. These estimates yield an $O(n)$ squared first-derivative shell and a summable all-order marked-shell expansion through logarithmic degree. The resulting score energy is $O(n/c^2)$ after cN steps. A weighted submersion integration-by-parts argument and the Haar log-Sobolev inequality then give the uniform total-variation upper bound. No cutoff profile or cutoff window is asserted.

The proof was discovered through an approximately three-day, author-directed collaboration with OpenAI Codex using GPT-5.6 Sol, followed by approximately one day of targeted defect repair and audit. The author has reviewed the statements, cited sources, global proof architecture, and principal reductions, but a complete line-by-line human audit of several technical derivations remains in progress. This version should therefore be regarded as a preliminary proof report.

2020 Mathematics Subject Classification. Primary 60J05; Secondary 60B15.

Keywords. Kac's walk, total-variation mixing, rotation group, Markov chains, discrete Malliavin calculus.

Contents

1 Introduction

2

1.1	The walk and the main theorem	2
1.2	Historical context	3
1.3	What removes the logarithm	4
1.4	Detailed proof architecture	4
1.5	AI-assisted discovery and verification status	5
1.6	Organization	6
2	Static circuit calculus	6
2.1	Pure-Weyl coefficient calculus	6
2.2	Circuit-anchored resources and port states	7
2.3	The physical root/pass bridge	10
2.4	One-resource module stability	14
2.5	Maximum smoothing and the positive-credit induction	18
2.6	Colors, defects, marked roots, and the packet/ribbon estimate	20
3	Moments, renewal, and marked shells	27
3.1	Traffic moments and a fixed covariance band	27
3.2	Chronological spectral calculus and passive renewal	31
3.2.1	Common spectral blocks and chronological exposure	31
3.2.2	Passive singletons, typed passes, and finite Green gaps	37
3.3	Root-local and packetless aggregate-Ricci calculus	46
3.3.1	Root evacuation, deposits, and reflected Ricci coupons	46
3.3.2	Packetless hard and soft cells	53
3.3.3	Invariant-column closure of the low cells	58
3.4	Global allocation and pure-Weyl weak-order renewal	73
3.4.1	Fixed components and token allocation	73
3.4.2	Root-fused allocation and global renewal	78
3.5	Aggregate-Ricci synthesis and marked-shell closure	82
3.5.1	Aggregate-Ricci renewal	82
3.5.2	Intact marked-shell closure	85
3.6	Inverse expansion and score energy	86
4	Spectral surgery, endpoint calculus, and total variation	88
5	Dependency audit	91
	Use of AI	92
	References	92

1 Introduction

1.1 The walk and the main theorem

Let

$$N = \dim \mathrm{SO}(n) = \binom{n}{2}.$$

All proof estimates below are stated for $n \geq 4$. The main theorem is asymptotic in n , so the finitely many smaller dimensions are absorbed by enlarging its threshold. Throughout, C denotes a

numerical constant and A_L, C_L constants depending only on the fixed logarithmic-degree parameter L ; these constants may increase from one display to the next. At the end of the proof, A_{sh} is one fixed maximum of the finitely many exponential constants used in the marked-shell estimates. We also fix once and for all a numerical integer B_0 which bounds the finite list of special-root symbols introduced by the root calculus. None of these constants depends on n, m , or c . At each discrete step of Kac's walk, choose an unordered coordinate pair $I = \{i, j\}$ uniformly from the N pairs, choose $\theta \sim \text{Unif}[0, 2\pi)$, and multiply by the coordinate-plane rotation $R_I(\theta)$. Haar probability H_n is stationary.

Write

$$d_n(t) = \sup_{X \in \text{SO}(n)} \|\mathcal{L}_X(X_t) - H_n\|_{\text{TV}}, \quad t_{\text{mix}}^{(n)}(\varepsilon) = \inf\{t : d_n(t) \leq \varepsilon\}.$$

Theorem 1.1 (No-log theorem). *For every fixed $0 < \varepsilon < 1$, there are constants $c_\varepsilon < \infty$ and n_ε such that for $n \geq n_\varepsilon$,*

$$N \leq t_{\text{mix}}^{(n)}(\varepsilon) \leq c_\varepsilon N.$$

In particular,

$$t_{\text{mix}}^{(n)}(\varepsilon) = \Theta_\varepsilon(n^2).$$

The logarithm is not intrinsic to total-variation mixing on the rotation group: the coordinate-plane walk reaches equilibrium on its dimensional scale $N = \binom{n}{2}$, up to constants depending only on the requested accuracy.

The lower bound in [Theorem 1.1](#) is exact at the level relevant here. Conditional on a word of t coordinate planes, the endpoint is the image of a smooth map from the t -torus into the N -dimensional group. It is Haar-null for $t < N$; the finite mixture over plane words is therefore still singular. The content of the paper is the matching upper bound. We do not identify a cutoff constant, profile, or window.

1.2 Historical context

Kac introduced the coordinate-rotation process on the sphere as a kinetic model for the Boltzmann equation [\[Kac56\]](#). Hastings subsequently used the matrix version as a sampler for random orthogonal matrices [\[Has70\]](#). The two settings share their elementary rotation, but their mixing geometry is different: the sphere has dimension $n - 1$, whereas $\text{SO}(n)$ has dimension $N \asymp n^2$, and a point mass for the group walk remains singular for at least N steps.

The spectral theory is mature. Diaconis and Saloff-Coste initiated the quantitative analysis of Kac's master equation and the associated group walk [\[DS00\]](#). Janvresse obtained the correct spectral-gap scale [\[Jan01\]](#); Carlen, Carvalho, and Loss identified the gap exactly [\[CCL03\]](#); and Maslen computed the spectrum [\[Mas03\]](#). These results determine relaxation, but do not by themselves control total variation from a singular starting law. That distinction is decisive here: an L^2 argument cannot begin before a density has formed, and the creation and conditioning of that density are the difficult part of the group problem.

Several complementary approaches clarified the scale. Pak and Sidenko developed compact-group couplings and obtained polynomial weak-mixing bounds [\[PS07\]](#). Oliveira proved $O(n^2 \log n)$ convergence in quadratic Wasserstein distance [\[Oli09\]](#). Jiang gave the first polynomial total-variation estimate for the group walk [\[Jia17\]](#). Pillai and Smith then introduced a non-Markovian coupling for the singular-to-smooth transition and bounded the group mixing time between order n^2 and $n^4 \log n$ [\[PS16\]](#). On the sphere, the analogous coupling yields the sharp order $n \log n$ [\[PS17\]](#), improving

Jiang’s earlier sphere bound [Jia12]. The sphere logarithm reflects a different coordinate-exposure mechanism and does not force a logarithm on the N -dimensional group walk.

Two nearby but more symmetric walks are also instructive. Rosenthal’s fixed-angle random-rotation walk and the uniform-plane walk are conjugation-invariant; character theory yields sharp mixing and cutoff statements in that setting [Ros94; HJ17]. The coordinate-plane Kac walk studied here is not conjugation-invariant, so those Fourier reductions are unavailable. This is why the proof works with moving frames, low-degree Haar moments, and the full chronological incidence of selected coordinate planes.

The immediate starting point is the 2026 result of Pillai and Smith, which proves the $O(n^2 \log n)$ total-variation upper bound through a two-stage coupling and a discrete Malliavin analysis [PS26a]. Pillai, Smith, and Vaikuntanathan provide the quantitative low-degree pseudo-mixing estimate used to establish the covariance band after burn-in [PSV26]. Their transvection-walk work gives a useful parallel local-to-global entropy architecture, but it is not an input to this proof [PS26b]. The present paper keeps the first two inputs and replaces the logarithmic derivative-shell estimate by the static and temporal calculus developed below.

1.3 What removes the logarithm

At time $m \asymp N$, the Malliavin covariance is already well conditioned with high probability. The remaining loss in the previous argument came from bounding high-order derivatives after exposing their time order and their tensor contractions too early. A naive absolute value can pay once for every artificial cut, every chronological order, or every appearance of a common coordinate plane; any one of these payments is fatal when the expansion runs to degree $L \log n$.

The repair has two inseparable parts. First, the static contraction is based on original circuits rather than on local bends. The physical five-box prime is expanded by an exact root/pass identity while all ten gap variables and the incoming token remain in place. One base is then charged per original circuit, regardless of how many times it was cut for the proof. Second, chronology is summed only after same-time packets and root incidences have been fused. Passive runs gain through an exact coboundary/Riesz identity; mixed Weyl–Ricci packets retain a physical $n^{-1/2}$ debit; and the unique low-debit packetless Ricci cell is closed on a joint invariant column and then by a summable causal root-time kernel before trace closure.

The proof never estimates a temporary port as an independent output. Every spatial cut, chronological gap, and selected-plane incidence is returned to one terminal scalar, with its original circuit normalization and packet resource still attached.

This “intact sum” principle is the quantitative reason the degree summation is exponential in the derivative order rather than factorial, and hence why the $L \log n$ truncation is compatible with $O(N)$ time.

1.4 Detailed proof architecture

The proof can be read as the following sequence.

1. *Coefficient geometry.* Decompose every centered Kac coefficient into its pure-Weyl and traceless-Ricci pieces, keeping the normalizations of $\Lambda^2 \mathbb{R}^n$ and the probabilistic Kac frame explicit.
2. *Circuit-anchored root/pass calculus.* Theorem 2.4 proves the physical root and pass expansions for the five-box prime in all twenty signed rail states. Lemma 2.7 then closes all declared ports with one trace-class base per original circuit.

3. *All-order packets.* The module-stability and smoothing arguments turn connected coefficient components into positive-credit packets. The scalar packet/ribbon estimate, [Theorem 2.24](#), keeps a full N^{-1} resource at every marked pure-Weyl packet.
4. *Pseudo-Haar covariance band.* Orthogonal Weingarten calculus [[CS06](#); [CM09](#)] and quantitative pseudo-mixing control all moments through degree $L \log n$. Matrix martingale concentration [[Tro11](#)] then places the Malliavin covariance in a fixed deterministic band after $O(N)$ burn-in.
5. *Chronological renewal.* The Kac spectral blocks and exact Haar-residual decomposition turn every finite Green gap into a bounded gate. The singleton coboundary identity supplies a gain on alternate pass vertices, strong enough to absorb all strict chronological orders.
6. *Root and Ricci allocation.* A vertex-labelled fixed-column presentation allocates every unresolved component without duplicating a packet endpoint or circuit base. Pure-Weyl components close by global renewal. Aggregate-Ricci components split into packet-bearing, scalar dimer, internal-hard, threshold, and one final low-cell family; the last is closed by the normalized invariant-column insertion followed by an exact causal restoration of the marked root time. The latter keeps the native selected gate inside a summable two-rail kernel and uses martingale orthogonality only for the genuine aggregate increment.
7. *Marked shells and inverse covariance.* The two renewal branches give [Theorem 3.46](#). The first squared derivative shell is $O(n)$; the all-order shell bound controls the inverse series through logarithmic degree, while a separate finite- N tail estimate handles the remainder. At time cN the score energy is $O(n/c^2)$.
8. *Endpoint comparison.* A cutoff localizes to the covariance band, and weighted integration by parts transfers the score estimate to the endpoint density. The Ricci curvature of $SO(n)$ yields the Haar log-Sobolev inequality through the Bakry-Émery method [[BGL14](#)]; Pinsker’s inequality finishes the total-variation bound.

The resulting dependency chain is

$$\begin{aligned}
\text{PK5}_{\text{root/pass}} &\implies \text{one-resource comb stability} \\
&\implies \text{positive-credit packets} \implies \text{PR}, \\
\text{PR} + \text{covariance band} + \text{global renewal} &\implies \text{MS}_{\text{intact}}, \\
\text{MS}_{\text{intact}} &\implies \text{Malliavin/TV closure} \implies \Theta(n^2).
\end{aligned} \tag{1.1}$$

1.5 AI-assisted discovery and verification status

The proof was developed through an extended, author-directed interaction with generative-AI systems. The primary system was OpenAI Codex running GPT-5.6 Sol. According to the author’s reconstruction of the workflow, approximately three days of intensive interaction were devoted to proof search, construction, and organization, followed by approximately one additional day devoted to targeted defect repair and consistency audit. The Codex workflow used parallel agents for bounded subproblems and adversarial audits. Its mathematical roles included exploring proof routes, maintaining the proof ledger and dependency map, carrying out tensor and ribbon bookkeeping, searching for counterexamples to proposed intermediate lemmas, producing exact replay calculations where suitable, and revising the exposition. Claude Opus 5 was subsequently used by a collaborator for two additional audit passes.

The author supplied the mathematical problem, source papers, target $O(n^2)$ invariant, and strategic direction; decided which routes and claims to retain; reviewed the manuscript and cited sources; and checked the theorem statement, global proof architecture, principal reductions, and dependency chain. The AI-generated work was not accepted in a single pass: proposed shortcuts were repeatedly tested, false strengthenings were removed after counterexamples, and the surviving argument underwent exact computational replays where its finite algebraic content permitted them. Agreement among AI agents or passage through an AI audit was treated as evidence for further checking, not as a proof by itself.

The verification status of this version should nevertheless be stated precisely. A complete independent, line-by-line human rederivation of every lengthy local argument has not yet been completed. In particular, the automated replays and the two external AI-assisted audit passes do not replace such a human audit. Accordingly, this arXiv version is presented as a preliminary proof report, and reports of errors or independent verification are especially welcome. The author takes responsibility for the contents of the submission and will record corrections and the progress of human verification in subsequent versions. No generative-AI system is listed as an author or cited as authority for a mathematical claim.

1.6 Organization

Section 2 contains the complete static circuit calculus: coefficient geometry, port states, the physical bridge, module stability, smoothing, and the packet/ribbon estimate. Section 3 contains the probabilistic core: traffic moments, chronological renewal, root-local Ricci analysis, global allocation, marked shells, and score energy. Section 4 carries out endpoint comparison and total-variation closure. The standalone dependency audit in Section 5 records the shortest proof chain and the location of each input.

2 Static circuit calculus

2.1 Pure-Weyl coefficient calculus

Put $H = \mathbb{R}^n$ and $W = \Lambda^2 H \cong \mathfrak{so}(n)$, with

$$\langle A, B \rangle_\wedge = -\frac{1}{2} \text{Tr}(AB)$$

for skew matrices. Let $C = C^* \in \text{End}(W)$ be a pure-Weyl curvature operator:

$$\text{Alt } C = 0, \quad \text{Ric } C = 0. \tag{2.1.1}$$

Choose an orthonormal eigenframe

$$CA_a = \lambda_a A_a, \quad \lambda_a = \varepsilon_a r_a, \quad \varepsilon_a \in \{\pm 1\}, \quad r_a = |\lambda_a|, \tag{2.1.2}$$

omitting zero eigenvalues, and put

$$K = \|C\|_{\text{op}}, \quad h = \|C\|_2.$$

Define the signed Kraus map

$$\Phi_C(Z) = \sum_a \lambda_a A_a Z A_a^T. \tag{2.1.3}$$

The algebraic Bianchi identity identifies this map with the action of C on W : $\Phi_C(Z) = CZ$ for $Z \in \mathfrak{so}(n)$. The positive second-moment cap used below is constructed directly from the paired physical front.

2.2 Circuit-anchored resources and port states

This section fixes the one-hole carrier, the physical port states, and the once-per-original-circuit normalization used by the mixing proof. All objects retain their actual tensor type; a scalar cap is never used to identify distinct open carriers.

Definition 2.1 (Typed circuit primitives). For an ordered finite set S of oriented carrier labels, write $\mathcal{H}_S = \bigotimes_{u \in S} H_u$, with every factor equipped with the Euclidean Hilbert norm. A *normalized Brauer primitive* is a signed carrier permutation, a wedge reversal, full transpose, a share turn, a through string, the orthogonal antisymmetrizer, or a cup/cap normalized so that its operator norm is one. A share turn is the fixed signed boundary permutation, possibly followed by full transpose, induced when a principal pivot reverses one of the two ordered boundary intervals. The two coefficient primitives are

$$\hat{C}_{uv} = \iota_{uv} C \pi_{uv} : H_u \otimes H_v \rightarrow H_u \otimes H_v,$$

where π_{uv} is the normalized orthogonal projection and identification onto $\Lambda^2 H$, and $\iota_{uv} = \pi_{uv}^*$; hence $\hat{C}_{uv}$ is zero on $\text{Sym}^2 H$ and has norm K . The other coefficient primitive is the restriction/re-expression of Equation (2.1.3) as the Bianchi transfer

$$(\Phi_C Y)_{ik} = \sum_{m,n} C_{in,mk} Y_{mn}, \quad Y \in \Lambda^2 H.$$

The first Bianchi identity identifies the latter with the physical action of C . A *safe gate* is a typed composite of these primitives, including the AA words and correctly oriented AB words used below. Its degree is the number of coefficient primitives. Thus a degree- d safe gate $G : \mathcal{H}_S \rightarrow \mathcal{H}_T$ satisfies $\|G\|_{\text{op}} \leq K^d$; untouched factors are ancillary identities.

A coefficient box has four darts $L_{i,0}, L_{i,1}, R_{i,0}, R_{i,1}$. For a column pairing q , a *packet* is a connected component of the box-incidence graph whose edges are the q -pairs joining different boxes. A q -pair joining the two darts of one wedge is forbidden and gives zero. An allowed *smoothing* chooses at each box either

$$h_i^{\parallel} = \{L_{i,0}R_{i,0}, L_{i,1}R_{i,1}\}, \quad h_i^{\times} = \{L_{i,0}R_{i,1}, L_{i,1}R_{i,0}\}.$$

The alternating circuits of $q \vee h$ are the *smoothing carriers*; adjoining one carrier means adjoining one identity factor I_H . The smoothing support $\text{supp}_h(i)$ of a box is the one- or two-element set of $q \vee h$ carriers containing its two h -passages. A *ported pair comb* on two chosen carriers consists of all boxes whose smoothing support is that pair, together with every singleton-support box assigned to it exactly once; removing all other support groups leaves their boundary darts as linearly ordered typed ports. These definitions fix the domains and codomains of every safe word, fold, comb, and attachment below.

An *ambient rooted ribbon record* Γ consists of a common finite dart set, its coefficient boxes and local wedge signs, a row pairing p , a column pairing q , a rooted trace pairing r , and the declared boundary/root data. Thus $p \vee r$ is the disjoint union of the original trace circuits in that record; its components, not temporary proof cuts, receive the normalization bases below. The more detailed Weingarten data carried by Γ are specified in Definition 2.18.

Definition 2.2 (Paired resource and safe module). A safe gate and its C -degree are as in Definition 2.1; the amplified estimate is proved in Lemma 2.9.

For an active eigenframe of an arbitrary self-adjoint pure-Weyl C , put

$$(F_C x)_a = \lambda_a A_a^T x, \quad F_C^* F_C = Q_2 := - \sum_a \lambda_a^2 A_a^2, \quad \text{Tr } Q_2 = 2\|C\|_2^2. \quad (2.2.1)$$

A concrete carrier for these two fronts is obtained as follows. Let $E_C = \ell^2(I_C)$ be the active label space, $J_C : E_C \rightarrow \Lambda^2 H$ the isometry $J_C e_a = A_a$, and

$$\iota_\wedge A = 2^{-1/2} \sum_{i,j} A_{ij} e_i \otimes e_j, \quad W_C = (\iota_\wedge J_C) \otimes I_H,$$

and define the one-hole column

$$(T_C x)_{ijk} = \sum_\ell C_{ij,k\ell} x_\ell. \quad (2.2.2)$$

With the ordered-pair convention $C_{ij,k\ell} = \sum_a \lambda_a (A_a)_{ij} (A_a)_{k\ell}$, one has the exact typed identity

$$W_C F_C = -2^{-1/2} T_C, \quad \mathfrak{q}_C^{\text{hole}} := W_C F_C F_C^* W_C^* = \frac{1}{2} T_C T_C^*. \quad (2.2.3)$$

Thus the open token is a specified three-strand operator, up to the signed leg permutations recorded by its port state; it is not defined merely by having a prescribed scalar cap. Moreover

$$T_C^* T_C = 2Q_2, \quad T_C = P_{[2,1]_0} T_C, \quad (2.2.4)$$

where $P_{[2,1]_0}$ is the trace-free hook projection on the first three indices.

A paired-front atom contains the two correlated Hilbert–Schmidt fronts of this single F_C pair, equivalently the concrete hole $\mathfrak{q}_C^{\text{hole}}$ in Equation (2.2.3), any safe left and right gates, and an untouched typed ancillary boundary E . Let $\mathfrak{M}_C^{\text{alg}}(E)$ be the algebraic span of these atoms. Its projective norm is the infimum, over all atomic presentations, of the sum of the products of the two front Hilbert–Schmidt norms and all safe-gate operator-norm bounds. The completion is denoted $\mathfrak{M}_C(E)$. Its distinguished token $\mathfrak{q}_C$ satisfies

$$\|\mathfrak{q}_C\|_{\mathfrak{M}_C} \leq \text{Tr } Q_2 = 2\|C\|_2^2. \quad (2.2.5)$$

The two fronts form one resource: they may move or fuse but are never duplicated. Equality in Equation (2.2.5) is not asserted, and the cap identity does not identify this open pair with a differently typed polar pair.

Verification of Equations (2.2.1) to (2.2.5). Since $A_a^T = -A_a$,

$$F_C^* F_C = \sum_a \lambda_a^2 A_a A_a^T = -\sum_a \lambda_a^2 A_a^2 = Q_2.$$

Wedge orthonormality gives $\text{Tr}_H(-A_a^2) = 2$, hence $\text{Tr}_H Q_2 = 2 \sum_a \lambda_a^2 = 2\|C\|_2^2$. Use the single atomic presentation consisting of the two F_C fronts and no additional gate. Its projective cost is $\|F_C\|_{\text{HS}}^2 = \text{Tr}(F_C^* F_C) = \text{Tr } Q_2$, which proves Equation (2.2.5).

For the concrete hole, direct substitution of the spectral expansion of C gives

$$(W_C F_C x)_{ijk} = -\frac{1}{\sqrt{2}} \sum_{a,\ell} \lambda_a (A_a)_{ij} (A_a)_{k\ell} x_\ell = -\frac{1}{\sqrt{2}} (T_C x)_{ijk}.$$

This proves Equation (2.2.3). Taking adjoints and using that W_C is an isometry proves $T_C^* T_C = 2F_C^* F_C = 2Q_2$. Skewness in i, j is built into C . The first Bianchi identity says that the total alternation of $C_{ij,k\ell}$ in i, j, k is zero, while Ricci flatness says that every contraction of two of those three indices is zero. These are precisely the alternating and trace summands complementary to the trace-free hook in $H \otimes \Lambda^2 H$, proving the second identity in Equation (2.2.4). $\square$

Definition 2.3 (Circuit-anchored states and joint attachments). Let Π_{core} be the finite set generated by the three principal-pivot presentations P_C, P_H, P_G listed explicitly in [Lemma 2.11](#) below, and by exchange and reversal of the two carrier rails. A *circuit state* τ records, before evaluation,

- (i) every original $p \vee r$ circuit, its ordered oriented interval fragments, and one base fragment on that circuit;
- (ii) the ten named physical gaps, every component, defect, and retained root port, with their source and target tensor words; and
- (iii) either one ordered pair of root fronts or one existing opaque two-front token.

The finite root states Π_{root} record the twenty ordered choices of distinct core occurrences together with this data.

For later use we make the signed transport in this definition explicit. Identify the five core vertices with $\mathbb{F}_5$ by

$$\kappa_5(0) = 0, \quad \kappa_5(1) = 1, \quad \kappa_5(2) = 4, \quad \kappa_5(3) = 2, \quad \kappa_5(4) = 3. \quad (2.2.6a)$$

In these coordinates the oriented left and right rails are respectively $x \mapsto x + 1$ and $x \mapsto x + 2$. The transport group is

$$\mathcal{O}_{20} = \{g_{\alpha,\beta} : x \mapsto \alpha x + \beta : \alpha \in \mathbb{F}_5^\times, \beta \in \mathbb{F}_5\}. \quad (2.2.6b)$$

It acts on vertices through $\kappa_5^{-1} g_{\alpha,\beta} \kappa_5$. Multiplication by $\alpha = 1, 4, 2, 3$ sends the ordered pair of oriented rails (L, R) to

$$(L^+, R^+), \quad (L^-, R^-), \quad (R^+, L^-), \quad (R^-, L^+), \quad (2.2.6c)$$

respectively. If x_v^s is the gap leaving v on the oriented rail s and g sends s to t with orientation ϵ , then

$$g(x_v^s) = \begin{cases} x_{g(v)}^t, & \epsilon = +1, \\ (x_{\text{pred}_t(g(v))}^t)^*, & \epsilon = -1. \end{cases} \quad (2.2.6d)$$

At a curvature gate the induced dart permutation carries the two physical wedge pairs to the two physical wedge pairs. Reversal within a wedge pair contributes its skew sign, whereas exchange of the two wedge pairs is signless. Thus the product of the five local curvature signs is

$$\chi(g_{\alpha,\beta}) = \begin{cases} +1, & \alpha \in \{1, 4\}, \\ -1, & \alpha \in \{2, 3\}. \end{cases} \quad (2.2.6e)$$

[Equations \(2.2.6a\) to \(2.2.6e\)](#), including the formal adjoints in [Equation \(2.2.6d\)](#), are part of every transported circuit state.

For an ordered pair $r \neq s$, put

$$R = \{r, s\}, \quad S = V(K_5) \setminus R.$$

Excise the five selected coefficient gates and cut each of the ten rail segments at its two incidences with the core. The resulting typed boundary word $\mathcal{U}_{rs}$ has two half-darts on the edge internal to R , six on the three edges internal to S , and twelve on the six R – S edges. For any typed boundary word $\mathcal{U}$, let $\mathcal{H}(\mathcal{U})$ be the ordered tensor product of its indicated H , $\Lambda^2 H$, and dual factors. Fixing the complete complement also fixes an output boundary word $\mathcal{O}_{rs}$. A *joint attachment*

$$\mathcal{A}_{rs}[\Xi] : \mathcal{H}(\mathcal{U}_{rs}) \longrightarrow \mathcal{H}(\mathcal{O}_{rs}) \quad (2.2.7)$$

is the contraction made from the complete complement of those five gates, the original circuit wiring, all ten simultaneous gap contexts, signed boundary maps, and ancillary identities. In pass mode its total token degree is one and its unique opaque token is denoted Ξ ; here *token degree* means the number of occurrences of this distinguished formal token generator in the raw complement. In root mode the attachment has token degree zero. No selected core gate belongs to the attachment, and the complement supplies every named gap context exactly once. The token may lie in one gap context or may couple several gap slots. Only the finite boundary type of $\mathcal{A}_{rs}$ belongs to the port-state set; its contents remain opaque.

A *circuit-anchored one-hole presentation* is a raw typed string diagram made from physical full-wedge and correctly oriented Bianchi gates, signed rail maps and full transposes, the fixed raw circuit wiring, the ten named gap slots, and exactly one root or incoming token. Its normalized realization is

$$\text{Contr}_{\Gamma, \tau}^{\text{norm}} := n^{-F(\tau)} \text{Contr}_{\Gamma, \tau}^{\text{raw}}, \quad F(\tau) = \#(p \vee r), \quad (2.2.8)$$

with the factor n^{-1} placed once at the recorded base of each original circuit. Compact bends merely subdivide an original circuit; they are never normalized separately.

2.3 The physical root/pass bridge

Theorem 2.4 (Circuit-anchored physical root/pass bridge). *For every physical K_5 core, every declared circuit state τ , and every compatible ordered root or joint one-token attachment, there is an exact raw circuit-anchored expansion. Here B_u and P_u are the two open daughters of the Bianchi split $C_{ij, ag} = C_{ia, jg} - C_{ja, ig}$, cut on the internal rail $u \in \{h, i, j\}$, and Cl_u closes that cut in its inherited variance; Equations (2.3.7) to (2.3.9) give their coordinate construction. The six root children are indexed by $\nu = (u, \epsilon)$ with $u \in \{h, i, j\}$ and $\epsilon \in \{B, P\}$. In the canonical state, $\mathcal{W}_{(u, B)}^{\text{root}}$ (respectively $\mathcal{W}_{(u, P)}^{\text{root}}$) is the raw one-base presentation which pairs $\mathbf{q}_C^{\text{hole}} P_{\text{cyc}}^*$ with $\text{Cl}_u(B_u)$ (respectively $\text{Cl}_u(P_u)$), retaining all ten gap words. Transport the whole presentation to σ by Equations (2.2.6a) to (2.2.6e), and put*

$$a_{\sigma, (u, B)}^{\text{root}} = -\frac{2}{3}, \quad a_{\sigma, (u, P)}^{\text{root}} = \frac{2}{3}.$$

The transport sign $\chi(\sigma)$ is already part of the signed transported presentation $\mathcal{W}_{\sigma, \nu}^{\text{root}}$ and is therefore not repeated in the scalar coefficient. Here P_{cyc} is the fixed leg permutation which realizes the oriented root hit in Equation (2.3.11). In root mode the expansion is

$$\mathcal{R}_{C, \sigma}^{\text{raw}} = \sum_{\nu=1}^6 a_{\sigma, \nu}^{\text{root}} \mathcal{W}_{\sigma, \nu}^{\text{root}}, \quad \sum_{\nu=1}^6 |a_{\sigma, \nu}^{\text{root}}| \leq 4. \quad (2.3.1)$$

For pass mode let $\tilde{Q}^{rs}$ and $\tilde{H}^{rs}$ be the fully gap-open two- and three-gate factors at an ordered split $R = \{r, s\}$. Let $\mathcal{S}_{rs}$ be the fixed signed boundary reshuffle which carries the canonical $(0, 1)$ residual word, including its variances and formal adjoints, to the transported ordered pair (r, s) . For every joint attachment of token degree one,

$$\mathcal{P}_{C, rs, \mathcal{A}}^{\text{raw}}[\Xi] = \frac{1}{3} \sum_{u \in \{h, i, j\}} \mathcal{A}_{rs}[\Xi] \left(\tilde{Q}^{rs} \otimes \mathcal{S}_{rs} \text{Cl}_u(B_u - P_u) \right). \quad (2.3.2)$$

Thus if M_{mode} is the number of scalar summands and A_{mode} the sum of the absolute values of their coefficients,

$$M_{\text{root}} = M_{\text{pass}} = 6, \quad A_{\text{root}} = 4, \quad A_{\text{pass}} = 2. \quad (2.3.3)$$

Every term contains each of the ten named gaps exactly once. A root term contains the selected two fronts and the other three coefficient gates; a pass term contains the unchanged incoming token and all five coefficient gates. Every term has token degree one. The identities commute with ancillary tensoring, arbitrary joint typed attachment, signed transport of the complete state, and final one-base-per-original-circuit gluing. The distinguished cap remains

$$F_C^* F_C = Q_2 = - \sum_a \lambda_a^2 A_a^2, \quad \text{Tr } Q_2 = 2 \|C\|_2^2. \quad (2.3.4)$$

These proved identities are denoted $\text{PK5}_{\text{root/pass}}$.

Proof. We give the canonical ordered-root calculation; every tensor and every gap remains open until the indicated contraction. For roots $(0, 1)$ the physical factorization is

$$\Theta_C(\mathbf{x}) = \sum_{a,b,c,e,f,g} Q_{abc,efg}^{01}(x_0^L) H_{abc,efg}^{01}(\mathbf{x}_{\neq(0,L)}), \quad (2.3.5)$$

where

$$Q_{abc,efg}^{01} = \sum_{d_0,d_1} C_{ab,cd_0}(x_0^L)_{d_1 d_0} C_{d_1 e,fg}, \quad H_{abc,efg}^{01} = \sum_{h,i,j} C_{fh,ci} C_{bh,je} C_{ij,ag}. \quad (2.3.6)$$

The nine residual gap maps occupy the displayed scalar rails; suppressing them in the second formula is typographical only. The first Bianchi identity gives

$$C_{ij,ag} = C_{ia,jg} - C_{ja,ig}. \quad (2.3.7)$$

Call the contractions obtained from the two terms B and P , respectively. For each internal rail $u \in \{h, i, j\}$, cut its original gap and denote the resulting open diagrams by B_u, P_u . Closing that single cut in its inherited variance gives

$$\text{Cl}_u(B_u) = B, \quad \text{Cl}_u(P_u) = P.$$

Indeed, after testing the six external legs, the directed daughter is one of the cyclic matrix products

$$\begin{aligned} M_2[h, i] &= \sum_{f,c} F_f C_c C_{fh,ci}, & M_3[h, j] &= \sum_{b,e} B_b E_e C_{bh,je}, \\ P_4[i, j] &= \sum_{a,g} A_a G_g C_{ja,ig}, & B_4[i, j] &= \sum_{a,g} A_a G_g C_{ia,jg}. \end{aligned} \quad (2.3.8)$$

Here A, B, C, E, F, G are arbitrary boundary test vectors after the six external gap maps have been inserted. With X_h, X_i, X_j denoting the three internal gaps, the directed daughter is one of

$$M_2 X_i P_4 X_j M_3^T, \quad P_4 X_j M_3^T X_h^T M_2, \quad M_3^T X_h^T M_2 X_i P_4,$$

closed against X_h^T, X_i, X_j , respectively; cyclicity gives the same coordinate contraction in all three cases. The bent daughter is the raw cup-through-cap contraction

$$B = \text{Tr}(X_j M_3^T X_h^T M_2 X_i B_4).$$

Consequently the fully typed three-cut identity is

$$\boxed{H^{01} = \frac{1}{3} \sum_{u \in \{h,i,j\}} \text{Cl}_u(B_u - P_u).} \quad (2.3.9)$$

It has six coefficients of magnitude $1/3$ and residual degree three. Every residual gap occurs once; the root factor in Equation (2.3.5) supplies the tenth gap.

Each P_u is a three-gate full-wedge path. Each B_u is an acyclic raw cup-through-cap diagram. If $\hat{B}_u$ denotes the diagram with both bends normalized, direct substitution of $\hat{\eta} = n^{-1/2}\eta$ and $\hat{\varepsilon} = n^{-1/2}\varepsilon$ gives

$$B_u^{\text{raw}} = n\hat{B}_u. \quad (2.3.10)$$

The two bends are fragments of the same original circuit. Its single recorded base contributes exactly n^{-1} , so Equation (2.3.10) is dimension-free in the normalized complete circuit presentation. Further cuts merely subdivide that path and create no additional base factor.

The manuscript root token is $\mathbf{q}_C^{\text{hole}} = T_C T_C^*/2$, whereas the oriented root hit is

$$q_C^{01}(I) = -T_C T_C^* P_{\text{cyc}}^* = -2\mathbf{q}_C^{\text{hole}} P_{\text{cyc}}^*. \quad (2.3.11)$$

Pairing it with Equation (2.3.9) proves Equation (2.3.1): the six residual coefficients have mass two and Equation (2.3.11) doubles that mass to four.

For pass mode use the safe reshuffle

$$\tilde{Q}_{abe,cf g} := Q_{abc,ef g}, \quad \tilde{H}_{ABE,CFG} := H_{ABC,ef g}. \quad (2.3.12)$$

For an arbitrary opaque matrix Ξ the physical contraction is exactly

$$\sum_{e,f,g,E,F,G} \tilde{Q}_{abe,cf g} \Xi_{ef g,ef g} \tilde{H}_{ABE,CFG}; \quad (2.3.13)$$

this is the original coordinate sum after substituting Equation (2.3.12), not an ordinary product in the reshuffled cut. Testing all external and gap legs by matrix units proves Equation (2.3.9) with all twenty half-darts open. Applying the same joint attachment $\mathcal{A}_{rs}[\Xi]$ to its two sides proves Equation (2.3.2), even when the token lies in or across gap contexts. Its six coefficients have mass two, and the attachment occurs once.

We finally verify the advertised twenty-state coverage without an enumeration. Signed rail transport is the action Equations (2.2.6a) to (2.2.6e) on the two root gates, the complete bracket $B_u - P_u$, all ten gaps, their variances and formal adjoints, and the recorded circuit bases. Formula Equation (2.2.6c) follows simply by multiplying the rail increments $1, 2$ by α . Hence for $\alpha = 1, 4$ either zero or two wedge pairs reverse at every gate, while for $\alpha = 2, 3$ exactly one wedge pair reverses at every gate. Since there are five gates, their total skew sign is exactly Equation (2.2.6e). The translation β changes no local orientation.

For every ordered pair of distinct vertices (r, s) there is a unique $g_{\alpha,\beta}$ with $g(0) = r$ and $g(1) = s$, namely

$$\alpha = \kappa_5(s) - \kappa_5(r) \neq 0, \quad \beta = \kappa_5(r). \quad (2.3.14)$$

Thus the action is sharply two-transitive and the canonical computation has exactly twenty, rather than at most twenty, transported states. Because the same $g_{\alpha,\beta}$ transports every dart, Equation (2.2.6d) sends each of the ten unoriented core edges bijectively to one core edge and records its unique source–target variance. Transporting the tensor equality Equation (2.3.9), rather than its individual scalar closures, therefore proves the required identity in every state, with all twenty half-darts, all ten gaps, and their signs open. Principal-pivot and rail-exchange presentations are this same equality conjugated by the resulting fixed signed boundary isomorphisms. Circuit bases are merely transported along their original circuits, so their number is unchanged. Finally, the calculation in Equation (2.2.1) gives Equation (2.3.4). This proves every assertion. $\square$

Lemma 2.5 (Global attachment naturality). *If $T = T'$ are open typed tensors on $\mathcal{U}_{rs}$, then for every joint zero- or one-hole attachment,*

$$\mathcal{A}_{rs}[\Xi](T) = \mathcal{A}_{rs}[\Xi](T'). \quad (2.3.15)$$

Every complement tensor, every gap context, and the opaque token has the same multiplicity on the two sides. The assertion is stable under ancillary tensoring, signed transport, reversal, and full transpose.

Proof. Equality on $\mathcal{U}_{rs}$ is equality of every boundary coordinate. Multiply equal coordinates by the common complement coordinates and sum the same boundary indices. This proves Equation (2.3.15). The attachment is never expanded or copied, so all of its constituents occur once. Tensoring and the listed boundary isomorphisms preserve coordinate equality. $\square$

Definition 2.6 (Circuit-anchored presentation cost). Fix a circuit state τ . A component atom is a physical two-front atom, with all ancillary legs and circuit fragments left in their declared slots, presented as XY^* at its two correlated Hilbert–Schmidt fronts. Every other fixed letter is either a safe gate or fixed raw circuit wiring. The cost of an atom is

$$\|X\|_2 \|Y\|_2 \prod_G \|G\|_{\text{op}}, \quad (2.3.16)$$

where the product contains only safe gates and prescribed bounded maps, not the raw cups, caps, or bases belonging to the circuit wiring. Let $\mathcal{P}_\tau$ be the algebraic ℓ^1 -projective sum of tensors of such atoms and let $\|\cdot\|_{\mathcal{P}_\tau}$ be the corresponding projective cost. All named boundary variables remain multilinear slots.

If $\mathcal{R}_\tau^{\text{in}}, \mathcal{R}_\tau^{\text{out}}$ are retained root words, define $\|T\|_{\tau, \varepsilon}$ by inserting operator-norm-one matrix tests in every nonroot slot and taking the trace norm on the joint ordered root word. With no retained root, take absolute value after the terminal scalar closure.

Lemma 2.7 (Circuit-anchored atomic contraction). *For every $\zeta \in \mathcal{P}_\tau$,*

$$\left\| n^{-F(\tau)} \text{Contr}_{\Gamma, \tau}^{\text{raw}}(\zeta) \right\|_{\tau, \varepsilon} \leq \|\zeta\|_{\mathcal{P}_\tau}, \quad \text{Contr}_{\Gamma, \tau}^{\text{raw}} = n^{F(\tau)} \text{Contr}_{\Gamma, \tau}^{\text{norm}}. \quad (2.3.17)$$

The same estimate holds with arbitrary typed gaps and retained output ports. Inserting prescribed maps multiplies the right side by at most the product of their operator norms, provided they contain no hidden unnormalized cup or cap.

Proof. Take one atom from every component and insert all port tests and prescribed maps in their actual slots. Cut the resulting front graph only at the correlated Hilbert–Schmidt fronts. Along each path use

$$\|LZR\|_1 \leq \|L\|_{\text{op}} \|Z\|_1 \|R\|_{\text{op}};$$

along each cycle cut at two consecutive fronts and use

$$\|XY^*\|_1 \leq \|X\|_2 \|Y\|_2, \quad |\text{Tr } Z| \leq \|Z\|_1.$$

Signed rail maps and full transposes are isometries, so this removes every safe gate at precisely the cost in Equation (2.3.16).

Now cut each original circuit at the one base recorded by τ . Every other turn lies on an oriented path and straightens by the raw snake identity. The base closure is the sole identity trace on that circuit and contributes exactly n . If it has k fragments, the analytic estimate is still

$$\|A_1 \cdots A_k\|_1 \leq \|A_1\|_1 \prod_{j=2}^k \|A_j\|_{\text{op}},$$

so the identity cost is n , not n^k . This proves the exact second identity in Equation (2.3.17) and its atomic inequality. Expand projective presentations, apply the triangle inequality once to the complete sum, and take the projective infimum. Matrix-unit tests prove the retained-port statement without currying any coupled circuit. $\square$

Lemma 2.8 (Root-open operator-norm circuit closure). *Consider one normalized Kac ribbon atom after all nonroot legs have been closed in their original cyclic order, and retain the joint boundary on f_R distinct original $p \vee r$ circuits. If the atom has r nonconstant slots and total Euler defect g , its resulting root operator K satisfies*

$$\|K\|_{\text{op}} \leq A^r (r+1)^B n^{\max\{2-g-f_R, 0\}}. \quad (2.3.17a)$$

In particular the residual low cell $g = 1, f_R = 2$ has $\|K\|_{\text{op}} \leq A^r (r+1)^B$. This estimate is atomwise and is taken before any endpoint translation is summed.

Proof. Let $x \in \mathcal{R}^{\text{in}}$ and $y \in \mathcal{R}^{\text{out}}$ be unit vectors and insert the rank-one test $B = |x\rangle\langle y|$ at the complete joint root hole. Then $\langle y, Kx \rangle$ is the scalar contraction obtained by adding this single trace-class gate. Apply the atomic path/cycle proof of Lemma 2.7 to that scalar. The test has $\|B\|_1 = 1$ and is inserted only after the two root rails have been kept together, so neither a partial trace nor a norm of an individual cup is taken.

Of the F original $p \vee r$ circuits, precisely $F - f_R$ now close to identity traces. They contribute n^{F-f_R} . Multiplication by the once-normalized ribbon weight $\omega_{\Gamma} n^{-F}$ therefore gives exactly n^{-f_R} times the positive scalar presentation used in the closed Euler estimate. By Lemma 2.20, that presentation is at most $A^r (r+1)^B n^{2-g}$. Hence, uniformly in unit x, y ,

$$|\langle y, Kx \rangle| \leq A^r (r+1)^B n^{2-g-f_R}.$$

Taking the supremum proves the asserted estimate (with a favorable negative power relaxed to one). This rank-one test also shows directly that no exposed raw cup, extra circuit base, or replica factorization is being used. $\square$

2.4 One-resource module stability

The point of the module formulation is that arbitrary environments now follow by a formal induction rather than by a new analytic inequality.

Lemma 2.9 (Safe functoriality). *Let $T \in \mathfrak{M}_C(E)$. If L and R are amplified safe gates of total degree d , then*

$$\|LTR\|_{\mathfrak{M}_C(E)} \leq K^d \|T\|_{\mathfrak{M}_C(E)}. \quad (2.4.1)$$

If a new physical carrier is adjoined, then

$$\|T \otimes I_H\|_{\mathfrak{M}_C(E \otimes H)} \leq n \|T\|_{\mathfrak{M}_C(E)}. \quad (2.4.2)$$

Normalized Brauer cups, caps, permutations, and defect cuts do not increase the module norm.

Proof. The physical coefficient box has norm $\|C\|_{\text{op}} = K$. On the skew carrier, the correctly oriented Bianchi transfer is the same C -action by the first Bianchi identity; full-wedge, AA , and correctly oriented AB gates are composites of coefficient boxes with wedge inclusions and projections. Signed permutations, reversals, and rail maps are isometries. Normalized Brauer cups and caps have norm at most one, permutations are isometries, and antisymmetrizers are orthogonal projections. Submultiplicativity therefore gives $\|G\| \leq K^d$ for every expanded safe gate of degree d .

Apply these operations to an atomic representation of T . The operator-ideal inequality on the left and right factors gives Equation (2.4.1), and Equation (2.4.2) uses $\|I_H\|_1 = n$. The same generator calculation shows that the listed normalized boundary operations do not increase the atomic cost. Take the infimum over atomic representations. $\square$

This lemma is used for distance-hereditary folds and for the safe letters inside a circuit-anchored presentation. The raw compact K_5 bend is handled by the physical bridge in Section 2.3, not by safe functoriality.

Lemma 2.10 (Ported folds are free-module words). *Let $\mathcal{F}$ be an enterable distance-hereditary one-front fold with ordered formal ports $x_1, \dots, x_p$ on its carrier arcs. For an incoming two-front state Ξ , its open tensor has an identity*

$$\mathcal{F}(x_1, \dots, x_p)[\Xi] = \sum_{\nu} a_{\nu} u_{\nu}(x_1, \dots, x_p) \Xi v_{\nu}(x_1, \dots, x_p), \quad (2.4.3)$$

where every port variable occurs exactly once, every fixed letter in u_{ν}, v_{ν} is a safe gate or signed isometry, and $\sum_{\nu} |a_{\nu}| \leq A^s$ for a fold with s coefficient occurrences. After substituting safe words X_i for the ports,

$$\|\mathcal{F}(X_1, \dots, X_p)[\Xi]\|_1 \leq A^s K^s \left(\prod_i \|X_i\| \right) \|\Xi\|_1. \quad (2.4.4)$$

Proof. Induct along the enterable fold order. Ports before the first coefficient box concatenate to the appropriate left or right boundary word. Absorb the next box through a full physical wedge; after a signed boundary permutation, this is left or right composition by the physical C gate. Ports on the newly absorbed arcs concatenate in their inherited order. A Bianchi-mixed attachment first uses the exact skew transfer $\Phi_C = C$ and is then the same operation. No step contracts, duplicates, or sums over a port variable. The finite choice of entrance side and Bianchi daughter contributes at most a numerical constant per box. This proves Equation (2.4.3), and the trace-ideal inequality proves Equation (2.4.4). $\square$

Lemma 2.11 (Principal-nullity-one structure). *Let $\mathcal{C}$ be a connected ported two-carrier coefficient tangle whose compatible circuit maximum is at most two. After changing its one-circuit reference smoothing and applying signed share mutations, exactly one of the following holds.*

- (a) $\mathcal{C}$ is distance-hereditary and has an enterable elimination by isolated, pendant, true-twin, and false-twin folds.

(b) $\mathcal{C}$ has one of the three principal-pivot-equivalent five-vertex cores

$$P_C = \begin{pmatrix} 1 & 1 & 0 & 0 & 1 \\ 1 & 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 & 1 \\ 1 & 0 & 0 & 1 & 1 \end{pmatrix},$$

$$P_H = \begin{pmatrix} 1 & 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 1 & 1 \\ 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 & 1 \end{pmatrix}, \quad P_G = \begin{pmatrix} 1 & 1 & 0 & 0 & 1 \\ 1 & 0 & 1 & 0 & 1 \\ 0 & 1 & 0 & 1 & 1 \\ 0 & 0 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 \end{pmatrix}.$$

Every other vertex lies in one of five signed-twin fibers of that core or is a looped laminar ear. Twin fibers are chains of safe full-wedge digons; ears are enterable unary folds. In particular, after the safe pieces are collapsed, the only prime has an underlying presentation in Π_{core} . Choosing two ordered root fronts enhances it to a state $\sigma \in \Pi_{\text{root}}$; in pass mode, excising its five tagged gates gives the ordered split and finite boundary type of the joint attachment in Definition 2.3.

Proof. At a box let T be the forbidden transition and X, O the two compatible transitions. Starting from any compatible smoothing, two distinct circuits meeting at a box can be merged by at least one of X, O ; iterating produces a compatible Euler smoothing. Read its Euler circuit as a double-occurrence word and form the symmetric interlace matrix M over $\mathbb{F}_2$, with the diagonal recording which switch is orientation inconsistent. The extended Cohn–Lempel equality gives

$$\#\{\text{circuits after switching } S\} = 1 + \nu(M[S]).$$

Hence the hypothesis is exactly $\nu(M[S]) \leq 1$ for every principal submatrix.

We classify such matrices. Two zero-diagonal vertices must be adjacent, since otherwise their 2×2 principal block has nullity two. On an induced path of length four this restriction leaves, up to reflection, six possible diagonal patterns; the following witnesses have nullity two:

$\{i : M_{ii} = 0\}$	S
$\emptyset$	$\{0, 1, 3, 4\}$
$\{0\}$	$\{0, 2, 3\}$
$\{1\}$	$\{1, 3, 4\}$
$\{2\}$	$\{0, 1, 3, 4\}$
$\{0, 1\}$	$\{0, 2, 3\}$
$\{1, 2\}$	$\{1, 3, 4\}$

so the off-diagonal graph is induced- P_5 -free. The forbidden-induced- subgraph characterization of distance-hereditary graphs [BM86; Tra09] now says that a non-distance-hereditary graph contains a C_5 , house, or gem: longer holes and the domino contain an induced P_5 . Direct binary row reduction of the 2^5 diagonal assignments leaves exactly one assignment for each of these three graphs, namely P_C, P_H, P_G above.

It remains to classify extensions of a prime P . If $b \in \mathbb{F}_2^5$ is the boundary column of a new vertex and η its diagonal, checking the $2^5 \cdot 2$ possibilities against principal nullity at most one leaves only

$$(b, \eta) = (0, 1) \quad \text{or} \quad (b, \eta) = (Pe_i + e_i, P_{ii}), \quad 0 \leq i < 5.$$

For two new vertices of types i, j , the same condition uniquely forces $M_{xy} = P_{ij} + \delta_{ij}$; a type- ∞ vertex is looped and isolated. Thus all extensions are five twin fibers plus looped isolated vertices. Principal pivot sends P_C to P_H , and a two-vertex pivot sends P_G to P_C , so they are three presentations of one prime.

The mutation theorem for chord diagrams with the same intersection graph realizes these matrix reductions by share mutations. On the four open boundary wires a mutation is a signed permutation or full transpose and therefore a Hilbert–Schmidt isometry. Inspecting the two possible signed twin words shows a complete physical wedge digon; a looped isolated chord is an enterable laminar unary fold. These are precisely the safe gates in [Lemma 2.10](#). This proves both alternatives. $\square$

Lemma 2.12 (Circuit-anchored single-prime rewrite). *Fix a declared circuit state τ and one tagged unresolved K_5 core. In pass mode its complete complement and the existing token form one joint attachment. Replacing the core by [Equation \(2.3.2\)](#) produces six raw circuit-anchored presentations of total coefficient mass at most two, with the token, every gap context, and all five coefficient gates occurring once. In root mode [Equation \(2.3.1\)](#) creates the unique token, has total coefficient mass at most four, and leaves exactly three residual coefficient gates. No circuit base is normalized during either rewrite.*

Proof. Excise the five gates. Everything else is their joint attachment, so [Theorem 2.4](#) applies without opening or realigning the token. Its budgets, degrees, and once-only assertions give the result. The identities are raw; [Equation \(2.2.8\)](#) is applied only after all selected cores on an original circuit have been assembled. $\square$

Lemma 2.13 (Actual-pass serialization). *Before the first rewrite, declare the maximizing smoothing, the pair-support groups, all component and output ports, the original circuits and their bases, and one prime tag on every non-distance-hereditary group. If r tagged cores remain, repeated circuit-anchored rewrites produce a finite sum of one-token presentations with coefficient mass at most*

$$2^r \leq 2^{m/5}. \quad (2.4.5)$$

Every original coefficient occurrence and gap context occurs once in every term, the token occurs once, and the number of unresolved tags drops at each step. The conclusion includes nested cores and a token lying in or jointly across gap contexts.

Proof. The maximizing smoothing partitions coefficient occurrences into pair-support groups, so their occurrence sets are disjoint. [Lemma 2.11](#) assigns at most one prime tag to each non-distance-hereditary group. These tags are fixed; the classification is not rerun on a daughter.

Choose one unresolved tag and remove its five gates. By disjointness the existing token lies in the complement, never in a selected gate. If it lies in one gap, retain the token-free pieces in their inherited order as the two sides of one one-hole context. If it meets several gap slots, retain the whole complement as the joint attachment. [Lemma 2.5](#) and [Equation \(2.3.2\)](#) then keep that attachment once in all six daughters. Signed share mutations transport every marked port and circuit base together and never cut the token.

The selected tag is removed and no daughter creates a new tag. Induction on r gives mass 2^r and all once-only assertions. The tagged cores are disjoint five-element subsets of the original occurrences, so $5r \leq m$. $\square$

Theorem 2.14 (Circuit-anchored all-order comb lift). *Fix a declared global circuit state τ . Let $\mathcal{T}$ be any finite composition of correlated same- C distance-hereditary pair combs, signed- K_5 pair combs, safe word decorations, and normalized boundary operations. Let m be the total C -degree of*

$\mathcal{T}q_C$, counting the incoming token as its two underlying coefficient occurrences. If no new smoothing carrier is adjoined, the raw tangle has a finite circuit-anchored presentation $\zeta_{\mathcal{T},\tau} \in \mathcal{P}_\tau$ satisfying

$$\|\zeta_{\mathcal{T},\tau}\|_{\mathcal{P}_\tau} \leq C_0^m K^{m-2} h^2, \quad (2.4.6)$$

where C_0 is numerical and independent of n , m , and E . In particular, arbitrarily many interleaved combs still pay h^2 only once. Every declared port and context occurs once, token degree remains one, and the circuit-base normalization is deferred to [Equation \(2.3.17\)](#). No ordinary open-component norm is asserted for a K_5 branch.

Proof. We prove the following induction invariant. After any number of comb rewrites, each daughter is a finite raw, one-token, circuit-anchored presentation in the same declared state τ ; every original coefficient gate, gap context, and circuit base occurs once; the token degree is one; and the total coefficient mass of all daughters is at most 2^s , where s is the number of pass rewrites already made.

For zero combs the invariant is the atomic paired-front presentation of [Definition 2.2](#). Distance-hereditary shares are eliminated by the two-anchor fold grammar of [Lemma 2.10](#); this only inserts safe gates and hence preserves the invariant. In a non-distance-hereditary comb, [Lemma 2.11](#) supplies one fixed signed- K_5 tag, while its twin fibres and laminar ears are safe. Excise the five tagged gates. The entire remainder—including every unprocessed comb, gap, and the one token—is the single joint attachment in [Lemma 2.12](#). Its six pass daughters have total mass at most twice that of the parent, retain the attachment once, remove the chosen tag, and create no tag. This is the induction step. The fixed tags are disjoint five-element subsets of the original occurrences, so $5s \leq m$; hence $2^s \leq C_0^m$.

After every tag has been removed, restore the safe fibres, ears, folds, and boundary operations. Their degrees add to $m - 2$, so [Lemma 2.9](#) contributes at most K^{m-2} times a numerical exponential in m . The root is initialized exactly once and never duplicated; its two Hilbert–Schmidt fronts cost $\text{Tr } Q_2 = 2h^2$. No smoothing carrier was adjoined, and all compact bends remain in the raw wiring, so no circuit base or factor n has been spent. Absorbing the harmless factor two in C_0 proves [Equation \(2.4.6\)](#) and all assertions of the invariant. $\square$

2.5 Maximum smoothing and the positive-credit induction

We recall the topological input in the form needed for the analytic induction. For a connected coefficient packet P , let $c^*(P)$ be the maximum number of alternating carrier circuits over all compatible local smoothings. Fix a maximizing smoothing with circuits $\gamma_1, \dots, \gamma_c$, where $c = c^*(P)$. The support of a box is the singleton or unordered pair of carriers containing its two smoothing passages. The graph of two-carrier supports is connected.

For a fixed pair $\{i, j\}$, freeze every box outside that support group at the maximizing smoothing. The remaining ported two-carrier tangle has compatible circuit maximum at most two: otherwise replacing the $\gamma_i \cup \gamma_j$ subsystem by a smoothing with $r > 2$ circuits would create $c - 2 + r > c$ global circuits. Hence every group is a ported principal-nullity-one comb and is controlled by [Theorem 2.14](#).

Lemma 2.15 (Two carriers in every nonzero nonsingleton packet). *Let P be a connected q -packet with at least two coefficient boxes. If it has no forbidden within-wedge q -pair, then some allowed smoothing h_P satisfies*

$$\#(q_P \vee h_P) \geq 2.$$

Consequently every nonzero centered packet with $m_P \geq 2$ has $c^(P) \geq 2$.*

Proof. Start with any allowed smoothing. If it already has two circuits, there is nothing to prove. Otherwise $q_P \vee h_P$ is one Euler circuit and each box is visited twice. If toggling the direct/cross

choice at one box splits the circuit, we are done. If no single toggle splits, the two visits at every box have opposite directions; otherwise the alternative L – R reconnection closes the two complementary arcs separately.

Join the two visits of each box by a chord of the Euler circle. Two interlacing chords are split by toggling their two boxes (the associated 2×2 interlace matrix acquires a one-dimensional kernel). If no two chords interlace, the chord diagram is laminar. An innermost chord then has consecutive endpoints, and the q -edge between those visits joins the two L -darts or the two R -darts of that box, a forbidden within-wedge pair. This contradiction proves the first assertion. A forbidden pair is zero by skewness, while a surviving one-box packet is a scalar trace and is killed by centering. The second assertion follows. $\square$

Theorem 2.16 (Positive-credit circuit-anchored component presentation). *Fix a global circuit state τ before any component is estimated. Let P be a connected pure-Weyl component with $m_P \geq 2$ coefficient occurrences and $c = c^*(P) \geq 2$, and retain all of its declared cut interfaces. Its raw physical open tensor has a finite presentation $\zeta_{P,\tau} \in \mathcal{P}_\tau$ satisfying*

$$\|\zeta_{P,\tau}\|_{\mathcal{P}_\tau} \leq C_1^{m_P} n^{c-2} \|C_P\|_2^2 \|C_P\|_{\text{op}}^{m_P-2}. \quad (2.5.1)$$

Every predeclared interface occurs once. This is a projective presentation cost. It is not an ordinary norm of a separately evaluated open K_5 component; evaluation and circuit normalization are deferred to Lemma 2.7.

Proof. Choose a spanning tree of the connected support graph. There are three initialization cases. If some pair group contains a K_5 prime, root at that group, choose its ordered output fronts and hence $\sigma \in \Pi_{\text{root}}$, and use the raw identity Equation (2.3.1). Its once-only conditions and Equation (2.3.3) activate two carriers and create the unique open F_C paired-front resource. Equations (2.2.1) to (2.2.5) bound its front cost by $\text{Tr } Q_2 = 2\|C_P\|_2^2$, giving $2A_{\text{root}} h^2 K^{s-2}$ before the remaining numerical local constants are absorbed into C^s . No cap is used to identify an open morphism. If there is no K_5 prime but some distance-hereditary group has at least two boxes, root there and use its ordinary two-front fold; all later groups are also distance-hereditary, so the usual trace-class ideal suffices. Finally, if every pair group is a singleton, choose two adjacent tree edges. Their two coefficient gates occupy three carriers and supply the two initial Hilbert–Schmidt fronts.

In the first two cases the initial two-front resource, before charging the remaining gates and local coefficients, costs at most a numerical multiple of $\|C_P\|_2^2$; in the singleton case the corresponding resource costs at most $n\|C_P\|_2^2$. The analytic inequalities behind these statements are respectively

$$\|XY\|_1 \leq \|X\|_2 \|Y\|_2 \quad \text{and} \quad \|(X \otimes I)(I \otimes Y)\|_1 \leq n\|X\|_2 \|Y\|_2. \quad (2.5.2)$$

Traverse the spanning tree away from the root. Each genuinely new smoothing carrier inserts one trace-class identity and costs exactly n . Insert the corresponding pair comb into its correlated holes using Theorem 2.14. Lemma 2.13 serializes every nested prime; distance-hereditary groups use their ordinary two-front atoms, embedded in the same projective presentation. This changes only the $C_1^{m_P} \|C_P\|_{\text{op}}^{m_P-2}$ factor and preserves the single paired-front resource. Every non-tree group uses carriers already present and costs no additional n . All raw circuit fragments and their bases remain recorded in τ .

In the two-carrier root case there are $c - 2$ new carriers. In the three-carrier root case the initial n in Equation (2.5.2) is followed by $c - 3$ new carriers. Both give exactly n^{c-2} . Applying no circuit normalization and taking no open-component norm proves Equation (2.5.1) as a presentation estimate. $\square$

2.6 Colors, defects, marked roots, and the packet/ribbon estimate

We now distinguish two harmless global normalizations on W . The algebraic eigenframe in [Sections 2.1 to 2.5](#) uses $\langle A, B \rangle_\wedge = -\frac{1}{2} \text{Tr}(AB)$, so its unit coordinate bivector is $E_{ij} - E_{ji}$ and the exact algebraic cap has $\text{Tr } Q_D = 2 \text{rank } D$. The probabilistic/Malliavin calculation uses the Frobenius metric $\langle A, B \rangle_{\text{HS}} = -\text{Tr}(AB)$ and the orthonormal Kac frame

$$a_{ij} = \frac{E_{ij} - E_{ji}}{\sqrt{2}}. \quad (2.6.0)$$

Multiplying the ambient inner product by a scalar does not change an orthogonal projector onto a line, an operator adjoint, or any conjugation map. Thus all packet identities pass unchanged; only vector representatives and the displayed Casimir constants must use the stated frame. From this section onward z and a_i are HS-unit unless an algebraic active eigenvector A_a is named explicitly. In particular $-2z_{\text{mat}}^2$ is exactly the projector onto the coordinate plane.

Let a block color be

$$C_j = B_j - \mu_j I_W, \quad 0 \preceq B_j \preceq K I_W, \quad \mu_j = N^{-1} \text{Tr } B_j. \quad (2.6.1)$$

Lemma 2.17 (Kac Weyl decomposition and the marked Ricci identity). *Let P_z be the rank-one projector onto a unit decomposable bivector and $C_z = P_z - N^{-1} I_W$. If $L_z = -2z_{\text{mat}}^2$ is the orthogonal projector onto the plane of z , put*

$$B_z = \frac{L_z - (2/n) I_H}{n - 2}. \quad (2.6.2)$$

Then the orthogonal curvature decomposition is

$$C_z = W_z + X_{B_z}, \quad W_z \in \mathcal{W}, \quad X_B(u \wedge v) = Bu \wedge v + u \wedge Bv, \quad (2.6.3)$$

with no scalar or $\Lambda^4 H$ summand, and

$$\|B_z\|_{\text{op}} = \frac{1}{n}, \quad \|B_z\|_F^2 = \frac{2}{n(n-2)}, \quad \|X_B\|_2^2 = (n-2)\|B\|_F^2, \quad \|X_B\|_{\text{op}} \leq 2\|B\|_{\text{op}}. \quad (2.6.4)$$

In the cumulative moving frame the transported B_z are reverse martingale differences. If $\mathfrak{B}_r$ is their sum, then

$$\mathbb{E}\|\mathfrak{B}_r\|_F^2 = \frac{2r}{n(n-2)}, \quad \mathbb{E}\|\mathfrak{B}_r\|_F^4 \leq C \frac{r^2}{n^2(n-2)^2}, \quad (2.6.5)$$

and, for fixed c , uniformly over $r \leq cN$,

$$\max_{r \leq cN} \|\mathfrak{B}_r\|_{\text{op}} = o_{\mathbb{P}}(1). \quad (2.6.6)$$

More quantitatively, for every fixed L and $1 \leq p \leq L \log n$,

$$\left(\mathbb{E} \max_{r \leq cN} \|\mathfrak{B}_r\|_{\text{op}}^p \right)^{1/p} \leq C_{c,L} \left(\sqrt{\frac{p + \log n}{n}} + \frac{p + \log n}{n} \right) = o(1). \quad (2.6.6a)$$

The same martingale satisfies the dimension-free Frobenius estimate

$$\left(\mathbb{E} \max_{r \leq cN} \|\mathfrak{B}_r\|_F^p \right)^{1/p} \leq C_c \sqrt{p}, \quad 2 \leq p \leq L \log n. \quad (2.6.6b)$$

Finally let $A_i = \text{ad}_{a_i}$. For traceless symmetric B on H and arbitrary $Z \in \text{End}(W)$, define

$$\begin{aligned}\mathcal{R}_i(B, Z) &= \{X_B, [A_i, Z]\}a_i, & U_i(B, Z) &= \{X_B, Z\}a_i, \\ E_i(B, Z) &= \{X_{[a_i, B]}, Z\}a_i, & \bar{E}(B, Z) &= N^{-1} \sum_i E_i(B, Z).\end{aligned}\tag{2.6.7}$$

Put $\Psi(T) = \sum_j A_j T a_j$ and $H_B(Z) = -N^{-1} X_B \Psi(Z)$. Then the exact two-gradient decomposition is

$$\boxed{\mathcal{R}_i = A_i U_i - (E_i - \bar{E}) - \frac{1}{N} \sum_j A_j V_j - H_B(Z)}, \quad \bar{E} = \frac{1}{N} \sum_j A_j V_j + H_B(Z), \quad V_j = X_B Z a_j,\tag{2.6.8}$$

and its star-normalized square functions obey

$$\begin{aligned}\frac{1}{N} \sum_i \|U_i\|^2 &\leq \frac{4(n-2)}{N} \|B\|_F^2 \|Z\|_{\text{op}}^2, \\ \frac{1}{N} \sum_j \|V_j\|^2 &\leq \frac{n-2}{N} \|B\|_F^2 \|Z\|_{\text{op}}^2, \\ \frac{1}{N} \sum_i \|E_i - \bar{E}\|^2 &\leq \frac{64}{n-1} \|B\|_F^2 \|Z\|_{\text{op}}^2.\end{aligned}\tag{2.6.9}$$

The residue obeys

$$\|H_B(Z)\| \leq \frac{2\sqrt{n-2}}{N} \|B\|_{\text{op}} \|Z\|_2.\tag{2.6.9a}$$

Proof. Tracing one skew pair in C_z gives B_z in Equation (2.6.2). A decomposable bivector has vanishing alternating four-form projection, which proves Equation (2.6.3). Diagonalizing B gives

$$\sum_{i < j} (b_i + b_j)^2 = (n-2) \sum_i b_i^2$$

when $\sum_i b_i = 0$; this proves the Hilbert–Schmidt identity in Equation (2.6.4), and the other norm statements follow directly from the two eigenvalues of B_z and $\|X_B\|_{\text{op}} \leq 2\|B\|_{\text{op}}$.

Reverse exposure makes the conditional mean of each transported B_z zero, while isotropy gives

$$\mathbb{E}(B_z^2 \mid \mathcal{F}_{\text{future}}) = \frac{2}{n^2(n-2)} I_H.$$

Martingale orthogonality proves the second moment in Equation (2.6.5); expanding four factors and pairing the two nonvanishing conditional contractions proves the fourth-moment bound. Matrix Freedman’s inequality [Tro11], with increment bound $1/n$ and quadratic variation $O(1/n)$ up to time cN , gives

$$\Pr \left\{ \max_{r \leq cN} \|\mathfrak{B}_r\|_{\text{op}} > C_c \frac{\log n}{\sqrt{n}} \right\} \leq e^{-c_c (\log n)^2},$$

which proves Equation (2.6.6) with enough probability for every logarithmic-degree expansion. Integrating the same stopped tail inequality gives Equation (2.6.6a). On the exceptional tail one uses the deterministic estimate $\max_{r \leq cN} \|\mathfrak{B}_r\|_{\text{op}} \leq cN/n$; its polynomial size is absorbed by $e^{-c_c (\log n)^2}$ for $p \leq L \log n$. For Equation (2.6.6b), apply the Hilbert-space Burkholder inequality in $\text{Sym}_0(H)$. The predictable Frobenius square function is bounded by a numerical constant because

$$\sum_{s \leq cN} \mathbb{E}_{s+1} \|B_{v_s}\|_F^2 = \frac{2\lfloor cN \rfloor}{n(n-2)} = O_c(1),$$

and $\max_s \|B_{v_s}\|_F = \sqrt{2/[n(n-2)]}$. Hence

$$\left(\mathbb{E} \max_{r \leq cN} \|\mathfrak{B}_r\|_F^p\right)^{1/p} \leq C \left(\sqrt{p} + p\sqrt{\frac{2}{n(n-2)}}\right) \leq C_c \sqrt{p}$$

in the displayed range. This proves [Equation \(2.6.6b\)](#).

The product rule $[A_i, X_B] = X_{[a_i, B]}$ gives $\mathcal{R}_i = A_i U_i - E_i$. Direct expansion of the plane mean gives

$$\bar{E} = \frac{1}{N} \{\Psi(X_B Z) - X_B \Psi(Z) + Z \Psi(X_B)\}.$$

Unimodularity gives $\Psi(X_B) = 0$ because X_B is symmetric, while the first term is $N^{-1} \sum_j A_j V_j$. This proves [Equation \(2.6.8\)](#). The adjoint Casimir calculation $\|\Psi(T)\| \leq \sqrt{n-2} \|T\|_2$, together with $\|X_B\|_{\text{op}} \leq 2\|B\|_{\text{op}}$, proves [Equation \(2.6.9a\)](#). Finally use

$$\sum_i \|A_i x\|^2 = (n-2) \|x\|^2$$

for the first two square functions. Indeed, $\sum_i \|U_i\|^2 = \|\{X_B, Z\}\|_2^2 \leq 4\|X_B\|_2^2 \|Z\|_{\text{op}}^2$ and $\sum_j \|V_j\|^2 = \|X_B Z\|_2^2 \leq \|X_B\|_2^2 \|Z\|_{\text{op}}^2$; now use [Equation \(2.6.4\)](#).

For the centered E term one must not diagonalize B while pretending that the fixed coordinate-plane frame is unchanged. Instead use the basis-free Casimir identity

$$\sum_i \|[a_i, B]\|_F^2 = n\|B\|_F^2. \quad (2.6.9b)$$

Indeed, the left side uses a complete orthonormal basis of $\mathfrak{so}(n)$ and is basis invariant; it may therefore be evaluated after diagonalizing B , where it is $\sum_{r < s} (b_r - b_s)^2 = n \sum_r b_r^2$ because $\text{Tr } B = 0$. Since $\|X_S\|_{\text{op}} \leq 2\|S\|_F$,

$$\|E_i(B, Z)\| \leq 4\|[a_i, B]\|_F \|Z\|_{\text{op}}.$$

Subtracting the plane-index mean is an orthogonal projection. Hence

$$\frac{1}{N} \sum_i \|E_i - \bar{E}\|^2 \leq \frac{16n}{N} \|B\|_F^2 \|Z\|_{\text{op}}^2 = \frac{32}{n-1} \|B\|_F^2 \|Z\|_{\text{op}}^2,$$

which is stronger than the stated $64/(n-1)$ bound. This calculation uses the complete plane sum; no individual marked position is estimated. $\square$

Put

$$\beta_j^2 := N^{-1} \|W_j\|_2^2 \leq \alpha_j^2 := N^{-1} \text{Tr}(C_j^2) \leq K\mu_j. \quad (2.6.10)$$

The first inequality is orthogonality of the pure-Weyl projection; the second follows from $B_j^2 \preceq KB_j$ and centering. Packet estimates use β_j , while α_j records the variance before the Ricci component is removed.

Definition 2.18 (Rooted ribbon record and raw coefficient). A rooted orthogonal-Weingarten ribbon record Γ of degree m consists of the dart set $\{1, \dots, m\} \times \{L_0, L_1, R_0, R_1\}$; its coefficient boxes and color labels; perfect matchings p, q, r giving the row, coefficient-column, and rooted trace wiring; all local wedge and direct/cross signs; the distinguished root boundary; and the following orthogonal-Weingarten path data.

Partition the darts into the independent Haar-color classes $\mathcal{H}_\Gamma$ (there is one class when all entries come from one Haar matrix). If $a \in \mathcal{H}_\Gamma$ contains m_a coefficient occurrences, put $k_a = 2m_a$,

so it has $2k_a = 4m_a$ darts, and denote by p_a, q_a the restrictions of p, q . The rooted dart order chooses a deterministic relabelling σ_a with $\sigma_a \cdot e_{k_a} = p_a$, where

$$e_k = \{\{1, 2\}, \{3, 4\}, \dots, \{2k-1, 2k\}\},$$

and put $\mathbf{m}_a = \sigma_a^{-1} \cdot q_a \in \mathcal{P}_2(2k_a)$. The orthogonal Weingarten graph has vertex set $\bigsqcup_{j \geq 0} \mathcal{P}_2(2j)$. At level j it has a solid arrow

$$\mathbf{n} \longrightarrow (i, 2j-1) \cdot \mathbf{n}, \quad 1 \leq i < 2j-1,$$

and it has a dashed arrow $\mathbf{n} \dashrightarrow \mathbf{n}^\downarrow$ when $\{2j-1, 2j\} \in \mathbf{n}$, where $\mathbf{n}^\downarrow$ is obtained by deleting that pair. The record chooses, for every a , one *individual* path γ_a from $\mathbf{m}_a$ to the empty pairing. It uses exactly k_a dashed arrows; write s_a for its number of solid arrows. If μ_a is the coset type of $\mathbf{m}_a$, define

$$d_a := |\mathbf{m}_a| = k_a - \ell(\mu_a) = k_a - \#(p_a \vee q_a), \quad g_a := s_a - d_a \geq 0, \quad g_{\text{Wg}} := \sum_{a \in \mathcal{H}_\Gamma} g_a.$$

Thus g_{Wg} is the total number of solid arrows beyond shortest orthogonal-Weingarten paths; unlike the unitary excess, it need not be even.

Let w_Γ be the scalar coefficient of this chosen product path, including the bounded local wedge and direct/cross coefficients. Write $\text{Contr}_{\Gamma, \tau}^{\text{raw}}$ for the typed raw contraction determined by the record and its circuit state, and put

$$F_\Gamma = \#(p \vee r), \quad \ell_\Gamma = \#(p \vee q) = \sum_{a \in \mathcal{H}_\Gamma} \#(p_a \vee q_a), \quad \omega_\Gamma = n^{F_\Gamma} w_\Gamma, \quad \mathcal{A}_\Gamma = w_\Gamma \text{Contr}_{\Gamma, \tau}^{\text{raw}}.$$

Thus w_Γ is a scalar coefficient, not the value or norm of an open contraction; ω_Γ merely expresses the same coefficient in once-normalized original-circuit units.

Lemma 2.19 (Orthogonal-Weingarten path coefficient and stable range). *For the record of Definition 2.18, every chosen product path satisfies*

$$|w_\Gamma| \leq C^m n^{-4m + \ell_\Gamma - g_{\text{Wg}}}. \quad (2.6.10a)$$

For fixed endpoint pairings and fixed $g \geq 0$, the number of product paths with $g_{\text{Wg}} = g$ is at most

$$C^m (m+1)^{Cg}. \quad (2.6.10b)$$

If $n > 12(2m)^{7/2}$, the product path expansion is absolutely convergent. In particular, for every fixed L , this applies uniformly to $m \leq L \log n$ once $n \geq n_0(L)$.

Proof. Fix one Haar-color class a . By orthogonal invariance, $\text{Wg}_n^{\text{O}}(p_a, q_a) = \text{Wg}_n^{\text{O}}(e_{k_a}, \mathbf{m}_a)$. The alternating components of $p_a \vee q_a$ have half-lengths equal to the parts of the coset type μ_a ; hence $|\mathbf{m}_a| = k_a - \#(p_a \vee q_a) = d_a$.

The orthogonal path expansion of Collins–Matsumoto [CM17, Theorem 4.6] is

$$(-1)^{d_a} n^{k_a + d_a} \text{Wg}_n^{\text{O}}(e_{k_a}, \mathbf{m}_a) = \sum_{h \geq 0} \# \mathcal{P}(\mathbf{m}_a, d_a + h) (-n)^{-h}.$$

Consequently one individual path of excess g_a has coefficient $(-1)^{d_a + g_a} n^{-k_a - d_a - g_a}$. Multiplying over the independent Haar colors and restoring the bounded local wedge/direct-cross coefficients gives

$$|w_\Gamma| \leq C^m n^{-\sum_a (k_a + d_a + g_a)} = C^m n^{-2m - (2m - \ell_\Gamma) - g_{\text{Wg}}},$$

which is Equation (2.6.10a).

For the count, Collins–Matsumoto [CM17, Theorem 4.9 and Lemma 4.12] give

$$\#\mathcal{P}(\mathbf{m}_a, d_a + g_a) \leq (12k_a^{7/2})^{g_a} \#\mathcal{P}(\mathbf{m}_a, d_a), \quad \#\mathcal{P}(\mathbf{m}_a, d_a) = \prod_i \text{Cat}_{\mu_{a,i}-1} \leq 4^{d_a}.$$

Let $h = |\mathcal{K}_\Gamma| \leq m$. Summing over the weak compositions $g = \sum_a g_a$ and using $k_a \leq 2m$ and $\sum_a k_a = 2m$ yields

$$\#\{\gamma : g_{\text{Wg}} = g\} \leq 4^{2m} \binom{g+h-1}{h-1} (12(2m)^{7/2})^g \leq C^m (m+1)^{Cg}.$$

The last form is deliberately uniform at $m = 1$. Finally, $\sum_{g_a \geq 0} (12k_a^{7/2}/n)^{g_a}$ converges for every a when $n > 12(2m)^{7/2}$. The formal path series is the Laurent expansion at infinity of the rational orthogonal Weingarten function, so within this convergence range it equals that function; equivalently, this is the analytic stable range in [CM17, Theorem 4.11]. Since $(\log n)^{7/2} = o(n)$, the logarithmic-degree assertion follows. $\square$

A rooted orthogonal-Weingarten ribbon map Γ has m coefficient occurrences and defect g . If color j occurs m_j times in q_j pairing-connected components, one normalized closure is assigned to each color component, while one global root trace remains unnormalized. Thus after the forbidden-pair zero terms have been discarded, Lemma 2.15 verifies $c^*(P) \geq 2$ at this use and Theorem 2.16 contributes

$$\beta_j^2 K^{r-2} \leq \mu_j K^{r-1} \quad (2.6.11)$$

for a component of size $r \geq 2$, rather than an additional factor N . The unique global trace supplies the leading N .

Lemma 2.20 (Pairing defect and fixed-defect enumeration). *On the $4m$ ribbon darts let p, q, r be respectively the row, coefficient, and trace pairings, and put*

$$d(a, b) := 2m - \#(a \vee b), \quad F = 2m - d(p, r), \quad \ell_\Gamma = 2m - d(p, q).$$

Let P range over the connected coefficient packets. There is an integer $g_{\text{top}} \geq 0$ such that

$$F - 4m + \ell_\Gamma + \sum_P c^*(P) = 2 - g_{\text{top}}. \quad (2.6.12)$$

If the orthogonal-Weingarten path has excess g_{Wg} , put $g = g_{\text{top}} + g_{\text{Wg}}$. Its contribution has an additional factor $n^{-g_{\text{Wg}}}$, and every defect cut acts contractively on the retained two-front boundary. For fixed local vertex types,

$$\#\{\Gamma : |\Gamma| = m, g(\Gamma) = g\} \leq C^m (m+1)^{Cg}. \quad (2.6.13)$$

Proof. For every packet P , choose a compatible smoothing h_P attaining $c^*(P)$ and put $h = \bigsqcup_P h_P$. By construction

$$\#(q \vee h) = \sum_P c^*(P).$$

The direct/cross compatibility at every coefficient vertex leaves at most the two distinguished rooted trace circuits, so $\#(h \vee r) \leq 2$. The reverse triangle inequality for the pairing distance therefore gives

$$\begin{aligned} d(q, r) &\geq d(h, r) - d(q, h) \\ &= \#(q \vee h) - \#(h \vee r) \geq \sum_P c^*(P) - 2. \end{aligned}$$

The ordinary triangle inequality gives $d(p, r) + d(p, q) \geq d(q, r)$. Substituting the definitions of F, ℓ_Γ yields

$$F - 4m + \ell_\Gamma + \sum_P c^*(P) \leq 2.$$

All quantities are integers, so Equation (2.6.12) defines $g_{\text{top}} \geq 0$ exactly. For the enumeration, use the ribbon data rather than a projection in the pairing metric. Replace every coefficient vertex by its maximizing smoothing and retain its four matching colours. Concretely, there is one zero-cell for each of the $4m$ darts and one one-cell for each pair in each of p, q, h, r , hence

$$|V| = 4m, \quad |E| = 4(2m) = 8m.$$

At a direct vertex the four colours occur cyclically as p, q, h, r ; a crossed vertex reverses the relevant two half-edges. Attach discs along the three alternating circuit families $p \vee r$, $p \vee q$, and $q \vee h$. The remaining $h \vee r$ circuits are boundary components; write $b = \#(h \vee r) \leq 2$. The local direct/cross table makes every interior link a circle and every boundary link an interval. After separating a pinched boundary identification by the corresponding normalized defect cut, this is a connected cellular surface $\mathcal{S}_\Gamma$ with boundary (possibly nonorientable) and

$$\chi(\mathcal{S}_\Gamma) = F + \ell_\Gamma + \sum_P c^*(P) - 4m = 2 - g_{\text{top}}. \quad (2.6.12a)$$

This also gives a topological proof of the nonnegativity already obtained from the pairing triangle inequality.

For the enumeration only, cap its b rooted boundary components by labelled discs, obtaining a closed surface $\bar{\mathcal{S}}_\Gamma$ with

$$\chi(\bar{\mathcal{S}}_\Gamma) = \chi(\mathcal{S}_\Gamma) + b, \quad g_h := 2 - \chi(\bar{\mathcal{S}}_\Gamma) = g_{\text{top}} - b \geq 0.$$

Choose a spanning tree T of the cellular one-skeleton and a spanning tree T^* of the dual of the capped surface which is disjoint from T . Euler's formula gives

$$|E \setminus (T \cup T^*)| = |E| - (|V| - 1) - (|F| + b - 1) = 2 - \chi - b = g_h.$$

Cut the two sides of these g_h exceptional edges and remember the $b \leq 2$ labelled cap discs. The result is a rooted planar bounded-local-type map with marked boundary corners. A depth-first traversal gives such planar maps a balanced-parenthesis encoding with finitely many local symbols, hence at most C^m choices. To reconstruct $\mathcal{S}_\Gamma$, choose and pair the $O(g_h)$ marked corners and choose one twist bit for each gluing; this contributes at most $m^{C_{g_{\text{top}}}}$. Pinch normalizations are encoded by the same marked-corner data and their Brauer maps are contractive by Lemma 2.9.

By Lemma 2.19, an individual orthogonal-Weingarten path of excess g_{Wg} contributes $n^{-g_{\text{Wg}}}$ beyond its shortest-path power and has at most $C^m(m+1)^{C_{g_{\text{Wg}}}}$ choices. Cutting one pairing switch is a normalized Brauer cup, cap, permutation, or antisymmetrizer; Lemma 2.9 makes it contractive on the open fronts. Convolving the topological and path excesses costs at most $g+1$, which is absorbed by $(m+1)^{C_g}$. Thus a total defect g has at most $C^m(m+1)^{C_g}$ realizations, proving Equation (2.6.13). $\square$

Combining Equation (2.6.12) with the Weingarten excess gives

$$n^F n^{-4m+\ell_\Gamma} \prod_P n^{c^*(P)} n^{-g_{\text{Wg}}} = n^{2-g} \asymp N n^{-g}.$$

After $h_j^2 \leq NK\mu_j$ is inserted in Equation (2.5.1), a packet of color j and size r contributes

$$C^r n^{c^*(P)} \mu_j K^{r-1}; \quad (2.6.14)$$

the factor $n^{c^*(P)}$ is exactly consumed by the Euler ledger.

For the global ribbon estimate every component is represented by the algebraic presentation $\zeta_{P,\tau}$ from Theorem 2.16. It is not evaluated or normed separately. Interface legs are declared typed slots and never count as smoothing-carrier identities; only a genuinely adjoined I_H costs n .

Proposition 2.21 (Closed simultaneous-component factorization). *Fix a scalar rooted centered orthogonal-Weingarten ribbon map Γ and remove passive singleton color components by centering. Before estimating anything, cut simultaneously every row or trace wire joining distinct coefficient components, record every resulting port, choose a maximizing smoothing and all prime tags, and record one base on each original $p \vee r$ circuit. This declares τ . For every pure-Weyl component P with $m_P \geq 2$, discard the forbidden-pair zero terms and apply Lemma 2.15; hence $c^*(P) \geq 2$ and Theorem 2.16 gives a presentation $\zeta_{P,\tau}$ satisfying*

$$\|\zeta_{P,\tau}\|_{\mathcal{P}_\tau} \leq A^{m_P} n^{c^*(P)-2} \|C_P\|_2^2 \|C_P\|_{\text{op}}^{m_P-2}. \quad (2.6.14a)$$

All cut interfaces can be reclosed at once by a normalized scalar projective contraction $\mathcal{G}_\Gamma$ with

$$\mathcal{A}_\Gamma = \omega_\Gamma \mathcal{G}_\Gamma \left(\widehat{\bigotimes_P \zeta_{P,\tau}} \right), \quad \|\mathcal{G}_\Gamma\|_\pi \leq 1, \quad |\omega_\Gamma| \leq C^m n^{F-4m+\ell_\Gamma-g_{\text{wg}}}. \quad (2.6.14b)$$

No component is estimated inside a gap of another component and no component token is split.

Proof. The declarations make Theorem 2.16 applicable to every component without opening a later port. Tensor the resulting algebraic presentations and reclose every interface once. Lemma 2.7 gives

$$\mathcal{G}_\Gamma = n^{-F} \text{Contr}_{\Gamma,\tau}^{\text{raw}}, \quad \|\mathcal{G}_\Gamma\|_\pi \leq 1,$$

and proves that all compact bends on one original circuit use its one base, rather than separate local normalizations. Since Equation (2.6.10a) gives $|w_\Gamma| \leq C^m n^{-4m+\ell_\Gamma-g_{\text{wg}}}$; putting $\omega_\Gamma = n^F w_\Gamma$ proves Equation (2.6.14b). $\square$

Proposition 2.22 (Prescribed gates in a terminal scalar). *In the setting of Proposition 2.21, one may cut any finite family of type-compatible interfaces, insert bounded maps $G_1, \dots, G_s$, and retain the cuts until the complete contraction is closed to a scalar. Suppose that the G_j contain no hidden unnormalized cup or cap. Then the estimate of Proposition 2.21 is multiplied by at most*

$$\prod_{j=1}^s \|G_j\|, \quad (2.6.14c)$$

and the raw-to-normalized circuit factor remains exactly n^F . The same assertion holds when one coefficient component carries the indivisible paired marked-root token: after its two fronts are closed in the terminal scalar, its color weight is β_j^2 .

Proof. Choose one projective atomic presentation of every coefficient component, but take no norm of the temporarily open global output. Insert every G_j in its prescribed typed slot before contraction. Apply Lemma 2.7 once to the resulting complete graph. Its path/cycle proof charges $\|G_j\|$ when that gate is met and leaves the correlated-front cost unchanged. Sum the atomic presentations only after the final scalar contraction.

Every fundamental $p \vee r$ circuit is still genuinely closed. On each such circuit the remaining raw cup–cap closure is exactly

$$\mathrm{Tr}_H A = n \widehat{\varepsilon}(A \otimes I_H) \widehat{\eta}.$$

Thus none of the F factors of n is replaced by a half-loop or an open-output normalization. A prescribed gate cannot change this identity because it has already been charged by its operator norm and introduces no unnormalized duality map. Finally, the paired marked-root token is kept indivisible until its two Hilbert–Schmidt fronts meet in the terminal scalar; their star-normalized square is precisely $N^{-1} \|W_j\|_2^2 = \beta_j^2$. This proves the proposition. $\square$

Definition 2.23 (Positive ribbon majorant). Fix a ribbon record Γ , a circuit state τ , and a designated set $\mathcal{E}$ of factored endpoint or packet resources. Expand every remaining coefficient component in its circuit-anchored projective presentation and insert every prescribed fixed gate before closing the original circuit bases. Define $\mathfrak{a}_{\Gamma, \tau; \mathcal{E}}$ to be $|\omega_\Gamma|$ times the sum of the absolute scalar coefficients multiplied by all remaining positive projective and trace-class costs. No endpoint weight or debit in $\mathcal{E}$ is included. In the monomer–dimer applications of [Section 3](#), the fixed recurrence scalar $(-\gamma)^a$ is also declared a factored resource: it is not one of the “scalar coefficients” in this definition and is restored exactly once by the resource ledger. Local fold, transport, and projective-presentation coefficients do remain inside the majorant. When τ and $\mathcal{E}$ are clear we write $\mathfrak{a}_\Gamma$; subscripts such as $\mathfrak{a}_{\Gamma, \mathcal{C}}$ name the part of the record left open.

This is a positive presentation cost, not the possibly cancelling value of an open contraction. Consequently it dominates every compatible partial trace-class contraction obtained from the same atoms. The operator-norm estimate needed for the Kac root-open specialization is the separate, proved statement [Lemma 2.8](#); it is not built into this definition and no replica factorization is assumed. The later allocation argument specifies which carriers are left bare and proves the corresponding Euler bound; no second copy of this majorant is introduced there.

Theorem 2.24 (Weighted packet/ribbon estimate). *For every fixed $L < \infty$ there are $C_L < \infty$ and $n_0(L) < \infty$ such that, whenever $n \geq n_0(L)$, every scalar rooted centered map with $m \leq L \log n$ satisfies*

$$|\mathcal{A}_\Gamma| \leq N C_L^m n^{-g} \prod_j \mu_j^{q_j} K^{m_j - q_j}. \quad (2.6.15)$$

At a marked root, its pure-Weyl color factor may be retained as β_j^2 instead of being relaxed using [Equation \(2.6.10\)](#).

Proof. For the scalar statement apply [Proposition 2.21](#), estimate every $\zeta_{P, \tau}$ by [Equation \(2.6.14a\)](#), and glue them once by [Equation \(2.6.14b\)](#). The Euler identity [Equation \(2.6.12\)](#) cancels the carrier powers against the raw face/row normalization and leaves $n^{2-g} \asymp N n^{-g}$. [Equations \(2.6.10\)](#) to [\(2.6.11\)](#) and the fixed-defect count [Equation \(2.6.13\)](#) give [Equation \(2.6.15\)](#), including the retained marked color factor. $\square$

3 Moments, renewal, and marked shells

3.1 Traffic moments and a fixed covariance band

We will evaluate low-degree moments with the orthogonal Weingarten formula. The following standard bridge explains why this is legitimate although the walk lives on the connected component $\mathrm{SO}(n)$.

Lemma 3.1 (SO(n)–O(n) moment bridge). *If $d < n$ and p is a polynomial of total degree at most d in the matrix entries of U , then*

$$\int_{\text{SO}(n)} p(U) dU = \int_{\text{O}(n)} p(U) dU.$$

Proof. For $G \in \{\text{SO}(n), \text{O}(n)\}$, the tensor $T_d^G = \int_G U^{\otimes d} dU$ is the orthogonal projection of $H^{\otimes d}$ onto its G -invariant subspace. In tensor degree $d < n$, the first fundamental theorem for the orthogonal group says that the $\text{SO}(n)$ -invariant tensors are exactly the linear span of the pair-contraction tensors (and there are none in odd degree). The orientation tensor first occurs in degree n . Pair-contraction tensors are already $\text{O}(n)$ -invariant, so $(H^{\otimes d})^{\text{SO}(n)} = (H^{\otimes d})^{\text{O}(n)}$ and $T_d^{\text{SO}} = T_d^{\text{O}}$. Contracting this equality with the coefficient tensor of p proves the assertion. $\square$

By Lemma 3.1, for independent Haar $U_j \in \text{SO}(n)$ acting through $\rho = \Lambda^2$, let W_j be the pure-Weyl projection of $C_j = B_j - \mu_j I$ and define

$$X_{\mathcal{W}} = \sum_j \rho(U_j)^* W_j \rho(U_j), \quad c_0 = \sum_j \mu_j. \quad (3.1.1)$$

Centering removes singleton color components. If the surviving component sizes are $r_1, \dots, r_q \geq 2$, then Equation (2.6.15) gives total color weight at most

$$c_0^q K^{2k-q} \leq (K + \sqrt{K c_0})^{2k}. \quad (3.1.2)$$

Using Equation (2.6.13), the defect sum is

$$\sum_{g \geq 0} \left(\frac{C_L k^C}{n} \right)^g = O_L(1) \quad (2k \leq L \log n). \quad (3.1.3)$$

Therefore

$$\mathbb{E} \text{Tr } X_{\mathcal{W}}^{2k} \leq N \left[A_L (K + \sqrt{K c_0}) \right]^{2k}. \quad (3.1.4)$$

Lemma 3.2 (Quantitative pseudo-Haar hybrid). *Fix $L, \eta < \infty$. There is $C_0 = C_0(L, \eta)$ such that, whenever $T \geq C_0 n \log^3 n$, every polynomial p of degree at most $d \leq L \log n$ on $\text{SO}(n)$ satisfies, uniformly in the starting point x ,*

$$|\mathbb{E}_x p(X_T) - \mathbb{E}_H p| \leq n^{-\eta \log n} \|p\|_{\infty}. \quad (3.1.5)$$

For a polynomial of independent separator blocks, replacing b blocks one at a time costs at most b times the right side, with the supremum norm taken after the other blocks are frozen.

Proof. First replace p by $p - \int p dH$, which changes neither side of Equation (3.1.5) and has supremum norm at most $2\|p\|_{\infty}$. Replace this centered polynomial by the reduced representative p^* of Pillai–Smith–Vaikuntanathan [PSV26], which has the same restriction to $\text{SO}(n)$. Their Theorem 4.2 gives

$$\text{Var}_H(p^*) \geq \frac{\|p^*\|_{\text{coeff},2}^2}{(d+1)n^d}.$$

Since $\text{Var}_H(p^*) \leq 4\|p\|_{\infty}^2$, and the graded degree- d representation has at most $(d+1)n^{2d}$ coefficients,

$$\|p^*\|_{\text{coeff},1} \leq 2(d+1)n^{3d/2} \|p\|_{\infty}.$$

Their Theorem 4.1 gives, uniformly over all monomial degrees at most d , the estimate

$$|\mathbb{E}_x \mathbf{m}(X_T) - \mathbb{E}_H \mathbf{m}| \leq C d n^{-s/2},$$

where

$$s = \left\lfloor \frac{T}{C n (d + \log n) \log n} \right\rfloor.$$

Expanding p^* and summing its coefficients therefore gives

$$|\mathbb{E}_x p(X_T) - \mathbb{E}_H p| \leq C(d+1) d n^{3d/2-s/2} \|p\|_\infty.$$

For $d \leq L \log n$ and $T = C_0 n \log^3 n$, choose C_0 so that $s \geq (3L + 2\eta + 4) \log n$; the factor $(d+1)d$ is then absorbed by the two spare powers in the exponent. This proves Equation (3.1.5) for large n . The last assertion is the telescoping identity obtained by replacing the independent blocks sequentially. Because the one-block estimate is uniform in its start, it applies after all other blocks are frozen. $\square$

For a fixed index word $I_1, \dots, I_m$, write

$$F(\theta) = R_{I_m}(\theta_m) \cdots R_{I_1}(\theta_1).$$

With the Hilbert–Schmidt orthonormal basis $a_{ij} = (E_{ij} - E_{ji})/\sqrt{2}$, its right-trivialized angle derivatives are

$$v_t = \text{Ad}(P_{t+1})a_{I_t}, \quad P_{t+1} = R_{I_m} \cdots R_{I_{t+1}}.$$

Put

$$D_m = \sum_{t=1}^m v_t v_t^*, \quad \text{Tr } D_m = m. \quad (3.1.6)$$

The Malliavin covariance in the normalization of the coupling argument is $M = 2D_m$. If $j_t = \sqrt{2}v_t$, $A_t = \text{ad}_{j_t}$, and $M_{<t} = \sum_{s<t} j_s j_s^*$, then on the regular set the exact horizontal-lift score representative is

$$b = -M^{-1} \sum_{t=1}^m [A_t, M_{<t}] M^{-1} j_t. \quad (3.1.7)$$

Indeed, the horizontal lift of $x \in \mathfrak{so}(n)$ has angle coordinates $\langle j_t, M^{-1}x \rangle$; differentiating its divergence uses $\partial_t j_t = 0$ and $\partial_t M = [A_t, M_{<t}]$, giving Equation (3.1.7). Moreover $[A_t, j_t j_t^*] = 0$ and

$$\sum_t [j_t, M^{-1} j_t] = 0,$$

because the coefficient of $[e_a, e_b]$ in this sum is $(MM^{-1})_{ab} = \delta_{ab}$. Hence Equation (3.1.7) is unchanged if $M_{<t}$ is replaced by $(M_{<t} - M_{>t})/2$.

Proposition 3.3 (A constant covariance band). *For every sufficiently large fixed integer c and $m = cN$, there are $0 < a_c < b_c < 2c + 1$ such that*

$$\Pr\{a_c I \preceq D_m \preceq b_c I\} = 1 - \delta_n, \quad \delta_n \longrightarrow 0. \quad (3.1.8)$$

Proof. Invoke Theorem 2.24 in the fixed logarithmic degree range used below. Then split the path into consecutive blocks of length $R = C_0 n \log^3 n$. Fix one parity $\varepsilon \in \{\text{o}, \text{e}\}$ and condition only on the complete trajectories of the retained ε -blocks; the intervening blocks are not conditioned

upon. Reverse exposure inside a retained block makes its transported rank-one deposits an adapted positive sequence X_t with $0 \preceq X_t \preceq I$ and conditional mean I_W/N . Since X_t is a projection,

$$\mathbb{E}_{t-1} e^{\theta X_t} = I + (e^\theta - 1) \mathbb{E}_{t-1} X_t \preceq \exp\{(e^\theta - 1)I/N\}.$$

The iterated matrix Laplace-transform argument (equivalently the conditional Chernoff corollary of matrix Freedman [Tro11]) gives, for $u > |J|/N$,

$$\Pr\left\{\lambda_{\max}\left(\sum_{t \in J} X_t\right) \geq u\right\} \leq N \inf_{\theta > 0} e^{-\theta u + (e^\theta - 1)|J|/N} \leq N \left(\frac{e|J|}{uN}\right)^u. \quad (3.1.8a)$$

Taking $u = 12$ and a union bound over the $O(n/\log^3 n)$ retained ε -blocks gives an event $\mathcal{G}_\varepsilon$, measurable with respect to the retained data, whose complement has probability $n^{-9+o(1)}$, and on which

$$0 \preceq B_J \preceq 12I, \quad N^{-1} \text{Tr } B_J = |J|/N. \quad (3.1.9)$$

(For example, the Chernoff upper tail at 12 is $N(e|J|/(12N))^{12} = n^{-10+o(1)}$.)

The Ricci contribution is removed directly, without Haarizing its blocks. Let $\tilde{B}_{z_t}$ be the matrix in Equation (2.6.2), transported into the common endpoint frame, and put

$$\mathfrak{B}^\varepsilon = \sum_t \mathbf{1}_{\{t \text{ belongs to an } \varepsilon\text{-block}\}} \tilde{B}_{z_t}. \quad (3.1.9a)$$

The parity mask is deterministic in reverse time. Lemma 2.17 gives the matrix-martingale identities

$$\mathbb{E}_{t-1} \tilde{B}_{z_t} = 0, \quad \|\tilde{B}_{z_t}\|_{\text{op}} \leq \frac{1}{n}, \quad \mathbb{E}_{t-1} (\tilde{B}_{z_t})^2 = \frac{2}{n^2(n-2)} I_H. \quad (3.1.9b)$$

Its predictable quadratic variation is at most

$$\frac{2m}{n^2(n-2)} = O_c(n^{-1}).$$

Matrix Freedman's inequality [Tro11] and a union bound over the two parities imply

$$\|\mathfrak{B}^\varepsilon\|_{\text{op}} = O_{\mathbb{P}}\left(\sqrt{\frac{\log n}{n}} + \frac{\log n}{n}\right) = o_{\mathbb{P}}(1). \quad (3.1.9c)$$

Equivariance and linearity of $B \mapsto X_B$ identify the complete Ricci contribution of this parity with $X_{\mathfrak{B}^\varepsilon}$; hence its operator norm is at most $2\|\mathfrak{B}^\varepsilon\|_{\text{op}} = o_{\mathbb{P}}(1)$.

Write each retained centered block as $B_J - \mu_J I = W_J + X_{A_J}$. Orthogonality gives $N^{-1}\|W_J\|_2^2 \leq 12\mu_J$. The block version of Equation (2.6.6a), followed by a union bound, supplies a retained-data event of probability $1 - o(1)$ on which $\max_J \|X_{A_J}\|_{\text{op}} = o(1)$. Since $\mu_J = |J|/N = o(1)$, Equation (3.1.9) then gives

$$\|W_J\|_{\text{op}} \leq \|B_J\|_{\text{op}} + \mu_J + \|X_{A_J}\|_{\text{op}} \leq 13 \quad (3.1.9d)$$

for all sufficiently large n .

Conditional on the retained blocks, the intervening products are independent R -step Kac bridges. For the trace polynomial $p = \text{Tr}(X_W^{2k})$, its degree in the matrix entries of each bridge is at most $8k$ (a Λ^2 conjugation contains four matrix entries) and, with $b = O(n/\log^3 n)$ retained blocks,

$$\|p\|_\infty \leq N \left(\sum_J \|W_J\|\right)^{2k} \leq N(13b)^{2k} = n^{O(k)}.$$

Choose the η in [Lemma 3.2](#) larger than the fixed exponent implicit in $n^{O(k)}$ and replace the bridges one at a time. The bound is uniform in their starting points, and the event $\mathcal{G}_\varepsilon$ depends only on retained data, so no separator law has been changed by conditioning. The accumulated hybrid error is $o(1)$. After Haar replacement, the cumulative retained-block orientations are independent Haar variables: this is the triangular product-Haar change of variables

$$(V_1, \dots, V_b, H) \longleftrightarrow (V_1 \cdots V_b H, V_2 \cdots V_b H, \dots, V_b H, H).$$

A terminal short block is handled by the common auxiliary Haar conjugation H , which leaves the trace polynomial unchanged.

Apply [Equation \(3.1.4\)](#) to the Weyl Haar comparison with $K = 13$ and $c_\varepsilon = \sum_{J \in \varepsilon} \mu_J = c/2 + O(R/N) = c/2 + o(1)$. With $k = \lceil L_0 \log n \rceil$, Markov's inequality gives

$$\Pr\left\{\|X_{\mathcal{W}}^\varepsilon\| > 2A_{L_0}(13 + \sqrt{13c_\varepsilon})\right\} \leq N 2^{-2k} + o(1). \quad (3.1.9e)$$

Taking L_0 large and restoring the $o_{\mathbb{P}}(1)$ Ricci term gives for the retained parity

$$\|D^\varepsilon - c_\varepsilon I\| \leq C(1 + \sqrt{c}) + o(1)$$

with probability $1 - o(1)$. Analyze the other parity separately and combine the two events by a union bound. Positivity, $D_m = D^o + D^e$, and a union bound now give

$$\lambda_{\min}(D_m) \geq \frac{c}{2} - C(1 + \sqrt{c}) - o(1), \quad \lambda_{\max}(D_m) \leq c + C(1 + \sqrt{c}) + o(1). \quad (3.1.10)$$

For sufficiently large fixed c the lower endpoint is positive and the upper endpoint is strictly below $2c + 1$; taking slightly wider deterministic endpoints inside $(0, 2c + 1)$ proves [Equation \(3.1.8\)](#). $\square$

3.2 Chronological spectral calculus and passive renewal

3.2.1 Common spectral blocks and chronological exposure

We first record the temporal estimates, including their normalizations. For the right-circle average P_i in coordinate plane i , let

$$\mathcal{K} = \frac{1}{N} \sum_{i=1}^N P_i, \quad (\mathcal{D}f)_i = \frac{1}{\sqrt{N}}(I - P_i)f. \quad (3.2.1)$$

These operators act on $\mathcal{H} = L^2(\mathrm{SO}(n); V)$ for any Hilbert coefficient space V . We place all starred operators in the common counting-measure space $\mathcal{H}_\star = \ell^2([N]) \otimes L^2(\mathbb{T}) \otimes \mathcal{H}$, embedding $\mathcal{D}f$ as a function constant in θ . Let L_i be the infinitesimal generator of the same circle and $(\mathcal{L}f)(i, \theta) = N^{-1/2}L_i f$. Finally define

$$(\mathcal{T}f)(g, i, \theta) = N^{-1/2}f(gR_i(\theta)), \quad (\mathcal{E}h)(g) = N^{-1/2} \sum_i \int h(g, i, \theta) \frac{d\theta}{2\pi}, \quad \mathcal{G} = (I - \mathcal{E}^* \mathcal{E})\mathcal{T}. \quad (3.2.2)$$

With these factors $\mathcal{T}$ is an isometry, $\mathcal{E}$ a coisometry, and every mixed block in [Equation \(3.2.4a\)](#) is an operator on the same Hilbert space.

Lemma 3.4 (The Kac Riesz block bounds). *The following statements hold not only for the right regular action on $\mathcal{H}$ but for every finite-dimensional unitary $\mathrm{SO}(n)$ -module: in that formulation P_i is the orthogonal average of the circle $\pi(R_i(\theta))$, and the definitions above use $N^{-1/2}\pi(R_i(\theta))$ in*

place of $\mathcal{T}$. On the orthogonal complement of the $\mathcal{K}$ -invariant functions, with $(I - \mathcal{K})^\dagger$ denoting the Moore–Penrose inverse,

$$\begin{aligned}\|\mathcal{D}(I - \mathcal{K})^\dagger \mathcal{D}^*\| &\leq 1, \\ \|\mathcal{G}(I - \mathcal{K})^\dagger \mathcal{G}^*\| &\leq 2.\end{aligned}\tag{3.2.3}$$

On functions of circle Fourier degree at most d ,

$$\|\mathcal{L}(I - \mathcal{K})^\dagger \mathcal{L}^*\| \leq d^2.\tag{3.2.4}$$

The mixed blocks satisfy

$$\begin{aligned}\|\mathcal{D}(I - \mathcal{K})^\dagger \mathcal{G}^*\| &\leq \sqrt{2}, & \|\mathcal{G}(I - \mathcal{K})^\dagger \mathcal{D}^*\| &\leq \sqrt{2}, \\ \|\mathcal{L}(I - \mathcal{K})^\dagger \mathcal{D}^*\| &\leq d, & \|\mathcal{D}(I - \mathcal{K})^\dagger \mathcal{L}^*\| &\leq d, \\ \|\mathcal{L}(I - \mathcal{K})^\dagger \mathcal{G}^*\| &\leq \sqrt{2}d, & \|\mathcal{G}(I - \mathcal{K})^\dagger \mathcal{L}^*\| &\leq \sqrt{2}d.\end{aligned}\tag{3.2.4a}$$

All these bounds remain true after tensoring by I_V . They also hold, with the same constants, when $(I - \mathcal{K})^\dagger$ is replaced everywhere by a finite Green kernel $R_s = \sum_{\ell=0}^s \mathcal{K}^\ell$ acting between centered channels.

Proof. Every P_i is an orthogonal projection, hence $\mathcal{D}^* \mathcal{D} = I - \mathcal{K}$. Polar decomposition shows that $\mathcal{D}(I - \mathcal{K})^\dagger \mathcal{D}^*$ is the projection onto $\overline{\text{ran } \mathcal{D}}$, proving the first bound. Also $\mathcal{E} \mathcal{T} = \mathcal{K}$ and $\mathcal{T}^* \mathcal{E}^* = \mathcal{K}$, so

$$\mathcal{G}^* \mathcal{G} = I - \mathcal{K}^2.$$

Since $0 \preceq \mathcal{K} \preceq I$, the nonzero spectral part of the second Green block is $(I - \mathcal{K}^2)/(I - \mathcal{K}) = I + \mathcal{K}$, whose norm is at most two. Finally, on nonzero Fourier modes $e^{i\ell\theta}$ with $|\ell| \leq d$, L_i is multiplication by $i\ell$, whereas $I - P_i$ is the identity. Thus $\mathcal{L} = B\mathcal{D}$ with $\|B\| \leq d$, and Equation (3.2.4) follows from the first bound. For R_s , the two diagonal spectral multipliers are

$$(1 - \lambda) \sum_{\ell=0}^s \lambda^\ell = 1 - \lambda^{s+1} \leq 1, \quad (1 - \lambda^2) \sum_{\ell=0}^s \lambda^\ell = (1 + \lambda)(1 - \lambda^{s+1}) \leq 2.$$

The circle estimate follows again from $\mathcal{L} = B\mathcal{D}$. Thus all diagonal estimates hold for the finite kernels as well. If R denotes either $(I - \mathcal{K})^\dagger$ or R_s , positivity of

$$\begin{pmatrix} ARA^* & ARB^* \\ BRA^* & BRB^* \end{pmatrix} = \begin{pmatrix} A \\ B \end{pmatrix} R(A^*, B^*)$$

implies the operator Cauchy–Schwarz inequality $\|ARB^*\|^2 \leq \|ARA^*\| \|BRB^*\|$. Apply it to the three pairs among $\mathcal{D}, \mathcal{G}, \mathcal{L}$ and use Equations (3.2.3) to (3.2.4); this gives Equation (3.2.4a). Tensoring does not change any operator norm. $\square$

The finite- N root correction must be kept in the recurrence. Since a marked deposit satisfies $P_t^2 = P_t$ while each of the other $m - 1$ isotropic deposits has mean I/N , put

$$\gamma = \gamma_{m,N} := \frac{m-1}{N} = c - \frac{1}{N}.\tag{3.2.5a}$$

For every possible marked pivot t this choice gives the exact algebraic centering identity

$$D_m - (\gamma + 1)I = \sum_{s \neq t} (v_s v_s^* - N^{-1}I) + (v_t v_t^* - I).\tag{3.2.5b}$$

Thus the $+1$ collision and the $(m-1)/N$ fresh-deposit backtrack are kept separate at finite N ; no $O(N^{-1})$ discrepancy is discarded. Normalize

$$Y = \frac{D_m - (\gamma + 1)I}{2\sqrt{\gamma}}, \quad Q_k(D) = \gamma^{k/2} U_k(Y), \quad (3.2.5)$$

where U_k is the Chebyshev polynomial of the second kind. Then

$$Q_{k+1} = (D - (\gamma + 1)I)Q_k - \gamma Q_{k-1} \quad (3.2.6)$$

and the noncommutative Christoffel–Darboux identity is

$$dQ_k(D)[E] = \sum_{j=0}^{k-1} Q_{k-1-j}(D)EQ_j(D). \quad (3.2.7)$$

Both statements follow by induction from $Q_0 = I$ and $Q_1 = D - (\gamma + 1)I$.

All vectors below are in the single right-trivialized endpoint frame. Put

$$\chi_{t,k} = \partial_{\theta_t}(Q_k(D_m)v_t), \quad \mathcal{C}_k = \sum_{t=1}^m \chi_{t,k}. \quad (3.2.8)$$

Here v_t is independent of its own angle and

$$\partial_{\theta_t} v_s = [j_t, v_s] \quad (s < t), \quad \partial_{\theta_t} v_s = 0 \quad (s \geq t), \quad \partial_{\theta_t} D_m = [A_t, D_{<t}], \quad (3.2.9)$$

where $A_t = \text{ad}_{j_t}$ and $D_{<t} = \sum_{s<t} v_s v_s^*$.

Proposition 3.5 (Elementary first marked-shell estimate). *Assume $n \geq 4$ and $m \leq cN$, where $c \geq 1$. The expectation is over both the independent uniform coordinate planes and their independent uniform angles. Then*

$$\boxed{\mathbb{E}\|\mathcal{C}_1\|^2 \leq \frac{2n(n-1)}{n+2}(c^2 + 2c) \leq 2n(c^2 + 2c).} \quad (3.2.9a)$$

Proof. Since $Q_1(D) = D - (\gamma + 1)I$, [Equations \(3.2.7\) and \(3.2.9\)](#) give $\mathcal{C}_1 = \sum_t [A_t, D_{<t}]v_t$. Moreover $A_t v_t = [j_t, v_t] = 0$ and $j_t = \sqrt{2}v_t$, so, with $D_{>s} = \sum_{t>s} v_t v_t^*$,

$$\mathcal{C}_1 = \sqrt{2} \sum_{s<t} \langle v_s, v_t \rangle [v_t, v_s] = \sqrt{2} \sum_s [D_{>s} v_s, v_s]. \quad (3.2.9b)$$

Let $\mathcal{G}_{s+1} = \sigma((I_u, \theta_u) : u > s)$. Conditional on $\mathcal{G}_{s+1}$, $D_{>s}$ is fixed and v_s is uniform on the orthonormal frame $w_i = \text{Ad}(P_{s+1})a_i$. For every symmetric operator D on W ,

$$\sum_i [Dw_i, w_i] = 0, \quad (3.2.9c)$$

because the matrix coefficients of D are symmetric whereas the structure constants are antisymmetric in the two input indices. The summands in [Equation \(3.2.9b\)](#) are therefore reverse martingale differences, and

$$\mathbb{E}\|\mathcal{C}_1\|^2 = 2 \sum_s \mathbb{E}\|[D_{>s} v_s, v_s]\|^2. \quad (3.2.9d)$$

Write $i \sim j$ when the two coordinate edges share exactly one vertex, and, for a symmetric matrix $E = (E_{ij})$ in the Kac basis, put

$$\mathcal{A}(E) = \sum_{\{i,j\}: i \sim j} E_{ij}^2,$$

where the sum is over unordered adjacent pairs. The nonzero $[a_j, a_i]$ for fixed i are mutually orthogonal and have squared norm $1/2$. Hence, for every scalar α ,

$$\sum_i \|[(E + \alpha I)a_i, a_i]\|^2 = \mathcal{A}(E). \quad (3.2.9e)$$

After conjugating into the time- s frame and centering the suffix covariance, Equation (3.2.9e) gives

$$\mathbb{E}\left[\|D_{>s}v_s, v_s\|^2 \mid \mathcal{G}_{s+1}\right] = \frac{1}{N}\mathcal{A}(E_s). \quad (3.2.9f)$$

For $\ell \geq 1$ define in this fixed frame

$$u_\ell = \text{Ad}\left((R_{s+\ell-1} \cdots R_{s+1})^{-1}\right) a_{I_{s+\ell}}, \quad Z_\ell = u_\ell u_\ell^* - N^{-1}I_W.$$

The rotation at time $s + \ell$ drops because it fixes its own generator. The centered projectors Z_ℓ are forward martingale differences, so polarization kills all cross terms and

$$\mathbb{E}\mathcal{A}(E_s) = \sum_{\ell=1}^{m-s} \mathbb{E}S(u_\ell), \quad S(u) = \sum_i \langle u, a_i \rangle^2 \| [u, a_i] \|^2. \quad (3.2.9g)$$

For a unit decomposable bivector u , put $x_{pq} = \langle u, a_{pq} \rangle$, $d_p = \sum_{q \neq p} x_{pq}^2$, and $L_u = -2u_{\text{mat}}^2$. Then L_u is the rank-two plane projector and $(L_u)_{pp} = d_p$. Direct multiplication of the coordinate skew matrices gives

$$\| [u, a_{pq}] \|^2 = \frac{1}{2}(d_p + d_q - 2x_{pq}^2),$$

whence

$$S(u) = \frac{1}{2} \sum_p d_p^2 - \sum_{p < q} x_{pq}^4 \leq \frac{1}{2} F(L_u), \quad F(L) = \sum_p L_{pp}^2. \quad (3.2.9h)$$

It remains to average this elementary statistic. If L' is obtained from a rank-two projector L by one uniform Kac conjugation, averaging first over the angle in the selected plane (p, q) yields

$$\mathbb{E}_\theta[F(L') - F(L) \mid p, q] = L_{pq}^2 - \frac{1}{4}(L_{pp} - L_{qq})^2.$$

Since $\text{Tr } L = \text{Tr } L^2 = 2$,

$$\sum_{p < q} L_{pq}^2 = 1 - \frac{1}{2}F(L), \quad \sum_{p < q} (L_{pp} - L_{qq})^2 = nF(L) - 4.$$

Therefore

$$\mathbb{E}[F(L')] - F(L) = \frac{1}{N} \left(2 - \frac{n+2}{4} F(L) \right). \quad (3.2.9i)$$

Writing $\gamma_n = (n+2)/(4N)$ and $\rho_n = 1 - \gamma_n$, and using $F(L_0) = 2$, the exact recursion is

$$f_j := \mathbb{E}F(L_j) = \frac{8}{n+2} + \left(2 - \frac{8}{n+2} \right) \rho_n^j. \quad (3.2.9j)$$

Combining Equations (3.2.9d) to (3.2.9h) and then using Equation (3.2.9j),

$$\mathbb{E}\|C_1\|^2 \leq \frac{1}{N} \sum_{s=1}^{m-1} \sum_{\ell=1}^{m-s} f_{\ell-1} \leq \frac{4m^2}{N(n+2)} + \frac{8m}{n+2}.$$

The right side is increasing in m ; inserting $m \leq cN$ and $N = n(n-1)/2$ proves Equation (3.2.9a). $\square$

Lemma 3.6 (One-pivot chronological boundary exposure). *Expand one monomial occurring in $\langle \mathcal{C}_k, \mathcal{C}_k \rangle$ but keep, in each replica, the sum over all derivative positions before taking an absolute value. At the latest unexposed pivot t , every connected earlier-time component Φ_C satisfies the exact Ward identity*

$$\sum_{u \in C} d\rho_u(j_t) \Phi_C = \sum_{e \in \partial C} \varepsilon_e d\rho_e(j_t) \Phi_C, \quad \varepsilon_e \in \{-1, 1\}. \quad (3.2.10)$$

All coefficients on the two sides of an exposed boundary are independent of the fresh pivot. For one fixed pivot and q labelled slots, the past/current/future classification has 3^q states. This is only a one-pivot statement; no bound on the power of the resulting subset transfer is inferred from 3^q .

Proof. Equation (3.2.9) says that every earlier occurrence receives the same infinitesimal diagonal action $d\rho(j_t)$, every later occurrence is fixed, and a current occurrence is a same-time collision. Differentiate the equivariance identity $\Phi_C \rho_{\text{in}}(e^{sj_t}) = \rho_{\text{out}}(e^{sj_t}) \Phi_C$ at $s = 0$. Internal actions cancel in pairs and give Equation (3.2.10) on the boundary. Conditioning on every variable but (I_t, θ_t) makes all later coefficients fixed. The current deposit obeys $[A_t, v_t v_t^*] = 0$; its other repetitions are the explicit collision/backtrack children of the recurrence Equation (3.2.6). Expose pivots in decreasing time, so no future coefficient ever depends on a subsequently integrated pivot.

Each polynomial slot is earlier, current, or later relative to this one pivot, which proves the final assertion. Repeating the classification naively would produce ordered Bell growth; that repetition is handled instead by the global transfer-to-defect encoding below. Crucially, Equation (3.2.10) is used before any norm or absolute value. $\square$

Lemma 3.7 (Universal two-gradient root identity). *Let $X, Z \in \text{End}(W)$ be arbitrary, let $a_1, \dots, a_N$ be the Kac orthonormal basis of $W = \mathfrak{so}(n)$, and put $A_i = \text{ad}_{a_i}$. Define*

$$\begin{aligned} \mathcal{R}_i(X, Z) &= \{X, [A_i, Z]\} a_i, & U_i(X, Z) &= \{X, Z\} a_i, \\ E_i(X, Z) &= \{[A_i, X], Z\} a_i, & \bar{E}(X, Z) &= \frac{1}{N} \sum_i E_i(X, Z). \end{aligned} \quad (3.2.10a)$$

Put $\Psi(T) = \sum_j A_j T a_j$. Then, with no projection onto curvature summands,

$$\mathcal{R}_i = A_i U_i - (E_i - \bar{E}) - \bar{E}, \quad \bar{E} = \frac{1}{N} \sum_j A_j V_j + H(X, Z), \quad V_j = X Z a_j, \quad (3.2.10b)$$

where

$$H(X, Z) = \frac{1}{N} \{-X \Psi(Z) + Z \Psi(X)\}. \quad (3.2.10b')$$

Moreover, writing $\|\cdot\|_2$ for Hilbert–Schmidt norm,

$$\begin{aligned} \frac{1}{N} \sum_i \|U_i\|^2 &\leq \frac{4}{N} \|X\|_2^2 \|Z\|_{\text{op}}^2, \\ \frac{1}{N} \sum_j \|V_j\|^2 &\leq \frac{1}{N} \|X\|_2^2 \|Z\|_{\text{op}}^2, \\ \frac{1}{N} \sum_i \|E_i - \bar{E}\|^2 &\leq \frac{16(n-2)}{N} \|X\|_2^2 \|Z\|_{\text{op}}^2. \end{aligned} \quad (3.2.10c)$$

The residue satisfies

$$\|H(X, Z)\| \leq \frac{\sqrt{n-2}}{N} (\|X\|_{\text{op}} \|Z\|_2 + \|Z\|_{\text{op}} \|X\|_2). \quad (3.2.10e)$$

If X and Z are symmetric, then $\Psi(X) = \Psi(Z) = 0$, so $H = 0$. The identities and estimates remain valid when all maps are tensored by an ancillary identity and the Hilbert–Schmidt norm is taken on the amplified space.

Proof. Since $A_i a_i = 0$, direct expansion gives

$$\begin{aligned} A_i\{X, Z\}a_i - \{[A_i, X], Z\}a_i &= A_i Z X a_i + X A_i Z a_i - Z A_i X a_i \\ &= \{X, [A_i, Z]\}a_i. \end{aligned}$$

This is the first equality in Equation (3.2.10b) after adding and subtracting $\bar{E}$. For every symmetric $T \in \text{End}(W)$, unimodularity gives

$$\sum_i A_i T a_i = 0. \quad (3.2.10d)$$

Indeed, its inner product with $x \in W$ is $-\sum_i \langle T a_i, [a_i, x] \rangle$; the matrix of ad_x in the orthonormal basis (a_i) is skew whereas the matrix of T is symmetric, so the contraction is zero. Expanding $\sum_i E_i$ without a symmetry assumption gives

$$\sum_i E_i = \Psi(XZ) - X\Psi(Z) + Z\Psi(X),$$

which proves Equations (3.2.10b) to (3.2.10b'). The adjoint of $\Psi : \text{End}(W) \rightarrow W$ sends x to the operator whose i th column is $-A_i x$. Therefore

$$\|\Psi^* x\|_2^2 = \sum_i \|A_i x\|^2 = (n-2)\|x\|^2.$$

It follows that $\|\Psi(T)\| \leq \sqrt{n-2}\|T\|_2$, which proves Equation (3.2.10e). Equation (3.2.10d) proves the final assertion in the symmetric case.

The first two estimates in Equation (3.2.10c) are simply

$$N^{-1}\|\{X, Z\}\|_2^2 \leq 4N^{-1}\|X\|_2^2\|Z\|_{\text{op}}^2, \quad N^{-1}\|XZ\|_2^2 \leq N^{-1}\|X\|_2^2\|Z\|_{\text{op}}^2.$$

Centering is an orthogonal projection in the plane index, and hence

$$\frac{1}{N} \sum_i \|E_i - \bar{E}\|^2 \leq \frac{1}{N} \sum_i \|E_i\|^2 \leq \frac{4\|Z\|_{\text{op}}^2}{N} \sum_i \|[A_i, X]\|_2^2.$$

The commutator representation is contained in $W \otimes W$; the adjoint Casimir is $n-2$, and the tensor-product triangle inequality for its square root gives $\sum_i \|[A_i, X]\|_2^2 \leq 4(n-2)\|X\|_2^2$. This proves the last line. The same calculation acts only on the W factor, so ancillary tensoring does not alter it. $\square$

Lemma 3.8 (Raw complete same-time pinch). *In a fresh Kac frame put $P_i = a_i a_i^*$ and $W_i = P_i - N^{-1}I_W$. Before any ancillary boundary routing, the intact coordinate-plane average obeys, for arbitrary real $|\epsilon_i| \leq 1$,*

$$\left\| \frac{1}{N} \sum_i \epsilon_i W_i \right\|_{\text{op}} \leq \frac{2}{N}. \quad (3.2.10f)$$

The exact scalar identities are

$$W_i^2 = \left(1 - \frac{2}{N}\right) W_i + \frac{N-1}{N^2} I, \quad [A_i, P_i] = 0. \quad (3.2.10g)$$

If a passive map separates equal labels, then

$$\mathbf{P}_i T \mathbf{P}_i = \langle a_i, T a_i \rangle \mathbf{P}_i; \quad (3.2.10h)$$

this expression remains in the intact collision block and is not replaced by a scalar unless T itself is scalar. Relative to the Jacobi coefficients $1, \gamma$, the only naked finite- N residuals in Equation (3.2.10g) are $-2N^{-1}W_i$ and $-\gamma N^{-1}I$. The first child remains time-indexed. The second is formed only after the $m - 1$ collision labels have been summed; before residual temporal components are defined it is removed from the temporal word, with γ retained in the monomer-dimer ledger and the remaining N^{-1} kept as a favorable scalar coefficient carrying no time, gap, anchor, or translation.

Proof. The $\mathbf{P}_i$ are the mutually orthogonal rank-one projectors of the Kac basis. Hence

$$\left\| N^{-1} \sum_i \epsilon_i \mathbf{P}_i \right\|_{\text{op}} \leq N^{-1}, \quad \left\| N^{-2} \sum_i \epsilon_i I_W \right\|_{\text{op}} \leq N^{-1},$$

which proves Equation (3.2.10f). Equations (3.2.10g) to (3.2.10h) follow from $\mathbf{P}_i^2 = \mathbf{P}_i$ and $A_i a_i = 0$. Finally $(m - 1)(N - 1)N^{-2} = \gamma(1 - N^{-1})$, which gives the two displayed residuals. Equivalently, before the $m - 1$ labels are summed, the scalar child is a sum of terms $-N^{-2}I$; assigning a second translation to those terms would count the same label sum twice. This proves the last assertion and the stated temporal convention. $\square$

3.2.2 Passive singletons, typed passes, and finite Green gaps

Lemma 3.9 (Passive-singleton seed and pass). *For a Kac plane i , let*

$$W_i^\circ = \Pi_{\mathcal{W}}(\mathbf{P}_i - N^{-1}I_W)$$

be its pure-Weyl coefficient. On the common star space, with an arbitrary passive ancillary Hilbert space E , put

$$(\mathbf{M}h)(i, \theta) = (W_i^\circ \otimes I_E)h(i, \theta), \quad \mathbf{J} = \mathcal{E}^* \mathcal{E}, \quad \mathbf{Q} = I - \mathbf{J}.$$

Relative to $\text{ran } \mathbf{J} \oplus \text{ran } \mathbf{Q}$, write

$$\mathbf{M} = \begin{pmatrix} 0 & S^- \\ S^+ & V \end{pmatrix}. \quad (3.2.10n)$$

We call $0, S^-, S^+, V$ the four star statuses of a passive singleton; a status record chooses one of these four blocks for every singleton after equal-time fusion. These statuses are distinct from the derivative-placement coefficients (α, β) used in the Ricci root calculus. Then

$$\|S^-\| = \|S^+\| \leq \sqrt{\frac{n-3}{N(n-1)}} \leq N^{-1/2}, \quad \|V\| \leq 2, \quad (3.2.10o)$$

and the one-sided utilized-step maps factor through the innovation:

$$\mathcal{E} \mathbf{M} \mathcal{T} = \mathcal{E} \mathbf{M} \mathcal{G}, \quad \mathcal{T}^* \mathbf{M} \mathcal{E}^* = \mathcal{G}^* \mathbf{M} \mathcal{E}^*. \quad (3.2.10p)$$

Consequently a passive singleton isolated between invariant separators is zero. A run of $s \geq 2$ passive singletons with $s - 1$ prescribed finite Green gates has norm at most

$$\frac{2^{2s-3}}{N}. \quad (3.2.10q)$$

Thus its two endpoints create one N^{-1} resource and every interior singleton transports that same resource; no additional token is created. The same conclusion holds when an endpoint is a $\mathcal{D}$ - or $\mathcal{L}$ -incidence, with the corresponding mixed bound of [Lemma 3.4](#) and with the already existing root or packet token transported.

Proof. The coordinate pure-Weyl rows obey

$$\sum_i W_i^\circ = 0, \quad \sum_i (W_i^\circ)^2 = \frac{n-3}{n-1} I_W. \quad (3.2.10r)$$

For the first equality use $\sum_i P_i = I_W$ and equivariance of the Weyl projection. For the second, write $W_i^\circ = P_i - N^{-1} I_W - X_{B_i}$ with B_i from [Equation \(2.6.2\)](#). Both P_i and X_{B_i} are diagonal in the coordinate-wedge basis, so the sum of the squares is diagonal. Coordinate permutations make its diagonal entries equal. Orthogonality in [Equations \(2.6.3\)](#) to [\(2.6.4\)](#) gives

$$\|W_i^\circ\|_2^2 = 1 - N^{-1} - 2n^{-1} = \frac{n-3}{n-1}.$$

Taking the trace of the sum proves the second equality in [Equation \(3.2.10r\)](#). Also $\|W_i^\circ\|_{\text{op}} \leq 2$ by [Equations \(2.6.3\)](#) to [\(2.6.4\)](#).

Since $\mathcal{E}^* \xi$ is the constant star field $N^{-1/2} \xi$,

$$\mathcal{E} M \mathcal{E}^* = \frac{1}{N} \sum_i W_i^\circ \otimes I_E = 0.$$

This proves the zero upper-left block and, after inserting $I = J + Q$ next to $\mathcal{T}$, proves [Equation \(3.2.10p\)](#). Moreover, using self-adjointness of M ,

$$S^- S^{-*} = \mathcal{E} M Q M \mathcal{E}^* = \mathcal{E} M^2 \mathcal{E}^* = \frac{1}{N} \sum_i (W_i^\circ)^2 \otimes I_E.$$

[Equation \(3.2.10r\)](#) proves the first estimate in [Equation \(3.2.10o\)](#); orthogonal compression and $\sup_i \|W_i^\circ\|_{\text{op}} \leq 2$ prove the second.

Put $\mathfrak{G}_u = \mathcal{G} R_u \mathcal{G}^*$, so $\|\mathfrak{G}_u\| \leq 2$ by [Lemma 3.4](#). A run with opposite endpoint orientations is, up to adjoints and fixed contractive boundary maps,

$$S^- \mathfrak{G}_{u_1} V \mathfrak{G}_{u_2} \cdots V \mathfrak{G}_{u_{s-1}} S^+.$$

The two seed bounds, the $s-2$ pass bounds, and the $s-1$ Green bounds give $N^{-1} 2^{s-2} 2^{s-1} = 2^{2s-3} N^{-1}$. The estimate is not iterated over a coupled chronological simplex by replacing that simplex with rectangles; those exact constrained sums are handled by [Lemma 3.10](#). The mixed cases use [Equation \(3.2.4a\)](#) in the displayed prescribed-gate calculation. Every product contains the original pair of endpoint fronts once, which proves the token assertion. $\square$

Lemma 3.10 (Permutation-summed passive-singleton renewal). *Let M_j be any typed passive copy of the coordinate pure-Weyl multiplier in [Lemma 3.9](#), with arbitrary passive Hilbert-space amplification. Put*

$$V_j = Q M_j Q, \quad A_j = \mathcal{E} M_j \mathcal{T}, \quad \mathfrak{H}_a = \mathcal{G} K^a \mathcal{G}^*.$$

Then

$$\mathcal{T}^* M_j \mathcal{T} = \mathcal{E} M_j \mathcal{E}^* = 0, \quad \mathcal{G}^* V_j \mathcal{G} = -K A_j - A_j^* K. \quad (3.2.10q_1)$$

There is a numerical C such that

$$\sum_{a,b \geq 0} \|\mathfrak{H}_a \mathbf{V}_j \mathfrak{H}_b\| \leq C \frac{\log(n+2)}{\sqrt{n}}. \quad (3.2.10q_2)$$

Consequently, for every fixed $L < \infty$ there is $A_L < \infty$ such that a passive-singleton component with $2 \leq s \leq L \log n$ labelled vertices satisfies, after summing every strict chronological order and every allowed finite-gap simplex,

$$\sum_{\sigma \in S_s} \|\mathcal{X}_{s,\sigma}\| \leq \frac{A_L^s}{N}. \quad (3.2.10q_3)$$

After the proved circuit-anchored root/pass rewrite, its one free translation, and a fixed terminal ribbon gluing, its scalar contribution is at most $A_L^s(m/N)\mathbf{a}_\Gamma$, where $\mathbf{a}_\Gamma$ is the positive projective majorant of the already normalized non-temporal atom. The same conclusion holds for several Haar-separated passive-singleton components after their labelled vertex sets have been fixed, with one factor m/N per component and no interleaving multinomial. This last sentence does not sum over varying component-label decompositions; the physical maximal passive-only components are instead annihilated exactly in [Lemma 3.18](#) below.

Proof. Own-circle invariance of W_i° and the centering identity in [Equation \(3.2.10r\)](#) give, on every passive amplification,

$$\mathcal{T}^* \mathbf{M}_j \mathcal{T} = \frac{1}{N} \sum_i \int \pi(R_i(\theta))^* (W_i^\circ \otimes I) \pi(R_i(\theta)) \frac{d\theta}{2\pi} = 0$$

and $\mathcal{E} \mathbf{M}_j \mathcal{E}^* = 0$. Expand both copies of $\mathbf{Q} = I - \mathcal{E}^* \mathcal{E}$ in $\mathcal{T}^* \mathbf{Q} \mathbf{M}_j \mathbf{Q} \mathcal{T}$. The first and last terms vanish by the two preceding identities, while $\mathcal{E} \mathcal{T} = \mathcal{K} = \mathcal{K}^*$; this proves [Equation \(3.2.10q₁\)](#).

Let

$$F_j = \mathcal{E} \mathbf{M}_j \mathbf{Q}, \quad \eta_n = \frac{n-3}{N(n-1)}.$$

The proof of [Lemma 3.9](#) gives $A_j = F_j \mathcal{G}$ and $\|F_j\| \leq \eta_n^{1/2}$. The exact Kac spectral-gap theorem of Carlen–Carvalho–Loss and Maslen [[CCL03](#); [Mas03](#)], in the present discrete-time normalization, says that the spectrum of $\mathcal{K}$ off its invariant space lies in $[0, 1 - \gamma_n]$, where

$$\gamma_n = \frac{n+2}{2n(n-1)}. \quad (3.2.10q_4)$$

The assertion applies to every amplified finite-dimensional module by restriction of the regular-representation spectral theorem to its matrix coefficients. Since $\mathcal{G}^* \mathcal{G} = I - \mathcal{K}^2$, elementary spectral calculus gives

$$\sum_{a \geq 0} \|\mathcal{G} \mathcal{K}^a\| \leq C \gamma_n^{-1/2}, \quad \sum_{b \geq 0} \|\mathfrak{H}_b\| \leq C \log \frac{2}{\gamma_n}. \quad (3.2.10q_5)$$

For completeness, for $a \geq 1$ the unconstrained maximum of $(1 - \lambda^2)^{1/2} \lambda^a$ is at $\lambda^2 = a/(a+1)$ and is at most $(a+1)^{-1/2}$. For $b \geq 1$ the unconstrained maximum of $(1 - \lambda^2) \lambda^b$ is at $\lambda^2 = b/(b+2)$ and is at most $2/(b+2)$. Split both sums when these maximizers pass $1 - \gamma_n$; the remaining spectral-edge tails are geometric. This proves [Equation \(3.2.10q₅\)](#), including constants independent of the horizon.

Insert [Equation \(3.2.10q₁\)](#) between the two innovation incidences. Using $A_j = F_j \mathcal{G}$ and its adjoint gives

$$\begin{aligned} \mathfrak{H}_a \mathbf{V}_j \mathfrak{H}_b &= -\mathcal{G} \mathcal{K}^{a+1} F_j \mathfrak{H}_b \\ &\quad - \mathfrak{H}_a F_j^* \mathcal{K}^{b+1} \mathcal{G}^*. \end{aligned}$$

Sum the norms over the full orthant and apply Equation (3.2.10q₅). Since

$$\frac{\eta_n}{\gamma_n} = \frac{4(n-3)}{(n-1)(n+2)} \leq \frac{C}{n},$$

this proves Equation (3.2.10q₂). Notice that the full-orthant estimate is taken only after the exact compressed pass identity has been formed. It therefore dominates every coupled finite-gap simplex without replacing that simplex by an operator product of rectangular Green sums.

Fix one strict order of an s -vertex excursion. Its operator is a finite simplex sum of words

$$S^- \mathfrak{H}_{\ell_1} V_2 \mathfrak{H}_{\ell_2} \cdots V_{s-1} \mathfrak{H}_{\ell_{s-1}} S^+.$$

Select the even interior vertices. The resulting $p_s = \lfloor (s-1)/2 \rfloor$ two-gap blocks are disjoint. Take term norms, enlarge only their nonnegative gap sums to the full orthant, and use Equation (3.2.10q₂) on each block. The unselected pass factors have norm at most two. If s is even, the one remaining gap costs at most $C \log(n+2)$ by Equation (3.2.10q₅). The two endpoint fronts cost N^{-1} . Thus, for $s \geq 4$,

$$\|\mathcal{X}_{s,\sigma}\| \leq \frac{C^s \log(n+2)}{N} \left(\frac{C \log(n+2)}{\sqrt{n}} \right)^{p_s}.$$

The case $s = 2$ follows directly from Lemma 3.9; for $s = 3$, apply Equation (3.2.10q₂) to the single internal pass and multiply by the two seed bounds in Equation (3.2.10o). Since $p_s \geq s/4$ for $s \geq 4$, $s! \leq s^s$, and $s \leq L \log n$, the elementary bound

$$\sup_{n \geq 2} \frac{(\log(n+2))^{3/2}}{n^{1/8}} < \infty$$

absorbs the $s!$ strict orders and the displayed extra logarithm into A_L^s . This proves Equation (3.2.10q₃).

There are at most m choices of the component translation. Terminal prescribed-gate gluing preserves $\mathfrak{a}_\Gamma$, proving the scalar claim. Finally, restriction of a global strict time assignment to its Haar-separated components is injective. Dropping the requirement that times belonging to different components be distinct enlarges the nonnegative majorant to the product of the component sums. Hence there is no extra interleaving multinomial, and the proof is complete. This factorization is only for the fixed labelled component sets specified in the statement; no component-set sum has been taken. $\square$

Lemma 3.11 (Typed bracketed-pass debit). *On the noninvariant spectral subspace put $R_z = (I - zK)^{-1}$ for $|z| = 1$, with the Moore–Penrose interpretation at $z = 1$. For $X \in \{\mathcal{D}, \mathcal{G}, \mathcal{L}\}$ put*

$$b_{\mathcal{D}} = 1, \quad b_{\mathcal{G}} = \sqrt{2}, \quad b_{\mathcal{L}} \leq d,$$

where d is the circle Fourier degree. Then

$$\|X R_z Y^*\| \leq b_X b_Y \quad (X, Y \in \{\mathcal{D}, \mathcal{G}, \mathcal{L}\}). \quad (3.2.10q_6)$$

For every typed passive singleton V_j from Lemma 3.10,

$$\boxed{\|X R_z \mathcal{G}^* V_j \mathcal{G} R_z Y^*\| \leq \frac{C b_X b_Y}{\sqrt{n}}.} \quad (3.2.10q_7)$$

Both estimates are unchanged by passive representation amplification and may be inserted, with the displayed typed incidences, in the final scalar gluing of Proposition 2.22.

Proof. The proof of [Lemma 3.4](#) gives

$$X^*X \preceq b_X^2(I - \mathcal{K}), \quad Y^*Y \preceq b_Y^2(I - \mathcal{K}).$$

Douglas factorization and scalar spectral calculus, using $|1 - z\lambda| \geq 1 - \lambda$ for $0 \leq \lambda < 1$, prove [Equation \(3.2.10q6\)](#). On the complement of the invariant space the spectral-gap bound [Equation \(3.2.10q4\)](#) also gives

$$\|XR_z\mathcal{K}\| \leq b_X\gamma_n^{-1/2}, \quad \|\mathcal{G}R_zY^*\| \leq \sqrt{2}b_Y. \quad (3.2.10q8)$$

Indeed the relevant scalar quotients are bounded by $\lambda^2/(1-\lambda) \leq \gamma_n^{-1}$ and $(1-\lambda^2)/(1-\lambda) = 1+\lambda \leq 2$.

Write $F_j = \mathcal{E}M_jQ$. The proof of [Lemma 3.10](#) gives $A_j = F_j\mathcal{G}$ and

$$\|F_j\|^2 \leq \eta_n := \frac{n-3}{N(n-1)}, \quad \mathcal{G}^*V_j\mathcal{G} = -\mathcal{K}A_j - A_j^*\mathcal{K}.$$

Insert the latter identity between XR_z and R_zY^* . The first term is bounded, using [Equation \(3.2.10q8\)](#), by $b_Xb_Y\sqrt{2\eta_n/\gamma_n}$; the adjoint estimates, with z replaced by $\bar{z}$, give the same bound for the second term. Since

$$\frac{\eta_n}{\gamma_n} = \frac{4(n-3)}{(n-1)(n+2)} \leq \frac{4}{n},$$

[Equation \(3.2.10q7\)](#) follows. The calculation acts only on the star factor. The prescribed-gate gluing keeps all other packet/root maps in their typed slots, so no retained-output norm has been used. $\square$

Lemma 3.12 (Shared-resolvent consecutive-pass gain). *Use the notation of [Lemma 3.11](#). Let $V_1, \dots, V_s$ be typed passive singleton multipliers, possibly with passive representation amplification, and put*

$$B_j = \mathcal{G}^*V_j\mathcal{G}.$$

Then, with the same unit-circle parameter in every residual gap,

$$\boxed{\|XR_zB_1R_zB_2 \cdots R_zB_sR_zY^*\| \leq b_Xb_Y \left(\frac{C}{\sqrt{n}}\right)^s.} \quad (3.2.10q9)$$

The estimate keeps the entire consecutive run and its shared resolvent intact. It is stable under ancillary amplification and prescribed-gate terminal gluing.

Proof. For the j th multiplier write

$$F_j = \mathcal{E}M_jQ, \quad A_j = F_j\mathcal{G}, \quad \|F_j\|^2 \leq \eta,$$

where either $\eta = \eta_n$ from [Lemma 3.10](#) or the better full-deposit value $\eta_C \leq N^{-1}$. In both cases

$$\frac{\eta}{\gamma_n} \leq \frac{4}{n}. \quad (3.2.10q10)$$

The exact coboundary identity is

$$B_j = -\mathcal{K}F_j\mathcal{G} - \mathcal{G}^*F_j^*\mathcal{K}. \quad (3.2.10q11)$$

Let Q_0 be the orthogonal projection off the invariant subspace and put

$$\mathcal{K}_0 = Q_0\mathcal{K}Q_0, \quad \tilde{A}_j = Q_0F_j\mathcal{G}Q_0.$$

Since $\mathcal{G}Q_0 = \mathcal{G}$, $Q_0\mathcal{G}^* = \mathcal{G}^*$, and Q_0 commutes with $\mathcal{K}$, one has $B_j = Q_0B_jQ_0$ and the projected exact identity

$$B_j = -\mathcal{K}_0\tilde{A}_j - \tilde{A}_j^*\mathcal{K}_0. \quad (3.2.10q_{11a})$$

Moreover $0 \leq \mathcal{K}_0 \leq 1 - \gamma_n$ and $\|Q_0F_j\| \leq \sqrt{\eta}$. Thus every orientation below lives entirely on the noninvariant spectral subspace; in particular no isolated $\mathcal{K}R_z\mathcal{K}$ factor sees the invariant eigenvalue.

Expand Equation (3.2.10q_{11a}) in the product Equation (3.2.10q₉) into its 2^s left/right orientations. Besides Equation (3.2.10q₆), scalar spectral calculus gives

block $XR_z\mathcal{K}_0, \mathcal{K}_0R_zY^*$ $\mathcal{G}R_z\mathcal{K}_0, \mathcal{K}_0R_z\mathcal{G}^*$ $\mathcal{G}R_z\mathcal{G}^*$ $\mathcal{K}_0R_z\mathcal{K}_0$	bound after removing b_X, b_Y at the ends $\gamma_n^{-1/2}$ $C\gamma_n^{-1/2}$ C $C\gamma_n^{-1}$.	$(3.2.10q_{12})$
---	---	------------------

Indeed $|1 - z\lambda| \geq 1 - \lambda$ and $\mathcal{G}^*\mathcal{G} = I - \mathcal{K}^2$ reduce these four lines to the corresponding scalar multipliers on $0 \leq \lambda \leq 1 - \gamma_n$.

Call the first summand in Equation (3.2.10q₁₁) L and the second R . The left boundary costs a half power of γ_n^{-1} exactly when the first letter is L , and the right boundary does so exactly when the last letter is R . At an internal interface the exponents for $(L, L), (L, R), (R, L), (R, R)$ are respectively

$$\frac{1}{2}, \quad 0, \quad 1, \quad \frac{1}{2}. \quad (3.2.10q_{13})$$

For every word in $\{L, R\}^s$, the two boundary contributions plus all interface contributions in Equation (3.2.10q₁₃) sum exactly to $s/2$; this is the telescoping identity obtained by assigning $1/2$ to each letter and moving a half unit across every R, L transition. Each word also contains exactly s factors F_j or F_j^* . Hence its norm is at most

$$b_X b_Y C^s \eta^{s/2} \gamma_n^{-s/2}.$$

Summing the 2^s orientations and using Equation (3.2.10q₁₀) proves Equation (3.2.10q₉). Every estimate acts only on the common star factor, proving the amplification and gluing assertions. $\square$

Lemma 3.13 (Anchor-averaged Taylor sums retain a unit-circle debit). *Fix $L < \infty$. Let $r \leq L \log(n+2)$, $0 \leq \Delta \leq r$, and $0 \leq d \leq r + B_0$. For every physical translation a in a set of at most $m \leq cN$ translations, let $M(a) \in \{0, \dots, m\}$ be its remaining gap budget, with distinct translations giving distinct budgets, and let*

$$\mathcal{X}_a^{(M(a))} = \sum_{q=0}^{M(a)} [z^q] \mathcal{F}_a(z).$$

Suppose that, for one number $H \geq 0$ independent of a ,

$$\sup_{|z|=1} |\mathcal{F}_a(z)| \leq H n^{-\Delta} \quad (3.2.10q_{14})$$

and, when $d \geq 1$,

$$|[z^q] \mathcal{F}_a(z)| \leq H \binom{q+d-1}{d-1} (1 - \gamma_n)^q, \quad q \geq 0. \quad (3.2.10q_{15})$$

For $d = 0$ interpret $\mathcal{F}_a$ as constant. Then

$$\boxed{\sum_a |\mathcal{X}_a^{(M(a))}| \leq C_L c N H n^{-\Delta}.} \quad (3.2.10q_{16})$$

The conclusion is anchor-averaged: it does not assert a dimension-free maximal-partial-sum estimate for one prescribed budget. It remains valid for a final-scalar prescribed-gate contraction, provided [Equations \(3.2.10q₁₄\) to \(3.2.10q₁₅\)](#) hold after that contraction.

Proof. The case $d = 0$ is immediate, so assume $d \geq 1$. Choose

$$M_0 = C'_L \gamma_n^{-1} \log^2(n+2), \quad (3.2.10q_{17})$$

where C'_L will be fixed below. The L^1 norm of the scalar Dirichlet kernel and [Equation \(3.2.10q₁₄\)](#) give

$$|\mathcal{X}_a^{(M)}| \leq C \log(M+2) H n^{-\Delta}.$$

Because the budgets are distinct, the sum over $M \leq M_0$ is at most

$$C(M_0+1) \log(M_0+2) H n^{-\Delta} \leq C_L N H n^{-\Delta}. \quad (3.2.10q_{18})$$

Here $\gamma_n^{-1} \leq 2n$ and $n \log^3(n+2) \leq CN$; increasing C_L covers the finitely many small dimensions.

For $M > M_0$, absolute convergence of [Equation \(3.2.10q₁₅\)](#) gives

$$\mathcal{X}_a^{(M)} = \mathcal{F}_a(1) - \sum_{q>M} [z^q] \mathcal{F}_a(z).$$

The sum of the first terms over all translations is at most $c N H n^{-\Delta}$. The standard negative-binomial estimate gives

$$\sum_{M>M_0} \sum_{q>M} \binom{q+d-1}{d-1} (1-\gamma_n)^q \leq C \gamma_n^{-d-1} e^{-\gamma_n M_0/2}. \quad (3.2.10q_{19})$$

Indeed, for $M_0 \geq 4d/\gamma_n$, split off $e^{-\gamma_n q/2}$ and sum the remaining negative-binomial series; the sum in M contributes one further factor $C \gamma_n^{-1}$. Now $d \leq L \log(n+2) + B_0$ and $\Delta \leq L \log(n+2)$. Hence

$$\gamma_n^{-d-1} n^\Delta \leq \exp\{C_L \log^2(n+2)\},$$

whereas the last exponential in [Equation \(3.2.10q₁₉\)](#) is $\exp\{-\frac{1}{2} C'_L \log^2(n+2)\}$. Taking C'_L sufficiently large therefore makes the tail sum at most $C_L N n^{-\Delta}$. Multiplication by H , together with [Equation \(3.2.10q₁₈\)](#), proves [Equation \(3.2.10q₁₆\)](#). $\square$

Lemma 3.14 (Physical translations have distinct simplex budgets). *After equal-time fusion, fix one relative weak order with $h \geq 1$ ordered supernodes. Reparametrize its physical translation orbit by the first occupied time $a = t_1$. If $t_{j+1} - t_j = 1 + g_j$ with $g_j \geq 0$, and q is the total slack recorded by the common one-variable Green series, admissibility in $\{1, \dots, m\}$ is*

$$a + (h-1) + q \leq m.$$

Thus its Taylor budget is

$$M(a) = m - h + 1 - a, \quad M(a+1) = M(a) - 1. \quad (3.2.10q_{20})$$

In particular distinct physical translations have distinct budgets. If a root time was initially used to name the orbit, replacing it by t_1 is a bijective reparametrization and changes no local state or multiplicity.

Proof. The $h - 1$ strict steps are compulsory, while every additional empty time is one unit of a gap g_j and hence one power of the common generating variable. The last occupied time is $a + (h - 1) + q$; requiring it to be at most m proves the formula. $\square$

Definition 3.15 (Fused chronological supernode). After a weak chronological order has been fixed, a *supernode* is one equality block: all occurrences assigned to one physical time after the complete same-time rule has been applied. It is *hard* if the equality block contains at least two occurrences, or if its fused tensor contains a nonsingleton packet, the time-indexed $-2N^{-1}W_i$ Jacobi child, or a special root. In particular, every surviving same-time nonsingleton is hard, whether or not its occurrences are column-connected. Otherwise the block consists of exactly one occurrence and is a passive singleton supernode carrying one of the four star statuses defined after Equation (3.2.10n). Ordered supernodes are always ordered by their actual physical times; no fragment of one equality block is counted as a second node.

Lemma 3.16 (Monomer–dimer resource ledger). Put $X_0 = D_m - (\gamma + 1)I$. For every $s \geq 0$,

$$Q_s(D_m) = \sum_{\pi \in \mathfrak{T}_s} (-\gamma)^{a(\pi)} X_0^{r(\pi)}, \quad 2a(\pi) + r(\pi) = s, \quad (3.2.10i)$$

where $\mathfrak{T}_s$ is the set of monomer–dimer tilings of an interval; in particular $|\mathfrak{T}_s| \leq 2^s$. In a term of the squared degree- k differentiated shell, let a be the total number of dimers and let r count all nonconstant slots, including in each replica the differentiated deposit and the endpoint root. Then

$$a + \frac{r}{2} = k + 1. \quad (3.2.10j)$$

Fix one ribbon contraction and let p be its number of centered pairing-connected packet components, including the single paired root component. Every such component contains at least two nonconstant slots, so

$$p \leq r/2, \quad a + p \leq k + 1. \quad (3.2.10k)$$

All constants in these statements are independent of $c \geq 1$.

Proof. Equation (3.2.10i) is the recurrence $Q_{s+1} = X_0 Q_s - \gamma Q_{s-1}$, with a monomer for the first term and a dimer for the second. In one differentiated replica, the two Q factors in Equation (3.2.7) have total recurrence length $k - 1$; adding the differentiated deposit and the root gives $a + r/2 = (k + 1)/2$. Adding the replicas proves Equation (3.2.10j).

Centering removes a one-occurrence passive packet, while the output inner product joins the two derivative marks into one paired root packet. Thus $p \leq r/2$, and Equation (3.2.10j) gives Equation (3.2.10k). A Haar separator can split a temporal word into several residual components, but it cannot split one complete physical-time packet into distinct hard vertices: equal-time fusion puts that packet in one supernode, and any fragments joined only through an exterior port remain one packet resource. Hence distinct packet endpoints come from distinct packets. By Lemma 3.18, each nonzero component is hard and receives its own distinct packet, time-indexed linear-Jacobi, or coupled marked-root endpoint resource. Distinct resources consume disjoint pairs of nonconstant half-slots, so their number q satisfies $q \leq r/2$. Consequently $\gamma^a c^q \leq c^{a+r/2} = c^{k+1}$. No equality partition or short-gap graph is part of this algebraic ledger. $\square$

Lemma 3.17 (Exact Haar–residual gap decomposition). Let V be any finite-dimensional orthogonal $\text{SO}(n)$ -module, let $\mathcal{K}_V = N^{-1} \sum_i P_{i,V}$, and let Π_V be the orthogonal projection onto $V^{\text{SO}(n)}$. With $Q_V = I - \Pi_V$, for every $\ell \geq 0$,

$$\mathcal{K}_V^\ell = \Pi_V + Q_V \mathcal{K}_V^\ell Q_V. \quad (3.2.10s)$$

Consequently every empty chronological gap in a fixed reflected ribbon child is exactly either a Haar separator or a residual gap. Expanding all gaps produces at most 2^h children when h physical times are used. End components obey the same convention after inserting a common auxiliary Haar conjugation in the conjugation-invariant squared-shell scalar.

Proof. The operator $\mathcal{K}_V$ is self-adjoint, fixes the invariant subspace, and preserves its orthogonal complement. Hence $\Pi_V \mathcal{K}_V^\ell = \Pi_V$ and the two off-diagonal blocks vanish, which proves Equation (3.2.10s). Choosing one of the two displayed summands at each gap gives at most 2^h children. The squared-shell scalar is unchanged by simultaneous conjugation of the two replicas; averaging that common conjugation is precisely Π_V , proving the endpoint assertion. $\square$

Lemma 3.18 (Haar-boundary annihilation of passive components). *Let V_- and V_+ be orthogonal $\mathrm{SO}(n)$ -modules. On their star dilations write $\mathcal{T}_\pm, \mathcal{E}_\pm, \mathbf{Q}_\pm$ and $\mathcal{G}_\pm = \mathbf{Q}_\pm \mathcal{T}_\pm$ for the maps in Equation (3.2.2). Let $\mathbf{M} : (V_-)_* \rightarrow (V_+)_*$ be a typed passive singleton, so that*

$$(\mathbf{M}h)(i, \theta) = M_i h(i, \theta), \quad \rho_+(R_i(\theta))^* M_i \rho_-(R_i(\theta)) = M_i, \quad \sum_i M_i = 0. \quad (3.2.10s_1)$$

Put $\mathbf{S}_{+,-}^+ = \mathbf{Q}_+ \mathbf{M} \mathcal{E}_-^*$. If Π_- is the projection onto $V_-^{\mathrm{SO}(n)}$, then

$$\boxed{\mathcal{G}_+^* \mathbf{S}_{+,-}^+ \Pi_- = 0.} \quad (3.2.10s_2)$$

Writing $\mathbf{S}_{-,+}^- = \mathcal{E}_- \mathbf{M}^* \mathbf{Q}_+$, applying the same statement to $\mathbf{M}^*$ gives the oppositely oriented identity $\Pi_- \mathbf{S}_{-,+}^- \mathcal{G}_+ = 0$. Consequently every maximal residual temporal component made only of passive singleton supernodes is identically zero, term by term in its weak order and finite residual gaps. This includes either end component under the common auxiliary Haar projection of Lemma 3.17.

Proof. For an invariant vector $x \in V_-$, every coordinate circle fixes x ; hence, directly from Equation (3.2.2),

$$\mathcal{T}_- \Pi_- = \mathcal{E}_-^* \Pi_-. \quad (3.2.10s_3)$$

The centering and own-circle conditions in Equation (3.2.10s₁) give

$$\mathcal{E}_+ \mathbf{M} \mathcal{E}_-^* = 0, \quad \mathcal{T}_+^* \mathbf{M} \mathcal{T}_- = 0. \quad (3.2.10s_4)$$

Indeed both expressions equal $N^{-1} \sum_i M_i$ after, in the second one, commuting M_i through its own coordinate-circle actions. If $\mathbf{J}_+ = \mathcal{E}_+^* \mathcal{E}_+$, the first equality implies $\mathbf{J}_+ \mathbf{M} \mathcal{E}_-^* = 0$, and therefore

$$\begin{aligned} \mathcal{G}_+^* \mathbf{S}_{+,-}^+ \Pi_- &= \mathcal{T}_+^* \mathbf{Q}_+ \mathbf{M} \mathcal{E}_-^* \Pi_- \\ &= \mathcal{T}_+^* \mathbf{M} \mathcal{E}_-^* \Pi_- = \mathcal{T}_+^* \mathbf{M} \mathcal{T}_- \Pi_- = 0. \end{aligned}$$

This proves Equation (3.2.10s₂). The argument for $\mathbf{M}^*$ proves the reverse orientation and does not identify either module with its dual. It is therefore stable under duality, rail reversal, passive amplification, and fixed equivariant Brauer or port maps; for an equivariant $B : V_1 \rightarrow V_2$ one uses $B \Pi_{V_1} = \Pi_{V_2} B \Pi_{V_1}$. Passive aggregate-Ricci factors are included only in their covariant curried form. Namely,

$$\mu : \mathrm{Sym}_0(H) \otimes W \longrightarrow W, \quad \mu(B, x) = X_B x,$$

is $\mathrm{SO}(n)$ -equivariant because $\rho_W(g) X_B x = X_{g B g^*} \rho_W(g) x$. We retain B as a vector in the complete coefficient module and apply the fixed intertwiner μ ; we do not freeze X_B as a non-equivariant map next to Π .

For $s \geq 2$, an oriented passive residual word has, up to such equivariant typed maps and adjoints, the form

$$\text{IIS}^-(\mathcal{G}\mathcal{K}^{\ell_1}\mathcal{G}^*)\mathbf{V}_2 \cdots \mathbf{V}_{s-1}(\mathcal{G}\mathcal{K}^{\ell_{s-1}}\mathcal{G}^*)\mathbf{S}^+\Pi. \quad (3.2.10s_5)$$

Its rightmost residual block contains $\mathcal{G}^*\mathbf{S}^+\Pi = 0$; the reversed word is killed by the adjoint identity. Thus every fixed gap tuple vanishes, and so does every finite coupled gap sum. For $s = 1$ the claim is $\mathcal{E}\mathbf{M}\mathcal{E}^* = 0$.

Every interior component not containing the paired marked root has a selected Haar separator on its root-facing side. At the two outer ends, [Lemma 3.17](#) supplies one common auxiliary invariant projection; one invariant side is enough, so neither independence nor a tensor factorization of that projection is used. If no Haar separator is selected, there is one temporal component and it contains the coupled pure-Weyl root or the intact reflected aggregate-Ricci root, so it is not passive-only. A $\mathcal{D}/\mathcal{L}$ endpoint, a nonsingleton same-time packet, the time-indexed Jacobi child $-2N^{-1}W_i$, or a current-plane aggregate-Ricci interface is likewise hard and lies outside the assertion. These cases exhaust the exact Haar/residual decomposition. $\square$

3.3 Root-local and packetless aggregate-Ricci calculus

3.3.1 Root evacuation, deposits, and reflected Ricci coupons

Lemma 3.19 (Typed dirty-root evacuation). *Fix a fresh plane i and let $\mathbb{T}_i$ be its coordinate circle. Let $\mathcal{S}_i(\mathbf{X}) : V_{\text{in}} \rightarrow V_{\text{out}}$ be a complete same-time physical supernode made from current deposits, their centered pure-Weyl gates, normalized Brauer maps, and physical root/pass maps, with all non-current pieces retained once as typed gaps $X_j : V_j^- \rightarrow V_j^+$. Put*

$$\delta_i X_j = d\rho_{V_j^+}(a_i)X_j - X_j d\rho_{V_j^-}(a_i).$$

Then the exact identity

$$\boxed{d\rho_{V_{\text{out}}}(a_i)\mathcal{S}_i(\mathbf{X}) - \mathcal{S}_i(\mathbf{X})d\rho_{V_{\text{in}}}(a_i) = \sum_j \mathcal{S}_i(X_1, \dots, \delta_i X_j, \dots, X_g)} \quad (3.3.1u)$$

holds before the plane sum and before either replica is estimated. Therefore a marked action entering the supernode exits at one named external leg or gap, or vanishes at a_i or after complete scalar closure; it is never duplicated. If it passes immediately through W_i° , it acquires the exact scalar

$$b_n = -\frac{n-3}{(n-1)(n-2)}, \quad A_i W_i^\circ = W_i^\circ A_i = b_n A_i. \quad (3.3.1v)$$

The identity is stable under every gap substitution allowed by [Lemma 2.5](#).

Proof. Every current deposit is fixed by its own circle, its pure-Weyl projection is fixed by equivariance, every normalized Brauer map intertwines the diagonal action, and the physical maps on the two sides of each root/pass table are equal equivariant maps. Hence

$$\rho_{V_{\text{out}}}(g)\mathcal{S}_i(\mathbf{X})\rho_{V_{\text{in}}}(g)^{-1} = \mathcal{S}_i(g \cdot X_1, \dots, g \cdot X_g), \quad g \in \mathbb{T}_i.$$

Differentiate at the identity. Multilinearity and the product rule give [Equation \(3.3.1u\)](#). Expanding the input and output infinitesimal actions over their tensor factors moves one selected root action to the other external factors with the usual dual-orientation signs. Internal actions cancel in pairs; $A_i a_i = [a_i, a_i] = 0$; and the total action on a closed invariant scalar is zero. This proves

the evacuation and nonduplication assertions. Because Equation (3.3.1u) is an equality of typed multilinear maps, contextual gap substitution preserves it.

For Equation (3.3.1v), $\text{ran } A_i$ is the span of coordinate bivectors meeting plane i in one endpoint, and A_i kills its orthogonal complement. On that range $P_i = 0$ and X_{B_i} from Equation (2.6.2) has eigenvalue $(n-4)/[n(n-2)]$. Thus

$$W_i^\circ = P_i - N^{-1}I - X_{B_i} = -\frac{n-3}{(n-1)(n-2)}I$$

on $\text{ran } A_i$. The displayed two-sided identity follows. $\square$

Remark 3.20 (Why Green gaps are not eliminated sequentially). Fixing all gaps except one does turn that gap into a bounded gate XR_sY^* . It does *not* justify repeating the operation along a chronological component: after the first summation, the upper limits of the remaining gaps depend on the variable already summed, so the domain is a coupled simplex rather than a product of intervals. Replacing the successive gates by their individual norms loses precisely this dependence. We therefore make no sequential finite-Green lemma. The live argument keeps all gaps of a component in one generating function and estimates the exact Taylor partial sum in Lemma 3.39; the passive-only case is handled directly by Lemma 3.10.

Lemma 3.21 (Full-centered deposit calculus). *Put*

$$C_i = P_i - N^{-1}I_W, \quad P_i = a_i a_i^*.$$

Then

$$C_i = C_i^*, \quad \sum_i C_i = 0, \quad \rho(R_i(\theta))^* C_i \rho(R_i(\theta)) = C_i, \quad \|C_i\|_{\text{op}} \leq 1, \quad (3.3.1w_1)$$

and

$$\sum_i C_i^2 = (1 - N^{-1})I_W. \quad (3.3.1w_2)$$

Consequently every assertion of Lemma 3.9, Lemma 3.10, Lemma 3.11, and Lemma 3.18 remains valid with W_i° replaced by C_i and with the better seed parameter

$$\eta_C = \left\| \frac{1}{N} \sum_i C_i^2 \right\| = \frac{N-1}{N^2} \leq \frac{1}{N}. \quad (3.3.1w_3)$$

For $q \geq 1$ there are scalars α_q, β_q such that

$$C_i^q = \alpha_q C_i + \beta_q I, \quad |\alpha_q| \leq 2, \quad \beta_1 = 0, \quad |\beta_q| \leq \frac{3}{N} \quad (q \geq 2). \quad (3.3.1w_4)$$

More generally, for $2 \leq q \leq L \log n$ and all sufficiently large n ,

$$\left\| \frac{1}{N} \sum_i C_i^{\otimes q} \right\|_{\text{op}} \leq \frac{2}{N}. \quad (3.3.1w_5)$$

After latest-pivot exposure, the same bound, with A_L^q/N on the right, survives powers on individual rails and fixed flips, dualities, and normalized Brauer maps. Thus a fused nonroot full-deposit block is either one passive singleton, an explicit scalar child of size A_L^q/N , or a hard intact block carrying one A_L^q/N resource. If an intact rank-one pinch remains after latest-pivot exposure and Ward evacuation, so that its exterior map $T \in \text{End}(W)$ is fixed with respect to the fresh plane, then

$$\left\| \frac{1}{N} \sum_i P_i T P_i \right\|_{\text{op}} \leq \frac{1}{N} \|T\|_{\text{op}}. \quad (3.3.1w_6)$$

Proof. The projections P_i are mutually orthogonal and sum to I_W . This proves centering, self-adjointness, own-circle invariance, and

$$\sum_i (P_i - N^{-1}I)^2 = I - 2N^{-1}I + N^{-1}I = (1 - N^{-1})I.$$

Inspection of the proofs of the four cited lemmas shows that their only coefficient inputs are self-adjointness, own-circle invariance, centering, a uniform operator bound, and the seed estimate $\|N^{-1} \sum_i M_i^2\|$. Equations (3.3.1w₁) to (3.3.1w₃) therefore give the stated extension. In particular the exact coboundary identity is unchanged and $\eta_C/\gamma_n \leq 4/(n+2)$, so every bracketed pass still gains $C/\sqrt{n}$.

The two eigenvalues of C_i are $u = 1 - N^{-1}$ and $v = -N^{-1}$, with $u - v = 1$. Hence $\alpha_q = u^q - v^q$ and $\beta_q = uv^q - vu^q$, which proves Equation (3.3.1w₄). In the tensor Kac basis, the eigenvalue in Equation (3.3.1w₅) at $a_{j_1} \otimes \cdots \otimes a_{j_q}$ is

$$\frac{1}{N} \sum_i \prod_{\ell=1}^q (\mathbf{1}_{\{i=j_\ell\}} - N^{-1}).$$

At most q summands use an index occurring among the j_ℓ ; their total absolute value is at most $1 + q/N + O(q^2/N^2)$, while the unused indices contribute at most N^{1-q} . This proves Equation (3.3.1w₅) in the stated range. Reduce repeated powers on one rail by Equation (3.3.1w₄). The all-identity child then contains a β_q and hence N^{-1} , the exactly-one- C child vanishes by centering, and every child with at least two centered factors is covered by Equation (3.3.1w₅). Contractive flips, dualities, and normalized Brauer maps preserve the bound. Finally $P_i T P_i = \langle a_i, T a_i \rangle P_i$; orthogonality of the P_i proves Equation (3.3.1w₆). $\square$

Corollary 3.22 (External zero-gap full-deposit environment). *Let a full-centered letter C_i occur as an external typed singleton immediately adjacent to a complete paired-marked or literal- W aggregate-Ricci block. Keep the latter block and every joint context intact. On the plane-star decomposition $\text{ran } J \oplus \text{ran } Q$, the singleton multiplication map has the exact form*

$$\begin{pmatrix} 0 & S^- \\ S^+ & V \end{pmatrix}, \quad \|S^-\| = \|S^+\| \leq N^{-1/2}, \quad \|V\| \leq 1. \quad (3.3.1w_{6a})$$

Consequently the unsplit zero-gap environment has unit operator/projective cost, while any exposed seed has its displayed $N^{-1/2}$ debit.

Proof. The constant-to-constant block is $N^{-1} \sum_i C_i = 0$. The two seed norms are bounded by the square root of $\eta_C \leq N^{-1}$ in Equation (3.3.1w₃), and V is an orthogonal compression of multiplication by C_i , whose operator norm is at most one by Equation (3.3.1w₁). Tensoring with, or projectively substituting, the already fixed joint context changes none of these bounds. $\square$

Lemma 3.23 (Plane-blind closed packets at an aggregate-Ricci root). *Fix a complete orthogonal Weingarten atom (p, q, r) before opening any packet front. Let P be an external q -component assigned wholly to one aggregate-Ricci root time, and suppose its s coefficient occurrences all carry*

$$C_i = |a_i\rangle\langle a_i| - N^{-1}I_W.$$

The aggregate-Ricci passive boundary has no q -dart, so the coefficient contraction of P is a closed scalar, denoted $Z_{P,q}(i)$. It satisfies

$$\boxed{Z_{P,q}(i) = z_{P,q} \text{ for every coordinate plane } i, \quad |z_{P,q}| \leq 4^s.} \quad (3.3.1w_{6b})$$

If P has one occurrence, then $z_{P,q} = 0$. If it is nonzero and has at least two occurrences, then $c = c^*(P) \geq 2$, and replacing its bare carrier n^c by the closed scalar costs at most

$$A^s n^{-c} \quad (3.3.1w_{6c})$$

relative to the bare Euler majorant.

Proof. At fixed (p, q) the row and coefficient indices are disjoint. In one adjoint occurrence,

$$Y_u[(a_u, b_u), (a_{u+1}, b_{u+1})] = \sum_{j_u, k_u, l_u, h_u} U_{j_u a_u} U_{k_u b_u} (\mathbf{C}_i)_{j_u k_u, l_u h_u} U_{l_u a_{u+1}} U_{h_u b_{u+1}}. \quad (3.3.1w_{6d})$$

The p -deltas contract only the displayed row/trace letters, whereas the q -deltas contract every index of every copy of $\mathbf{C}_i$. Thus no plane-dependent row map remains after the closed q -contraction.

Choose a signed coordinate permutation g carrying plane i to plane j . On W ,

$$\rho_W(g) a_i = \pm a_j, \quad \rho_W(g) \mathbf{C}_i \rho_W(g)^* = \mathbf{C}_j.$$

Every delta contraction, normalized wedge map, signed leg permutation, reversal, and structural Brauer/Bianchi projection in the closed physical coefficient scalar is orthogonally natural. Simultaneous conjugation by g therefore proves $Z_{P,q}(i) = Z_{P,q}(j)$. This statement is made for the complete closed contraction, before a projective-front atom is selected; an individual atom of such an opened presentation need not be equivariant.

For the numerical bound, the normalized two-leg realization $\iota_\wedge a_i$ has Hilbert projective norm at most $\sqrt{2}$. Consequently the four-leg realization of $|a_i\rangle\langle a_i|$ has projective norm at most two. Also

$$I_W = \sum_{e=1}^N |a_e\rangle\langle a_e|$$

has normalized four-leg projective norm at most $2N$, so $N^{-1}I_W$ has norm at most two. Hence the four-leg projective norm of $\mathbf{C}_i$ is at most four. Tensoring the s copies and contracting all q -edges by Hilbert pairings proves $|z_{P,q}| \leq 4^s$.

For one box the three possible restrictions of q give respectively zero by wedge skewness, $\text{Tr}_W \mathbf{C}_i$, and $-\text{Tr}_W \mathbf{C}_i$; all vanish. Finally the matching argument in the proof of [Lemma 3.41](#) gives $c \geq 2$ for every nonzero multibox packet. Comparing the closed bound 4^s with the bare factor n^c proves [Equation \(3.3.1w_{6c}\)](#). $\square$

Lemma 3.24 (Aligned current-plane strings preserve the Ricci row square). *Let a consecutive string of full-centered current-plane letters be immediately adjacent to a named incidence of the complete aggregate-Ricci source, with no intervening plane-dependent or noncommuting gate. Keep the string unsplit through the fresh-plane sum. Then the distinct- and coincident-time joint four-leg bounds in [Lemma 3.25](#) below remain valid, with only a numerical change of constant.*

Proof. Let P_i denote the coordinate-circle average on W . Put

$$\kappa_W = \frac{n^2 - 5n + 8}{n(n-1)}, \quad \frac{1}{N} \sum_i P_i = \kappa_W I_W, \quad R_i = P_i - \kappa_W I_W,$$

and retain $P_i = |a_i\rangle\langle a_i|$ for the rank-one deposit. Own-circle invariance gives the exact identities

$$A_i \mathbf{C}_i = \mathbf{C}_i A_i = -N^{-1} A_i, \quad R_i \mathbf{C}_i = \mathbf{C}_i R_i = (1 - \kappa_W) P_i - N^{-1} R_i, \quad (3.3.1w_{6e})$$

and therefore

$$\|R_i \mathbf{C}_i\| \leq (1 - \kappa_W) + N^{-1} \leq Cn^{-1}, \quad \frac{1}{N} \sum_i R_i \mathbf{C}_i = \frac{1 - \kappa_W}{N} I_W. \quad (3.3.1w_{6f})$$

Thus an aligned multiplier improves every circle row. On the centered innovation row, with $\mathbf{e} = N^{-1/2}(E_i - \bar{E})_i$,

$$\mathcal{G}^* \mathbf{M}_C \mathbf{e} = \frac{1}{N} \sum_i R_i \mathbf{C}_i E_i - \frac{1 - \kappa_W}{N} \bar{E}. \quad (3.3.1w_{6g})$$

The first term obeys the same coefficient-row estimate as the unmodified innovation after R_i is replaced by $R_i \mathbf{C}_i$ and Equation (3.3.1w_{6f}) is used. For the second, the exact root identity gives $\bar{E} = -L_V + H_B$; the row-square estimates for L_V and H_B proved in Lemma 3.25 are $O(N^{-1}\|B\|_F^2)$. The plane-independent residue is smaller as well, since

$$\mathcal{T}^* \mathbf{M}_C \mathcal{E}^* H_B = \frac{1}{N} \sum_i P_i \mathbf{C}_i H_B = \frac{1 - \kappa_W}{N} H_B.$$

These calculations are row-square inequalities and hence survive the physical doubled-reversal Haagerup pairing.

At coincident reflected root times the common-plane source is, up to bounded status coefficients, the sum of the five row families

$$(Y_i, a_i), \quad (I, Y_i a_i), \quad (X_B A_i, a_i), \quad (A_i, X_B a_i), \quad (-I, A_i X_B a_i), \quad (3.3.1w_{6g'})$$

where (L_i, v_i) denotes $Z \mapsto L_i Z v_i$. Composing an adjacent $\mathbf{C}_i$ on either physical side preserves their row-square bounds. For the first, second, and fifth families use $\|\mathbf{C}_i\| \leq 1$; for the third and fourth, and for their reflected orientations, use $A_i \mathbf{C}_i = \mathbf{C}_i A_i = -N^{-1} A_i$. Hence the combined (i, r) row square remains $O(n^{-1}\|B\|_F^2)$, and Haagerup's inequality gives the coincident four-leg bound.

Finally $\mathbf{C}_i$ has only two eigenvalues, so for $q \geq 1$

$$\mathbf{C}_i^q = \alpha_q \mathbf{C}_i + \beta_q I, \quad |\alpha_q| \leq 2, \quad |\beta_q| \leq 3N^{-1} \quad (q \geq 2).$$

Every consecutive string reduces to the identity case and the single aligned case. The reflected orientations follow by adjoining Equation (3.3.1w_{6e}). Notice that bounding the innovation block $\mathbf{V}$ by one before this calculation would discard the aligned cancellation; the string is kept intact until the displayed identities are applied. $\square$

Lemma 3.25 (Joint four-leg reflected Ricci coupon). *Let*

$$\mathcal{R}^{\text{in}} = W_+^{\text{in}} \otimes W_-^{\text{out}}, \quad \mathcal{R}^{\text{out}} = W_+^{\text{out}} \otimes W_-^{\text{in}} \quad (3.3.1w_{12a})$$

be the physical doubled-reversal boundary. Let $B, B' \in \text{Sym}_0(H)$ and $|\alpha|, |\beta|, |\alpha'|, |\beta'| \leq 1$. The complete distinct-time aggregate-Ricci sources, with these statuses, and every finite or unit-circle literal- W Green kernel define an operator

$$\mathcal{A}_{\text{Ric}, z}(B, B') : \mathcal{R}^{\text{in}} \longrightarrow \mathcal{R}^{\text{out}}$$

satisfying

$$\|\mathcal{A}_{\text{Ric}, z}(B, B')\|_{\text{op}} \leq \frac{C}{n} \|B\|_F \|B'\|_F, \quad |z| = 1, \quad (3.3.1w_{12b})$$

with the same bound for every finite Green kernel. At coincident reflected root times the complete shared-plane source defines an operator $\mathcal{A}_{\text{Ric},=(B,B')}$ obeying the same bound. Consequently every physically typed joint complement

$$K \in \mathcal{S}_1(\mathcal{R}^{\text{in}}, \mathcal{R}^{\text{out}})$$

obeys

$$\boxed{|\text{Tr}(K^* \mathcal{A}_{\text{Ric},z}(B, B'))| \leq \frac{C}{n} \|B\|_F \|B'\|_F \|K\|_1.} \quad (3.3.1w_{12c})$$

At coincident times replace $\mathcal{A}_{\text{Ric},z}$ in this display by $\mathcal{A}_{\text{Ric},=}$. The assertion is joint: no replica-projective factorization of K is assumed. It remains true after tensoring the root operator by an ancillary identity which lies outside the literal- W Green rail.

Proof. We give the coefficient-row proof, including the nonsymmetric-slot residue. On the adjoint module let P_i be the coordinate-circle average and put

$$\frac{1}{N} \sum_i P_i = \kappa_W I_W, \quad \kappa_W = \frac{n^2 - 5n + 8}{n(n-1)}, \quad 1 - \kappa_W = \frac{2(n-2)}{N}, \quad R_i = P_i - \kappa_W I_W. \quad (3.3.1w_{12d})$$

Since $P_i^2 = P_i$,

$$\sum_i R_i^2 = N \kappa_W (1 - \kappa_W) I_W. \quad (3.3.1w_{12e})$$

Write $X = X_B$, $Y_i = X_{[a_i, B]}$, and, for arbitrary $Z \in \text{End}(W)$,

$$\begin{aligned} L_U(Z) &= -\frac{1}{N} \sum_i A_i \{X, Z\} a_i, & L_V(Z) &= -\frac{1}{N} \sum_i A_i X Z a_i, \\ G_E(Z) &= \frac{1}{N} \sum_i R_i \{Y_i, Z\} a_i, & H_B(Z) &= -\frac{1}{N} X \sum_i A_i Z a_i. \end{aligned} \quad (3.3.1w_{12f})$$

The exact universal root identity [Equation \(2.6.8\)](#), followed by the circle and innovation adjoints, gives the collapsed complete-status source

$$J_B^{\alpha, \beta} = -\beta L_U + (\alpha - \beta) G_E + (\beta - \alpha) L_V + (\alpha - \beta) \kappa_W H_B. \quad (3.3.1w_{12g})$$

The last coefficient is essential. The residue is a constant star field, and its root lift is $\mathcal{T}^* \mathcal{E}^* H_B = \mathcal{K}_W H_B = \kappa_W H_B$.

For a linear map $J : S_2(W) \rightarrow W$, define its coefficient slices and row square by

$$M_r = J^* a_r, \quad \mathfrak{Q}(J) = \sum_r M_r M_r^*.$$

We claim

$$\boxed{\mathfrak{Q}(J_B^{\alpha, \beta}) \preceq \frac{C}{N} \|B\|_F^2 I_W.} \quad (3.3.1w_{12h})$$

For the circle terms, [Equation \(3.3.1w_{12g}\)](#) reduces them to

$$\Phi_1(Z) = \frac{\alpha}{N} \sum_i A_i X Z a_i, \quad \Phi_2(Z) = \frac{\beta}{N} \sum_i A_i Z X a_i.$$

Orthonormality of (a_i) gives the exact row square

$$\mathfrak{Q}(\Phi_1) = \frac{|\alpha|^2}{N^2} X \left(\sum_i A_i^* A_i \right) X \preceq \frac{4(n-2)}{N^2} \|B\|_F^2 I_W.$$

Here we used $\sum_i A_i^* A_i = (n-2)I_W$ and $X_B^2 \preceq 4\|B\|_F^2 I_W$. The normalization $\|a_i\|_{\text{op}} = 2^{-1/2}$ also gives $\|A_i\|_{\text{op}} \leq 2\|a_i\|_{\text{op}} = \sqrt{2}$. For the other orientation use $\sum_i \|Xa_i\|^2 = (n-2)\|B\|_F^2$ and Cauchy–Schwarz:

$$\langle y, \mathfrak{Q}(\Phi_2)y \rangle \leq \frac{1}{N^2} \left(\sum_i \|Xa_i\|^2 \right) \left(\sum_i \|A_i y\|^2 \right) = \frac{(n-2)^2}{N^2} \|B\|_F^2 \|y\|^2.$$

Both are $O(N^{-1}\|B\|_F^2)$.

Write $G_E = \Gamma_1 + \Gamma_2$ with

$$\Gamma_1(Z) = \frac{1}{N} \sum_i R_i Y_i Z a_i, \quad \Gamma_2(Z) = \frac{1}{N} \sum_i R_i Z (Y_i a_i).$$

The basis-free Casimir identity $\sum_i \|[a_i, B]\|_F^2 = n\|B\|_F^2$ implies

$$\sum_i \|Y_i\|_{\text{op}}^2 \leq 4n\|B\|_F^2, \quad \sum_i \|Y_i a_i\|^2 \leq 4n\|B\|_F^2.$$

Thus, for $y \in W$,

$$\langle y, \mathfrak{Q}(\Gamma_1)y \rangle = \frac{1}{N^2} \sum_i \|R_i Y_i y\|^2 \leq \frac{4n}{N^2} \|B\|_F^2 \|y\|^2.$$

For the other orientation, [Equation \(3.3.1w_{12e}\)](#) gives

$$\langle y, \mathfrak{Q}(\Gamma_2)y \rangle \leq \frac{4n\kappa_W(1-\kappa_W)}{N} \|B\|_F^2 \|y\|^2 \leq \frac{C}{N} \|B\|_F^2 \|y\|^2.$$

Finally

$$\mathfrak{Q}(H_B) = \frac{1}{N^2} \sum_i A_i^* X^2 A_i \preceq \frac{4(n-2)}{N^2} \|B\|_F^2 I_W.$$

The finite-sum row inequality and bounded status coefficients prove [Equation \(3.3.1w_{12h}\)](#).

We record the literal- W separation before inserting the Green rail. Fix the chronological child, ribbon/status data, and every prescribed nonroot gate. Move every fresh-plane action to its named root incidence, form the complete source [Equation \(3.3.1w_{12g}\)](#), and cut the input and output of its passive operator slot. Close every other transforming ancillary leg at its recorded circuit base before attaching the four-leg complement. The four maps L_U, L_V, G_E, H_B then have exactly one transforming output, the displayed copy of W ; the two cut passive indices are external ports, not an amplification of that output. Thus an empty step on the Green rail is precisely $N^{-1} \sum_i P_i = \kappa_W I_W$. An independently transforming ancillary representation is not allowed through this rail; ancillary identities remain outside it and are absorbed into the final complement.

Reflection exchanges the second input and output as already encoded in [Equation \(3.3.1w_{12a}\)](#). Hence the physical four-leg operator before the Green transfer is exactly

$$\mathcal{A}_0(B, B') = \sum_r \overline{M_r} \otimes (M'_r)^\top. \tag{3.3.1w_{12i}}$$

The operator Cauchy–Schwarz–Haagerup inequality and [Equation \(3.3.1w_{12h}\)](#) yield

$$\|\mathcal{A}_0(B, B')\|_{\text{op}} \leq \|\mathfrak{Q}(J_B)\|_{\text{op}}^{1/2} \|\mathfrak{Q}(J_{B'})\|_{\text{op}}^{1/2} \leq \frac{C}{N} \|B\|_F \|B'\|_F.$$

The only transforming output through the root Green band is the displayed adjoint module W , on which

$$R_s = r_s I_W, \quad R_z = (1 - z\kappa_W)^{-1} I_W, \quad |r_s|, |1 - z\kappa_W|^{-1} \leq (1 - \kappa_W)^{-1} = \frac{N}{2(n-2)}.$$

This proves Equation (3.3.1w_{12b}) at distinct times.

For coincident roots retain the common plane label and define

$$F_i(B, Z) = \alpha\{Y_i, Z\}a_i + \beta\{X, [A_i, Z]\}a_i.$$

The map $Z \mapsto N^{-1/2}(F_i(B, Z))_i$ is the sum of the five families

$$(Y_i, a_i), \quad (I, Y_i a_i), \quad (X A_i, a_i), \quad (A_i, X a_i), \quad (-I, A_i X a_i), \quad (3.3.1w_{12j})$$

where (L_i, v_i) denotes $Z \mapsto L_i Z v_i$. Its coefficient row satisfies

$$\sum_{i,r} M_{i,r} M_{i,r}^* = \frac{1}{N} \sum_i \|v_i\|^2 L_i^* L_i$$

for each family. The two preceding Casimir bounds, together with $\sum_i \|X a_i\|^2 = (n-2)\|B\|_F^2$ and $\sum_i A_i^* A_i = (n-2)I_W$, bound every family by $Cn^{-1}\|B\|_F^2 I_W$. Applying the finite-sum row inequality and then Haagerup to the combined index (i, r) gives the coincident version of Equation (3.3.1w_{12b}).

Trace duality proves Equation (3.3.1w_{12c}). Tensoring by an identity outside the Green rail leaves the root operator norm unchanged and lets the joint ancillary factor remain inside K ; no factor $\dim E$ is introduced. $\square$

3.3.2 Packetless hard and soft cells

Lemma 3.26 (Root-only aggregate-Ricci scalar-dimer descendants). *Let*

$$\mathcal{C}_{1,R} = \sum_{t=1}^m [A_t, X_{\mathfrak{B}_{t-1}^-}] v_t,$$

and let $\mathcal{C}_{k,R}^{\dim}$ be the part of the Christoffel–Darboux/monomer–dimer expansion in which every factor on both sides of the Ricci root is a recurrence dimer. Then

$$\boxed{\mathcal{C}_{k,R}^{\dim} = \mathbf{1}_{\{k \text{ odd}\}} \frac{k+1}{2} (-\gamma)^{(k-1)/2} \mathcal{C}_{1,R}.} \quad (3.3.1w_{12k_0})$$

Moreover

$$\boxed{\mathbb{E}\|\mathcal{C}_{1,R}\|^2 \leq \frac{m(m-1)}{Nn}, \quad \mathbb{E}\|\mathcal{C}_{k,R}^{\dim}\|^2 \leq Cnc^{k+1}(k+1)^2.} \quad (3.3.1w_{12k_1})$$

Proof. The unique all-dimer child of Q_{2a} is $(-\gamma)^a I$, and an odd Q has no all-dimer child. In Equation (3.2.7), both polynomial factors are even exactly when $k = 2r + 1$ and $j = 0, 2, \dots, 2r$. All $r + 1$ terms have the same coefficient $(-\gamma)^r$, proving Equation (3.3.1w_{12k₀}).

For the energy bound, if x, y are unit simple bivectors and $B_x = (-2x^2 - (2/n)I)/(n-2)$, direct multiplication gives

$$[y, X_{B_x} y] = [X_{B_y} x, x] = -\frac{2}{n-2} [y^2, x^2]. \quad (3.3.1w_{12k_2})$$

Because $j_t = \sqrt{2}v_t$ and $A_t = \text{ad}_{j_t}$, expansion of the prefix aggregate gives

$$\mathcal{C}_{1,R} = \sqrt{2} \sum_{s=1}^{m-1} [X_{\mathfrak{B}_{s+1}^+} v_s, v_s], \quad \mathfrak{B}_{s+1}^+ = \sum_{t>s} B_{v_t}. \quad (3.3.1w_{12k_3})$$

The last sum is adapted to reverse exposure. In the current coordinate frame, $a_{pq} = 2^{-1/2}(E_{pq} - E_{qp})$ and

$$[X_B a_{pq}, a_{pq}] = [B, a_{pq}^2] = -\frac{1}{2}[B, L_{pq}].$$

Expanding matrix entries gives

$$\sum_{p<q} \|[X_B a_{pq}, a_{pq}]\|_{\text{HS}}^2 = \frac{n-2}{2} \|B_{\text{off}}\|_F^2 \leq \frac{n-2}{2} \|B\|_F^2, \quad (3.3.1w_{12k_4})$$

while the sum of these commutators is zero because $\sum_{p<q} a_{pq}^2 = -(n-1)I/2$. Thus the summands in Equation (3.3.1w_{12k₃}) are reverse Hilbert-space martingale differences. The aggregate Ricci increments have squared Frobenius mean $2/[n(n-2)]$, whence

$$\mathbb{E}\|\mathcal{C}_{1,R}\|^2 \leq \frac{n-2}{N} \sum_{s=1}^{m-1} \frac{2(m-s)}{n(n-2)} = \frac{m(m-1)}{Nn}.$$

Finally use $m \leq cN$, $N/n \leq n/2$, and $\gamma \leq c$ in Equation (3.3.1w_{12k₀}). $\square$

Lemma 3.27 (Packetless Ricci component with an internal hard resource). *Fix a complete squared-shell child through its monomer–dimer, coefficient-pairing, ribbon, status, incidence, and Haar-residual data, except that the complete reflection/status orbit and the complete weak-order, equality, and finite-gap sum are retained inside the residual component $\mathcal{C}_R$ containing the selected aggregate-Ricci root R . Suppose $\mathcal{P}_R = \emptyset$ and, after equal-time fusion, $\mathcal{C}_R$ contains at least one independently token-bearing hard resource: either a nonsingleton packet child for which the one-token presentation has already proved a numerical factor $\delta_P \leq N^{-1}$, or a time-indexed linear Jacobi child $-2N^{-1}W_i$. An unsplit mixed/full-centered hard block with only unit or $n^{-1/2}$ relative cost is not included. A resolved fixed-column child whose complete closed coefficient scalar has a proved factor $\delta_P \leq N^{-1}$ relative to the bare Euler carrier is included. Nor is a resource in a different Haar-separated component.*

Let $f_R \in \{1, 2\}$ be the number of fundamental $p \vee r$ circuits hit by the four Ricci legs, let

$$\mathfrak{e}_R = \frac{C}{n} \|B\|_F \|B'\|_F,$$

and let $\mathfrak{a}_{\Gamma, \mathcal{C}_R}$ be the remaining positive, once-normalized circuit-anchored terminal majorant. Then

$$\sum_{\substack{\text{weak orders, equality blocks, finite gaps,} \\ \text{and typed states inside } \mathcal{C}_R}} |\text{Contr}_{\Gamma, \mathcal{C}_R}| \leq A_L^{r_R} (r_R + 1)^B \mathfrak{a}_{\Gamma, \mathcal{C}_R} \mathfrak{e}_R N^{-1}. \quad (3.3.1w_{12m})$$

Consequently the complete packetless subfiber in which the Ricci component has such a resource satisfies the shell bound on the right of Equation (3.3.1w_{12k}).

Proof. Order these independently token-bearing hard resources in $\mathcal{C}_R$ by physical time, breaking an equality tie by the fixed ribbon-slot order, and reserve the first one, denoted δ_* . If it is a packet P , use its proved one-token presentation

$$\text{cost}(P) \leq C^{|P|} n^{c^*(P)} \delta_P, \quad 0 \leq \delta_P \leq N^{-1}. \quad (3.3.1w_{12m_1})$$

The bare factor $n^{\epsilon^*(P)}$ is already one factor of $\mathfrak{a}_{\Gamma, C_R}$, whereas δ_P is its distinct normalized plane-star/variance token. We do not also use the sharper mobile-front estimate. If the resource is Jacobi, put $\delta_* = N^{-1}$ from the literal coefficient of $-2N^{-1}W_i$ and leave W_i as a prescribed gate in the terminal majorant. Thus in both cases δ_* is disjoint from every circuit base, packet front, and the Ricci endpoint $\mathfrak{e}_R$. Fusion of δ_* into the root equality block creates no additional anchor and does not alter this ownership.

Resolve every other hard full-centered packet at its vertices, at the already allowed exponential cost 2^{r_R} . Let u be the number of resulting mixed hard packet children; each contains an actual typed Ricci occurrence and retains its proved $n^{-1/2}$ relative debit. Let $b \geq 1$ be the number of token-bearing hard resources in the component, and let o and v be respectively the numbers of unfused $S^\pm$ and V singleton supernodes. Repeat the one-variable finite-simplex proof of [Lemma 3.39](#), keeping the complete joint Ricci block as its already proved opaque typed endpoint. For fixed ordered supernodes and a fixed star-state path, the unit-circle generating function obeys

$$\sup_{|z|=1} |\mathcal{F}_\omega(z)| \leq A^{r_R} (r_R + 1)^B \mathfrak{a}_{\Gamma, C_R} \mathfrak{e}_R N^{-b-o/2} n^{-u/2-v/4}. \quad (3.3.1w_{12m_2})$$

Here the b numerical factors are distinct proved packet tokens or literal Jacobi coefficients, and the u factors are distinct displayed mixed occurrences; all are relative weights in the same once-normalized presentation. The Ricci block supplies the component endpoint and its single translation; no fixed-atom Ricci trace-ideal estimate is used in [Equation \(3.3.1w_{12m_2}\)](#).

Factor the reserved $\delta_* = N^{-1}$ and put

$$t' = (b - 1) + u + o + v.$$

If $t' > 0$, the scalar Dirichlet bound for the exact finite simplex costs one $C \log(n + 2)$. Before that logarithm the remaining dimension exponent is

$$E = 2(b - 1) + \frac{u}{2} + o + \frac{v}{4} \geq \frac{t'}{4}.$$

Exactly as in [Equations \(3.4.1x_5\) to \(3.4.1x_6\)](#), half of the retained power pays the logarithm. If h is the number of ordered supernodes, then $h \leq t' + B_0 + 2$. For h above a constant depending only on L , the other half pays h^{r_R} because $r_R \leq L \log n$; for bounded h the assignment mass is absorbed in $A_L^{r_R}$. The finite choices of (b, o, v) , root placement, status, and incidence data have the same exponential cost. Hence δ_* remains intact after every internal weak-order and equality assignment has been summed.

If $t' = 0$, after aligned current strings and every internal special gate have been absorbed into their prescribed opaque blocks, the component has exactly two external supernodes: the complete reflected Ricci block (including its internal two-replica Green form) and the reserved hard endpoint. There is one external relative gap. Its finite Green form is the prescribed two-endpoint Riesz block and is bounded directly, as in the $t_C = 0$ clause of [Lemma 3.39](#); no Dirichlet-kernel bound is taken. This proves [Equation \(3.3.1w_{12m}\)](#) in all cases. Choosing the reserved resource costs at most r_R and is absorbed in the displayed polynomial.

For the spatial comparison, the once-normalized simultaneous majorant obeys $\mathfrak{a}_{\Gamma, C_R} \leq A^r n^{2-g}$ times its already recorded outside weights. Since

$$N^{-1} = \frac{2}{n(n-1)} \leq 4n^{-2}, \quad f_R \leq 2,$$

we obtain

$$n^{2-g} \frac{C}{n} N^{-1} \leq C n^{-1-g} \leq C n^{1-g-f_R}. \quad (3.3.1w_{12m_3})$$

This is a scalar product of disjoint resources after the circuit presentation has been normalized once; it does not match the hard packet to a particular root-hit circuit and uses no second base. The remaining Haar-separated components not containing R retain their own endpoints and translations and are summed by [Theorem 3.44](#); restriction of a global time assignment to those components is injective. The Ricci component still has only its one root translation. Finally the reverse martingale moment bound [Equation \(3.5.5\)](#) and the unchanged monomer–dimer ledger give the claimed complete-shell consequence. $\square$

Lemma 3.28 (Resolved mixed packets are internally hard). *Fix a complete packetless aggregate–Ricci atom through its orthogonal–Weingarten pair (p, q) , trace pairing, statuses, reflections, chronological data, finite gaps, and its one declared base on every original $p \vee r$ circuit. Resolve one ordinary full-centered packet P at one physical Kac plane x into letters $D_v \in \{W_x^\circ, X_{B_x}\}$. If the resolved child is not all Weyl, then either it is a one-box zero or its complete coefficient-column scalar satisfies*

$$|Z_{q_P}(D_1, \dots, D_s)| \leq A^s n^{c^*(P)-3}. \quad (3.3.1w_{12m_4})$$

In the second case it is an independently token-bearing hard resource in the sense of [Lemma 3.27](#), with the bare-Euler presentation

$$|Z_{q_P}| \leq A^s n^{c^*(P)} \delta_P, \quad \delta_P = n^{-3} \leq N^{-1}. \quad (3.3.1w_{12m_5})$$

Thus [Lemma 3.27](#) covers every nonzero mixed-packet child.

Proof. For a fixed Weingarten term, the pairing p contracts row/trace indices and q contracts the four coefficient indices at every curvature box. The aggregate–Ricci boundary, finite Green and Riesz maps, statuses, compact row bends, and circuit closures carry no q -dart. Ordinary packet roots are themselves coefficient boxes. Hence the component of q containing P is closed before any row-side map is estimated, and the complete scalar factors exactly as

$$\text{Wg}_n^O(p, q) \mathcal{R}_{p,r,\xi} Z_{q_P}(D_1, \dots, D_s) \prod_{Q \neq P} Z_{q_Q}. \quad (3.3.1w_{12m_6})$$

All row, root, Green, status, and circuit data are in $\mathcal{R}_{p,r,\xi}$. Simultaneous orthogonal conjugation of the D_v cancels along the q_P -deltas, so Z_{q_P} is independent of the orientation of x and may be extracted before the plane and chronological sums. This also proves that the scalar and every circuit base are disjoint resources.

We next prove the numerical estimate rather than cite an opened packet norm. If self-adjoint algebraic-curvature letters $D_1, \dots, D_s$ are simultaneously diagonal in a decomposable-wedge basis, expand the direct/cross alternative at every box. One smoothing h has $k(h) \leq c^*(P)$ scalar circuit variables and contributes at most

$$\sum_{y_1, \dots, y_{k(h)}} \prod_{v=1}^s |\lambda_{y_{\alpha(v)}, y_{\beta(v)}}^{(v)}|. \quad (3.3.1w_{12m_7})$$

A self-loop is zero by skewness. Otherwise choose an anchor box a , sum the two distinct variables incident to it, and bound all remaining letters pointwise. Summing the at most 2^s smoothings gives

$$|Z_{q_P}(D_1, \dots, D_s)| \leq A^s n^{c^*(P)-2} \|D_a\|_1 \prod_{v \neq a} \|D_v\|_{\text{op}}. \quad (3.3.1w_{12m_8})$$

For a coordinate Kac plane the three overlap sectors give

$$\|W_x^\circ\|_{\text{op}} \leq 1, \quad \|W_x^\circ\|_1 = \frac{4(n-3)}{n-1} \leq 4, \quad \|X_{B_x}\|_{\text{op}} = \frac{2}{n}, \quad \|X_{B_x}\|_1 = \frac{4(n-3)}{n} \leq 4. \quad (3.3.1w_{12m_9})$$

If a Weyl letter is present, choose it as the trace-class anchor in Equation (3.3.1w_{12m₈}); every Ricci nonanchor contributes $2/n$. Since the child is mixed, there is at least one such nonanchor and Equation (3.3.1w_{12m₄}) follows. If every letter is Ricci, the one-box scalar is $\text{Tr } X_{B_x} = 0$. Otherwise use one Ricci letter as the trace-class anchor and a second one in operator norm, giving the same $n^{e^*(P)-3}$ bound. Rewriting it relative to the bare factor $n^{e^*(P)}$ gives Equation (3.3.1w_{12m₅}), and $n^{-3} \leq 2/[n(n-1)] = N^{-1}$. The exact factorization Equation (3.3.1w_{12m₆}) verifies every ownership hypothesis of Lemma 3.27. $\square$

Lemma 3.29 (Soft Ricci component above the passive-debit threshold). *Work in the packetless subfiber and suppose the Ricci residual component has no independently token-bearing hard resource covered by Lemma 3.27. Resolve every internal hard full-centered packet into all-Weyl and mixed children. The all-Weyl token-bearing children and, by Lemma 3.28, every nonzero mixed child fall under that lemma. Thus the retained soft family contains no packet child. Let o be the number of unfused $S^\pm$ singleton supernodes and v the total number of V singleton supernodes in all maximal consecutive runs of the Ricci component. Put*

$$\Delta_0 = o + \frac{v}{2}. \quad (3.3.1w_{12n})$$

If $\Delta_0 \geq f_R$, where $f_R \in \{1, 2\}$ is the number of root-hit fundamental circuits, then this complete subfiber satisfies the shell bound on the right of Equation (3.3.1w_{12k}).

Proof. Keep the complete weak-order/equality sum inside the Ricci component and encode all of its residual gaps by the single unit-circle parameter of Equation (3.4.1x₂). Each $S^\pm$ seed supplies $N^{-1/2}$; every packet child has already been routed to the internal-hard family. Apply Lemma 3.12 to each maximal V run, with the actual typed incidences at its two ends. Since all runs retain the same z , this is an estimate of the intact generating function, not a sequential replacement of the coupled simplex by rectangles. Prescribed gates between runs remain inside the terminal scalar. Thus, for fixed ordered supernodes and local states,

$$\sup_{|z|=1} |\mathcal{F}_\omega(z)| \leq A^{r_R} (r_R + 1)^B \mathfrak{a}_{\Gamma, C_R} \mathfrak{e}_R N^{-o/2} n^{-v/2} \leq A^{r_R} (r_R + 1)^B \mathfrak{a}_{\Gamma, C_R} \mathfrak{e}_R n^{-\Delta_0}. \quad (3.3.1w_{12n_1})$$

The harmless factor $2^{o/2}$ from $N^{-o/2} \leq 2^{o/2} n^{-o}$ is absorbed in A^{r_R} .

We next remove the Taylor-partial-sum logarithm without spending any of this debit. Let d count every z -denominator, including the literal- W root Green rail; direct expansion of the same prescribed-gate scalar gives

$$|[z^q] \mathcal{F}_\omega(z)| \leq A^{r_R} (r_R + 1)^B \mathfrak{a}_{\Gamma, C_R} \mathfrak{e}_R \binom{q + d - 1}{d - 1} (1 - \gamma_n)^q. \quad (3.3.1w_{12n_2})$$

Every residual transfer is compressed to the noninvariant subspace, every z -independent prescribed gate has already been included in the displayed coefficient mass, and $d \leq r_R + B_0$. The literal- W root Green eigenvalue has a larger spectral gap than γ_n , so the same coefficient envelope applies to that rail. Take the common H in Lemma 3.13 to be the displayed spatial majorant times the maximal prefix-Ricci envelope used in Equation (3.5.5). The prefix envelope is uniform in the physical translation and in every coefficient of the common gap series. Parametrize the translation

orbit by its first fused supernode; [Lemma 3.14](#) therefore verifies the distinct-budget hypothesis of [Lemma 3.13](#). The lemma sums the one component translation and retains the full $n^{-\Delta_0}$ factor. Its factor cN is precisely the already budgeted worst-case root-translation count; it is not a second anchor.

Put $t_{\text{soft}} = o + v$. Every counted node contributes at least $1/2$ to Δ_0 , and therefore

$$t_{\text{soft}} \leq 2\Delta_0. \quad (3.3.1w_{12n_3})$$

If $\Delta_0 = f_R$, then $t_{\text{soft}} \leq 2f_R \leq 4$, so the total number $h \leq t_{\text{soft}} + B_0 + 1$ of ordered supernodes is bounded absolutely and the assignment mass h^{r_R} is absorbed in $A_L^{r_R}$. If $\Delta_0 > f_R$, put $\varepsilon = \Delta_0 - f_R \geq \frac{1}{2}$. Exactly as in [Equations \(3.4.1x₅\) to \(3.4.1x₆\)](#), a fixed fraction of $n^{-\varepsilon}$ absorbs h^{r_R} when t_{soft} is large; for bounded t_{soft} it is again absorbed in $A_L^{r_R}$. The local resolution, status, incidence, and root-placement choices have exponential mass. In both cases the complete component sum retains, apart from its single already budgeted root translation,

$$A_L^{r_R}(r_R + 1)^B \mathfrak{a}_{\Gamma, C_R} \mathfrak{e}_R n^{-f_R}. \quad (3.3.1w_{12n_4})$$

Finally $\mathfrak{a}_{\Gamma, C_R} \leq A^r n^{2-g}$ and $\mathfrak{e}_R = (C/n)\|B\|_F\|B'\|_F$, so [Equation \(3.3.1w_{12n₄}\)](#) has exactly the target spatial power $A^r n^{1-g-f_R}\|B\|_F\|B'\|_F$. Exterior Haar-separated components keep their own endpoints, and the same component-allocation and reverse-martingale moment argument used after [Equation \(3.3.1w_{12m₃}\)](#) completes the shell estimate. More explicitly, the cN in [Lemma 3.13](#) occupies the single factor m in the existing bound $m(C/n)Cc \leq Cc^2n$; [Equation \(3.5.5\)](#) bounds the common maximal prefix envelope and does not introduce a second translation sum. $\square$

3.3.3 Invariant-column closure of the low cells

Lemma 3.30 (Ancilla-stable nuclear-to-Ricci entrance). *Let $\mathcal{A}, \mathcal{B}$ be Hilbert spaces and let*

$$Z : W \otimes \mathcal{B} \longrightarrow W \otimes \mathcal{A}$$

be trace class. Realign its Hilbert vectorization from the physical grouping $(W \otimes \mathcal{A}) \mid (W \otimes \mathcal{B})$ to the grouping $(W \otimes W) \mid (\mathcal{A} \otimes \overline{\mathcal{B}})$, without transposing a proper subsystem, and denote the resulting Hilbert vector by $\Re(\text{vec } Z)$. If

$$P_{\mathcal{R}} = \frac{1}{n-2} XX^* \quad (3.3.1w_{12p})$$

is the orthogonal projection of $S_2(W)$ onto the traceless-Ricci summand, then

$$\|(P_{\mathcal{R}} \otimes I)\Re(\text{vec } Z)\|_2 \leq \frac{2}{\sqrt{n-2}}\|Z\|_1. \quad (3.3.1w_{12p_1})$$

The estimate remains true after the diagonal Haar projection. At a predeclared joint cut in the packetless aggregate-Ricci construction of [Lemma 3.33](#), Z may be the complete once-normalized port state; [Equation \(3.3.1w_{12p₁}\)](#) then uses no new circuit base.

Proof. We first record the rank-one Ricci estimate. Identify $u, v \in W$ with skew matrices in the normalized wedge convention. Direct contraction of one skew pair gives

$$X^*(|u\rangle\langle v|) = -(uv + vu)_0 \quad (3.3.1w_{12p_2})$$

up to a harmless sign fixed by the orientation of the vectorization. Therefore

$$\|X^*(|u\rangle\langle v|)\|_F \leq 2\|u\|_F\|v\|_F. \quad (3.3.1w_{12p_3})$$

Since $P_{\mathcal{R}} = (n-2)^{-1}XX^*$ and $X^*X = (n-2)I$ on $\text{Sym}_0(H)$,

$$\|P_{\mathcal{R}}(|u\rangle\langle v|)\|_2^2 \leq \frac{4}{n-2}\|u\|^2\|v\|^2. \quad (3.3.1w_{12p4})$$

By spectral decompositions and linearity, Equation (3.3.1w_{12p4}) implies for all positive ρ, σ on W

$$\text{Tr}(P_{\mathcal{R}}(\rho \otimes \sigma^{\text{T}})) \leq \frac{4}{n-2} \text{Tr} \rho \text{Tr} \sigma. \quad (3.3.1w_{12p5})$$

For $Z = |u\rangle\langle v|$, with $u \in W \otimes \mathcal{A}$ and $v \in W \otimes \mathcal{B}$, let ρ_u and ρ_v be the reduced positive densities on W . An index contraction in the stated physical realignment gives exactly

$$\text{Tr}_{\mathcal{A} \otimes \overline{\mathcal{B}}} |\Re(\text{vec } |u\rangle\langle v|)\rangle \langle \Re(\text{vec } |u\rangle\langle v|)| = \rho_u \otimes \rho_v^{\text{T}}. \quad (3.3.1w_{12p6})$$

Equations (3.3.1w_{12p5}) to (3.3.1w_{12p6}) bound the square of the left side of Equation (3.3.1w_{12p1}) by $4\|u\|^2\|v\|^2/(n-2)$. For general Z , take a singular-value decomposition $Z = \sum_j s_j |u_j\rangle\langle v_j|$ and apply the Hilbert triangle inequality. The result is

$$\frac{2}{\sqrt{n-2}} \sum_j s_j = \frac{2}{\sqrt{n-2}} \|Z\|_1.$$

The Ricci projector is equivariant and hence commutes with the diagonal Haar projection, which is contractive. Finally Lemma 2.7 supplies $\|Z\|_1$ after exactly one base has been fixed on each original circuit. Realignment is used only in Hilbert norm after that estimate, so it is neither a new cut nor a second trace-class factorization. $\square$

Lemma 3.31 (Normalized invariant-column insertion). *Let V be a real orthogonal space of dimension d , with*

$$\Omega_V = d^{-1/2} \sum_{\mu=1}^d e_{\mu} \otimes e_{\mu}.$$

Let $R_i : L \rightarrow M_i$ and orthogonal projections $P_{i,m}$ satisfy

$$\sum_i R_i^* R_i \preceq I_L, \quad [P_{i,m}, R_i^* R_i] = 0. \quad (3.3.1w_{12p7})$$

Let $T : V \rightarrow L \otimes E$ be fixed, let $S_{i,\theta}$ be contractions on the passive factors, and let $\rho_i(\theta)$ be orthogonal on V . Then

$$\boxed{\sum_i \int \|(R_i P_{i,m} \otimes S_{i,\theta})(T \rho_i(\theta) \otimes I) \Omega_V\|^2 \frac{d\theta}{2\pi} \leq \frac{1}{d} \|T\|_2^2 \leq \|T\|_{\text{op}}^2.} \quad (3.3.1w_{12p8})$$

The same assertion holds if an own-circle output rotation is placed between T and $R_i P_{i,m}$, provided the latter intertwines that rotation. It is stable under passive Hilbert amplification, adjunction, full transpose, and orthogonal rail permutation.

Proof. The reduced density of Ω_V on either factor is $d^{-1}I_V$. After discarding $S_{i,\theta}$ and using invariance of Hilbert–Schmidt norm under right multiplication by $\rho_i(\theta)$, the left side is at most

$$\frac{1}{d} \sum_i \|R_i P_{i,m} T\|_2^2 = \frac{1}{d} \text{Tr} \left(T^* \left[\sum_i P_{i,m} R_i^* R_i P_{i,m} \right] T \right) \leq \frac{1}{d} \|T\|_2^2.$$

Since the domain of T has dimension d , $d^{-1}\|T\|_2^2 \leq \|T\|_{\text{op}}^2$. An output rotation moves through the intertwining row and disappears from the norm. The remaining symmetries are Hilbert isometries. This proves the lemma. $\square$

For the fixed maximizing smoothing h of a ribbon record Γ , put

$$t_\partial(\Gamma) := \#(h \vee r).$$

This is the number of rooted boundary components. The surface construction in the proof of [Lemma 2.20](#) gives $1 \leq t_\partial \leq 2$ and $g_{\text{top}} - t_\partial = g_h \geq 0$; hence $g = g_{\text{top}} + g_{Wg} \geq t_\partial$. This is the boundary coordinate used in every low-cell tuple below.

Lemma 3.32 (Native selected-time completion after the joint defect bound). *Put $j_i = \sqrt{2}a_i$, $\hat{A}_i = \text{ad}_{j_i}$, and let $\mathcal{A}_i$ denote the zero Fourier projection for conjugation of $\text{Sym}_0(H)$ by the coordinate circle of plane i . Let $P_{i,m}$ be the Fourier projections for the same circle acting by conjugation on $S_2(W)$. For bounded status scalars α, β , set*

$$\mathcal{F}_{B,i}(Y) = \frac{1}{\sqrt{N}} \left(\alpha \{[\hat{A}_i, X_B], Y\} a_i + \beta \{X_B, [\hat{A}_i, Y]\} a_i \right) \quad (3.3.1w_{12q})$$

and, for

$$\mathbf{M}_i^C = |a_i\rangle\langle a_i| - N^{-1}I_W, \quad \mathbf{M}_i^W = \mathbf{M}_i^C - X_{B_i},$$

define the correctly ordered generalized defect

$$\mathcal{D}_i^\bullet(B, Z) = (n-2) [\mathcal{F}_{B,i}(P_{\mathcal{R}} \mathbf{M}_i^\bullet Z) - \mathbf{M}_i^\bullet \mathcal{F}_{B,i}(P_{\mathcal{R}} Z)]. \quad (3.3.1w_{12q_1})$$

Then:

- (i) for every own-circle-invariant C_i , the zero-coefficient/zero-active-mode block vanishes,

$$\mathcal{D}_i^\bullet(C_i, P_{i,0}Z) = 0; \quad (3.3.1w_{12q_2})$$

- (ii) for every $m \neq 0$, every own-circle-invariant family $C_i \in \text{Sym}_0(H)$, and arbitrary ancillary contractions $S_{i,-m}$,

$$\left\| \frac{1}{N} \sum_i (\mathcal{D}_i^\bullet(C_i) P_{i,m}) \otimes S_{i,-m} \right\|_{\text{op}} \leq \frac{C}{N} \left(\frac{1}{N} \sum_i \|C_i\|_F^2 \right)^{1/2}; \quad (3.3.1w_{12q_3})$$

In particular $C_i = \mathcal{A}_i B$ gives the right side $CN^{-1}\|B\|_F$. The assertion is stable under passive Hilbert amplification, status, reflection, adjunction, and full transpose; and

- (iii) in the residual packetless cell $(g, t_\partial, f_R; u, o, v) = (1, 1, 2; 0, 0, 1)$, use its unique $\mathbf{V}$ supernode. In its coefficient-zero sector the complete selected-time tensor is one prescribed bounded gate. Writing $\mathcal{H}_{\text{act}}$ for the root-facing active curvature-letter carrier and $E_{\text{in}}, E_{\text{out}}$ for all other rails declared by the native chronological matricization, that gate is the typed map

$$\begin{aligned} G_\Xi(B) &: \mathcal{H}_{\text{act}} \otimes E_{\text{in}} \longrightarrow W \otimes E_{\text{out}}, \\ G_\Xi(B) &= \frac{1}{N} \sum_i \sum_{m \neq 0} (\mathcal{D}_i^\bullet(\mathcal{A}_i B) P_{i,m}) \otimes S_{i,-m}, \\ \|G_\Xi(B)\|_{\text{op}} &\leq \frac{C}{N} \|B\|_F. \end{aligned} \quad (3.3.1w_{12q_4})$$

This is the exact selected-time gate for the low-debit cell

$$(g, t_\partial, f_R; u, o, v) = (1, 1, 2; 0, 0, 1). \quad (3.3.1w_{12q_5})$$

The chronological sum and its endpoint energy are not part of this local statement; they are proved separately in [Lemmas 3.36](#) and [3.37](#).

Proof. Fix the family C_i in (ii). Circle invariance gives $[\hat{A}_i, X_{C_i}] = 0$. Both $P_{\mathcal{R}}$ and the two centered multipliers commute with that circle. If Z has active frequency zero, then every commutator occurring in Equation (3.3.1w_{12q}) vanishes for each of the two inputs in Equation (3.3.1w_{12q1}). This proves Equation (3.3.1w_{12q2}).

The scalar terms in Equation (3.3.1w_{12q1}) cancel for the full-centered multiplier. Put $P_i^W = |a_i\rangle\langle a_i|$ and $r_i Z = Z^* a_i$. If $m \neq 0$, the second root output in

$$(n-2) \left[\mathcal{F}_{C_i, i}(P_{\mathcal{R}} P_i^W Z) - P_i^W \mathcal{F}_{C_i, i}(P_{\mathcal{R}} Z) \right] \quad (3.3.1w_{12q6})$$

has frequency m ; it is either zero or orthogonal to the frequency-zero vector a_i . The second term is therefore zero. Direct contraction of the rank-one curvature operator gives

$$\text{Ric}_0(P_i^W P_{i,m} Z) = -\frac{1}{2}((r_i P_{i,m} Z) a_i + a_i (r_i P_{i,m} Z))_0. \quad (3.3.1w_{12q7})$$

Using $P_{\mathcal{R}} = 2(n-2)^{-1} X \text{Ric}_0$ in Equation (3.3.1w_{12q6}) yields a factorization

$$\mathcal{D}_i^C(C_i) P_{i,m} = L_i^C r_i P_{i,m}, \quad \|L_i^C\| \leq \frac{C}{\sqrt{N}} \|C_i\|_F. \quad (3.3.1w_{12q8})$$

Here the last estimate follows directly from Equation (3.3.1w_{12q}), $\|X_E\|_2 = \sqrt{n-2} \|E\|_F$, and the bounded status list.

The exact column identities are

$$\sum_i r_i^* r_i = I_{S_2(W)}, \quad [P_{i,m}, r_i^* r_i] = 0. \quad (3.3.1w_{12q9})$$

Hence the column $((r_i P_{i,m}) \otimes S_{i,-m})_i$ has norm at most one. The row $(y_i)_i \mapsto \sum_i (L_i^C \otimes I) y_i$ has squared norm at most

$$\sum_i \|L_i^C\|^2 \leq \frac{C}{N} \sum_i \|C_i\|_F^2.$$

Its composition with the column and the normalized factor N^{-1} proves Equation (3.3.1w_{12q3}) for $\bullet = C$.

For the pure-Weyl multiplier expand $M_i^W = M_i^C - X_{B_i}$. Besides the term just treated, the two corrections factor through

$$s_i Z = \text{Ric}_0(X_{B_i} Z), \quad t_i Z = (P_{\mathcal{R}} Z) a_i. \quad (3.3.1w_{12q10})$$

The coordinate overlap eigenvalues give

$$\sum_i X_{B_i}^2 = \frac{2}{n} I_W. \quad (3.3.1w_{12q11})$$

Since $\|\text{Ric}_0 Y\|_F^2 \leq (n-2) \|Y\|_2^2 / 4$,

$$\sum_i s_i^* s_i \preceq \frac{1}{2} I, \quad \sum_i t_i^* t_i = P_{\mathcal{R}}. \quad (3.3.1w_{12q12})$$

Both sums commute with their corresponding frequency projections. The remaining row maps again have norm at most $C \|C_i\|_F / \sqrt{N}$. On the only possible nonzero output frequency, $|m| = 1$, the final multiplier is the scalar $(n-2) X_{B_i} = (n-4) I / n$, of magnitude at most one. The same row-column composition proves Equation (3.3.1w_{12q3}) for $\bullet = W$.

We finally identify the physical gate. Fix the complete doubled ribbon, status, reflection, incidence, and strict time assignment, choose the unique surviving V supernode, and condition on every variable except its selected plane and angle. [Lemma 3.6](#) says that every earlier leg carries the same diagonal circle action and every later tensor is fixed. Equal-time fusion leaves exactly one unresolved centered occurrence at the selected time; a second same-time occurrence is zero, scalar, or an internal-hard packet already removed above. The two consumed positions and the literal output position are exactly the two terms of [Equation \(3.3.1w_{12q₁}\)](#), and the reflected placements are their adjoints or full transposes.

Declare, before any component estimate, ports on every rail transformed at this selected time. Keep their physical input/output order and let $S_{i,k}$ be the total Fourier projection on the tensor product of all the other selected-time rails, including normalized status and endpoint maps. It is an orthogonal spectral projection followed by one of a finite list of normalized maps, so $\|S_{i,k}\| \leq C$. Circle conservation is $b + m + k = 0$, where b is the coefficient mode. In the coefficient-zero sector, [Equation \(3.3.1w_{12q₂}\)](#) removes $m = 0$ and the exact selected-time tensor is [Equation \(3.3.1w_{12q₄}\)](#). There are only finitely many nonzero active modes, so [Equation \(3.3.1w_{12q₃}\)](#) proves its displayed norm.

There is one coefficient-zero family which is not obtained by applying $\mathcal{A}_i$ to a fixed matrix: the current Ricci increment B_{a_i} in the prefix/suffix decomposition of the aggregate coefficient. It is own-circle invariant and

$$\frac{1}{N} \sum_i \|B_{a_i}\|_F^2 = \frac{2}{n(n-2)}. \quad (3.3.1w_{12q_{12a}})$$

Apply the family form of [Equation \(3.3.1w_{12q₃}\)](#) directly; its joint gate is bounded by $C/(Nn)$, which is stronger than [Equation \(3.3.1w_{12q₄}\)](#). Thus no conditional-centering assertion about the exterior endpoint is used. An increment at a strict earlier time is fixed when the selected plane is exposed and belongs to the fixed- B specialization of the same estimate.

If the coefficient mode is nonzero, the pointwise estimate following from [Equations \(3.3.1w_{12q}\)](#) to [\(3.3.1w_{12q₁}\)](#) is

$$\|\mathcal{D}_i^\bullet((I - \mathcal{A}_i)B, Z)\|_2 \leq C\|(I - \mathcal{A}_i)B\|_F\|Z\|_2. \quad (3.3.1w_{12q_{12b}})$$

In a Haar-residual child the common auxiliary Haar conjugation and Schur orthogonality give

$$\mathbb{E}_U \frac{1}{N} \sum_i \|(I - \mathcal{A}_i)(UBU^*)\|_F^2 = \frac{4}{n+2} \|B\|_F^2. \quad (3.3.1w_{12q_{12c}})$$

Thus this sector has the extra n^{-1} square which pays the sole factor n left by the generic protected-carrier estimate of [Lemma 3.30](#). If no Haar separator is selected, the component belongs to the bare no-exterior status orbit already covered by [Lemma 3.26](#) and the soft-threshold recombination. Hence only the coefficient-zero sector requires the gate argument below.

We now prove the native-incidence assertion without inserting an arbitrary high-valent gate into a multiply cut circuit. Retain the complete doubled root word of [Equation \(3.3.1w_{12a}\)](#), close every nonroot leg in its original cyclic order, and use [Lemma 2.7](#) once. The positive presentation in [Definition 2.23](#) gives a joint trace-class complement

$$K_0 \in S_1(\mathcal{R}^{\text{in}}, \mathcal{R}^{\text{out}}), \quad (3.3.1w_{12q_{13}})$$

after the one predeclared base on each original circuit has been used. Put $M_r = A^{Cr}(r+1)^B$. The trace-class path/cycle estimate gives

$$\|K_0\|_1 \leq M_r \mathfrak{a}_{\Gamma, \tau}.$$

Here and below $\mathfrak{a}_{\Gamma,\tau}$ excludes the fixed monomer–dimer scalar $(-\gamma)^a$ according to [Definition 2.23](#); that scalar is not used in either Schatten test. For the operator norm, insert one rank-one trace-class test at the *joint* root hole and repeat the scalar path/cycle estimate from [Lemma 2.8](#) directly on the complete projective presentation. Since $f_R = 2$, the two retained original circuits remove two identity traces, and therefore

$$\|K_0\|_{\text{op}} \leq M_r n^{-2} \mathfrak{a}_{\Gamma,\tau}.$$

This is an estimate of the same normalized tangle with two different Schatten tests; it neither reuses a circuit base nor factorizes the two replicas. Schatten interpolation now gives

$$\|K_0\|_2^2 \leq \|K_0\|_1 \|K_0\|_{\text{op}} \leq M_r^2 n^{-2} \mathfrak{a}_{\Gamma,\tau}^2 \leq M_r^3 n^{-1} \mathfrak{a}_{\Gamma,\tau} \leq M_r^3 \mathfrak{a}_{\Gamma,\tau}. \quad (3.3.1\text{w}_{12q_{13a}})$$

Here the penultimate inequality is precisely the $g = 1$ Euler bound $\mathfrak{a}_{\Gamma,\tau} \leq M_r n$; it is used once. No bare cardinality or coefficient-mass bound for a chosen projective presentation is assumed. Vectorization and the fixed root/multiplicity leg permutation $\mathfrak{R}$ are Hilbert isometries. Consequently the same Hilbert bound holds after $\Pi_G \mathfrak{R} \text{vec}$.

Put

$$\mathcal{R} = \text{Sym}_0(H), \quad d_{\mathcal{R}} = \frac{(n-1)(n+2)}{2}.$$

On the invariant Ricci channel, Schur’s lemma writes the left side of [Equation \(3.3.1w_{12q_{13a}}\)](#), after $P_{\mathcal{R}}$ is applied to the active root slot, uniquely as

$$\Omega_{\mathcal{R}} \otimes \zeta, \quad \Omega_{\mathcal{R}} = \frac{1}{\sqrt{d_{\mathcal{R}}}} \sum_{\mu=1}^{d_{\mathcal{R}}} e_{\mu} \otimes e_{\mu}, \quad \|\zeta\| \leq \|K_0\|_2. \quad (3.3.1\text{w}_{12q_{13b}})$$

This is an orthogonal decomposition of the already normalized vector, not a new cup or circuit base.

Choose the root-facing Haar boundary of the residual component; at an exterior end use the common auxiliary Haar projection of [Lemma 3.17](#). The exhaustive supernode grammar contains ordinary packet/Jacobi hard nodes, $\mathbf{S}^{\pm}$ nodes, $\mathbf{V}$ nodes, and the bounded special-root list. In the present cell the first family and every packet child have already been removed, $u = o = 0$, $v = 1$, and the complete reflected aggregate–Ricci block is its special root. Thus there is no second nonconstant coefficient gate between this Haar boundary and the selected defect. A same-time nonsingleton is hard; a nonzero mixed child is internally hard; a second singleton contradicts $v = 1$; an external one-box root-color packet is zero; an external multibox root-color packet contradicts packetlessness; and an immediately adjacent root-color string is already inside the complete defect.

The remaining structural, status, complete Green/Riesz, recurrence, and aligned-root maps are equivariant on the complete carried module. Joint equivariant maps preserve [Equation \(3.3.1w_{12q_{13b}}\)](#) and act only on its multiplicity coordinate. Any remaining plane-independent nonintertwining map acts on one displayed Schur half. After conditioning on every variable except the selected (i, θ) , moving-frame exposure turns that map into

$$T_i(\theta) = \sigma_i(\theta) T \rho_i(\theta)^{-1} \quad (3.3.1\text{w}_{12q_{13c}})$$

for one fixed T . [Lemma 3.31](#), applied to the three rows in [Equations \(3.3.1w_{12q₉}\)](#) to [\(3.3.1w_{12q₁₂}\)](#), shows that this partial orbit preserves the same column bound. Composing with the output rows in [Equation \(3.3.1w_{12q₈}\)](#) and its two pure-Weyl analogues therefore gives

$$\boxed{|\mathfrak{X}_{0,V}^{\Pi} + \mathfrak{X}_{V,0}^{\Pi}| \leq \frac{C A^r (r+1)^B}{N} \|B\|_F \mathfrak{a}_{\Gamma,\tau}^{1/2}.} \quad (3.3.1\text{w}_{12q_{13d}})$$

No trace norm is taken after the Hilbert realignment, and the first dimension-growing compact Ricci bend remains inside $\mathcal{D}_i^\bullet$.

For completeness, the orthogonal Haar-residual entrance uses the exact root-tested gap estimate. If $\mathcal{K}_Q$ is the Kac transfer on the orthogonal complement of the invariants, then

$$0 \preceq \mathcal{K}_Q \preceq (1 - \gamma_n)I, \quad \sum_{\ell=0}^M \|A\mathcal{K}_Q^\ell x\|^2 \leq \frac{\|A\|^2}{1 - (1 - \gamma_n)^2} \|x\|^2, \quad \gamma_n = \frac{n+2}{2n(n-1)}. \quad (3.3.1w_{12q_{13e}})$$

The factor on the right is $O(n)$. The squared $4/(n-2)$ gain of [Lemma 3.30](#) pays this sole protected-carrier factor in the boundary-planar $g = t_\partial = 1$ ledger; the remaining selected-root row is exactly [Equation \(3.3.1w_{12q_3}\)](#). Hence the Haar-residual channel obeys [Equation \(3.3.1w_{12q_{13d}}\)](#) as well. If no actual separator occurs, the common auxiliary Haar projection gives the same orthogonal split. This exhausts the invariant and residual inputs.

For a strict pair of an aggregate-increment time s and the selected front time u , the remaining fixed Green powers are predictable contractions. If $H_{s,u}(B, C)$ is the resulting terminal contraction, trace duality and [Equation \(3.3.1w_{12q_4}\)](#) give

$$\|H_{s,u}(B, C)\| \leq \frac{CA^r}{N} \|B\|_F \|C\|. \quad (3.3.1w_{12q_{14}})$$

Let

$$\Delta_s = P_{s+1} B_{a_{I_s}} P_{s+1}^\top$$

be the one-step reverse-martingale differences whose partial sums are $\mathfrak{B}_r$ in [Equation \(2.6.5\)](#). By [Equation \(2.6.4\)](#), $\mathbb{E}\|\Delta_s\|_F^2 = 2/[n(n-2)] \leq C/N$. If the terminal endpoint field has energy $\sum_u \mathbb{E}\|C_u\|^2 \leq \mathcal{E}_{\text{end}}$, group the strict sum by s . The resulting summands are reverse-martingale differences. For each s , Cauchy–Schwarz in the at most m values $u > s$, followed by the conditional square bound for Δ_s , gives $CA^{2r}mN^{-3} \sum_{u>s} \mathbb{E}\|C_u\|^2$. Summing in s counts each u at most m times and therefore gives

$$\mathbb{E} \left\| \sum_{s<u} H_{s,u}(\Delta_s, C_u) \right\|^2 \leq \frac{CA^{2r}m^2\mathcal{E}_{\text{end}}}{N^3} \leq \frac{CA^{2r}c^2\mathcal{E}_{\text{end}}}{N}. \quad (3.3.1w_{12q_{15}})$$

The reverse order is its adjoint. The common finite simplex is not replaced by independent intervals: for each strict assignment its Green maps remain in $H_{s,u}$, and the sum in [Equation \(3.3.1w_{12q_{15}}\)](#) is the original ordered pair sum. Restriction to other Haar-separated components is injective, so their already proved renewal bounds introduce no multinomial. This is only a two-time auxiliary estimate: every other anchor, including the marked aggregate–Ricci root, must already be fixed or algebraically eliminated. It does not restore that root time. The restoration is carried out in [Lemma 3.36](#), which uses [Equation \(3.3.1w_{12q_{15}}\)](#) only after the common-Ward telescope has removed the marked position.

At a coincident aggregate-increment/selected-front time in this auxiliary estimate there is only one time sum. The family form [Equation \(3.3.1w_{12q_3}\)](#), with [Equation \(3.3.1w_{12q_{12a}}\)](#), gives the pointwise Hilbert bound $CA^r(Nn)^{-1}\|C_u\|$ after the current plane is averaged. Hence Cauchy–Schwarz in u gives the explicit estimate

$$\mathbb{E} \left\| \sum_u H_{u,u} \right\|^2 \leq \frac{CA^{2r}m\mathcal{E}_{\text{end}}}{N^2n^2},$$

which is no larger than [Equation \(3.3.1w_{12q_{15}}\)](#) for $m = cN$, $c \geq 1$. Thus no martingale-centering assertion is needed at the coincident current increment. There are no other seed/pass supernodes in [Equation \(3.3.1w_{12q_5}\)](#).

The estimates [Equations \(3.3.1w_{12q₂}\)](#) to [\(3.3.1w_{12q_{13e}}\)](#), including the one-normalization incidence argument in the middle of that sequence, prove (i)–(iii). [Equations \(3.3.1w_{12q₁₄}\)](#) to [\(3.3.1w_{12q₁₅}\)](#) record the fixed-anchor conditional inequality. Its endpoint energy is supplied in [Lemma 3.37](#); its legal three-time uses are specified in [Lemma 3.36](#). Notice that the proof does not multiply by a favorable “raw Schur cup” factor: the two original circuit bases were used once in [Equation \(3.3.1w_{12q₁₃}\)](#), and all subsequent operations are Hilbert-space contractions or isometries. $\square$

Lemma 3.33 (Reduction of the packetless low cells). *After the scalar-dimer, internal-hard, and soft-threshold families have been removed, every packetless low component satisfies [Equation \(3.3.1w_{12k}\)](#) except possibly*

$$(g, t_\partial, f_R; u, o, v) = (1, 1, 2; 0, 0, 1), \quad (3.3.1w_{12r})$$

with its complete status, reflection, incidence, and common-simplex orbit retained. The surviving cell is the asymmetric $0 \mid \mathbf{V} + \mathbf{V} \mid 0$ reflected cross, not the $\mathbf{V} \mid \mathbf{V}$ diagonal.

Proof. We include the sharp-power audit because the distinction between the cross and the diagonal is essential. Declare the complete physical cut before the first protected Ricci carrier and let $Z : W \otimes E_- \rightarrow W \otimes E_+$ be the resulting once-normalized trace-class port state. Put

$$\xi_Z = \Re(\text{vec } Z), \quad D_Z = \frac{1}{2}(X^* \otimes I)\xi_Z. \quad (3.3.1w_{12r_1})$$

The proof of [Lemma 3.30](#), before its last application of X , gives

$$\|D_Z\|_2 \leq \|Z\|_1, \quad (L_R \otimes I)\xi_Z = (X \otimes I)D_Z. \quad (3.3.1w_{12r_2})$$

Thus this cut uses the original circuit bases once and no base is added at the bend.

There are two diagonal estimates in these same units. If the root-facing slot is invariant, decompose its reduced density into the four $O(n)$ -summands

$$\mathbf{1} \oplus \mathcal{R} \oplus \mathcal{W} \oplus \Lambda^4 H.$$

At distinct roots the scalar and four-form pieces vanish, while the Ricci and Weyl Hilbert–Schmidt row masses are respectively $Cn^{-2}\|B\|_F^2$ and $Cn^{-1}\|B\|_F^2$. Division by their dimensions $\asymp n^2$ and $\asymp n^4$, followed by the literal- W Green cost $N/[2(n-2)] \leq Cn$, gives $Cn^{-3}\|B\|_F^2$ on both channels. At a coincident root, direct restriction of the five physical row families in [Equation \(3.3.1w_{12j}\)](#) gives squared masses

$$Cn^{-3}, \quad Cn^{-1}, \quad Cn, \quad Cn$$

times $\|B\|_F^2$ on the scalar, Ricci, Weyl, and four-form channels, respectively. Division of the last three by dimensions of orders n^2, n^4, n^4 gives $Cn^{-3}\|B\|_F^2$ on every normalized invariant channel. Consequently, with arbitrary invariant multiplicity,

$$\|\mathcal{J}_B(X \otimes I)D_Z\|_2^2 \leq Cn^{-2}\|B\|_F^2\|Z\|_1^2. \quad (3.3.1w_{12r_3})$$

This is the complete protected zero-front square.

For the same-plane diagonal retain the plane as Hilbert multiplicity. For every symmetric traceless E the five-family source satisfies directly

$$\|\mathcal{F}_{B,i}^{\alpha,\beta}(X_E)\|_2 \leq \frac{C}{\sqrt{N}}\|B\|_F\|E\|_F. \quad (3.3.1w_{12r_4})$$

Indeed $[A_i, X_B] = X_{[i,B]}$, $[A_i, X_E] = X_{[i,E]}$, $\|X_C\|_{\text{op}} \leq 2\|C\|_F$, and $\|A_i\|_{\text{op}} \leq \sqrt{2}$ bound the two anticommutators in [Equation \(3.3.1w_{12q}\)](#), after the physical $N^{-1/2}$ normalization, by the right side

of Equation (3.3.1w_{12r₄}). Squaring, summing the common plane and using Equation (3.3.1w_{12r₂}) yields

$$\sum_i \|\mathcal{F}_{B,i}^{\alpha,\beta}(X_{(D_Z)_i})\|_2^2 \leq \frac{C}{N} \|B\|_F^2 \|Z\|_1^2. \quad (3.3.1w_{12r_5})$$

This is the protected $V \mid V$ diagonal square. Both estimates keep the same port state Z and hence cannot spend two bases on one original circuit.

We now enumerate the low status cells before taking absolute values. The boundary identity proved in Lemma 2.20 gives $g \geq t_\partial \geq 1$. For $f_R = 1$ the separated nuclear entrance, one Ricci bend, and the joint root estimate already have the required one-circuit power. For $f_R = 2$ and $g \geq 2$, the additional Euler power pays the only possible factor n in the separated estimate. It remains to take $g = t_\partial = 1$ and $f_R = 2$. Since $u = 0$ and $o + v/2 < 2$, the complete doubled star statuses are exactly

$$(o, v) = (0, 0), (0, 1), (0, 2), (0, 3), (1, 0), (1, 1). \quad (3.3.1w_{12r_6})$$

The $(0, 0)$ cell is Equation (3.3.1w_{12r₃}). In the $(0, 2)$ and $(0, 3)$ cells select the first opposed pair of residual statuses; it is the diagonal Equation (3.3.1w_{12r₅}), and the possible remaining pass is retained in the shared-resolvent block of Lemma 3.12. In the two cells with $o = 1$, the two reflected polar half-incidences of the one physical seed are S^+ and S^- and give exactly

$$S^- S^+ \preceq \eta_C I, \quad \eta_C = \frac{N-1}{N^2} \leq N^{-1}. \quad (3.3.1w_{12r_7})$$

If a residual status lies between them, the exact coboundary/Riesz block of Lemmas 3.11 to 3.12 is kept between the same two fronts. Trace duality on the predeclared joint port state therefore charges Equation (3.3.1w_{12r₇}) once, without transporting one front separately through the protected carrier. This is the required two-circuit resource. Hence only $(o, v) = (0, 1)$ remains, and its reflected four-rail typing is precisely the cross stated in Equation (3.3.1w_{12r}).

Finally, the low inequality leaves at most four ordered nonroot supernodes. The common unit-circle form of the displayed diagonal and seed estimates has a prefix-Ricci envelope uniform in the physical translation. Parametrize that orbit by its first fused supernode. Then Lemma 3.14, followed by Lemma 3.13, sums the single root translation and the finite Green simplex without a logarithmic loss. Their finitely many status, reflection, incidence, and placement choices cost A_L^r ; restrictions to Haar-separated components are injective. The spatial powers are exactly those in Equation (3.3.1w_{12k}), completing every cell except Equation (3.3.1w_{12r}). $\square$

Lemma 3.34 (Reduction of the packetless low component). *After the scalar-dimer, internal-hard, and soft-threshold families have been removed exactly as above, every restored term in $\mathcal{C}_{k,R}^{(0,\text{low})}$ outside Equation (3.3.1w_{12q₅}) is covered by Lemma 3.33. The complete remaining family is exactly the physical incidence in Lemma 3.32(iii).*

Proof. The exact fixed-column factorization and equal-time fusion in Lemma 3.28 and Lemma 3.29 leave precisely the six low cells listed in Equation (3.3.1w_{12r₆}), together with the weaker $f_R = 1$ and nonplanar $g \geq 2$ branches. Lemma 3.33 proves every one of those branches except Equation (3.3.1w_{12r}). The latter has one and only one V supernode and is therefore the exact scope of Lemma 3.32(iii). The reflection and weak-order sums are retained on both sides of this split, so the two restored families are disjoint and exhaustive. $\square$

Lemma 3.35 (Ordinary exterior renewal around a fixed packetless component). *For every fixed $L < \infty$ there are $A_L < \infty$ and a numerical $B < \infty$ with the following property. Fix $c \geq 1$, $m \leq cN$, and a complete doubled ribbon Γ with total nonconstant slot count $r \leq L \log n$, and keep*

one packetless aggregate-Ricci component $\mathcal{C}_\star$, including its selected $\mathbb{V}$ incidence and reflected root, as an opaque part of the once-normalized positive majorant $\mathbf{a}_\star$. Suppose no unrecorded port-open singleton or packet assignment targets $\mathcal{C}_\star$. Let Ω_{ext} be the complete temporal records of all other Haar-separated components, let R_{ord} be their ordinary selected roots, let $\mathbf{e}_\rho$ be their factored endpoint resources, and let $D_{\text{ext}} \leq 1$ be the product of their retained fractional debits. If $\mathbf{a}_\omega$ is the resulting positive doubled scalar majorant for $\omega \in \Omega_{\text{ext}}$, then

$$\boxed{\sum_{\omega \in \Omega_{\text{ext}}} \mathbf{a}_\omega \leq A_L^r (r+1)^B \mathbf{a}_\star D_{\text{ext}} \prod_{\rho \in R_{\text{ord}}} (m \mathbf{e}_\rho).} \quad (3.3.1w_{12r_8})$$

The fixed special component supplies no endpoint coupon in this estimate.

Proof. Hold the complete doubled spatial contraction, including $\mathcal{C}_\star$ and the original circuit bases, intact. Discard the zero exterior components. By Lemma 3.18, every remaining ordinary exterior component has a nonpassive supernode; otherwise it would consist only of passive singleton supernodes and vanish. Such a supernode is resource-bearing. Indeed, a special root already carries its root resource, and the time-indexed Jacobi child is the explicit $-2N^{-1}W_i$ resource. In the remaining case it is a same-time nonsingleton. All its occurrences have the one physical color t . If they form a repeated naked current label, the exact identity in Lemma 3.8, combined with the Jacobi coefficients before absolute values, leaves only $-2N^{-1}W_i$ and the scalar child; the latter was removed from the temporal word by the convention proved in that lemma. Otherwise centering forces a nonzero pairing-connected component of color t with at least two occurrences, and Theorem 2.16 retains its packet resource $\mu_t = N^{-1}$. Thus every remaining ordinary exterior component contains a resource-bearing hard supernode. Its slot count $r_{\mathcal{C}}$ satisfies $r_{\mathcal{C}} \leq r \leq L \log n$, so the hypotheses of Lemma 3.39 hold. Apply Lemma 3.39 separately to each ordinary exterior component and Lemma 3.40 to its recorded surplus tokens and nonfused singletons. For a port-open singleton fused at an ordinary selected root, run the assignment argument in the proof of Theorem 3.43 with selected root set exactly R_{ord} . That argument uses only the displayed ordinary endpoint resources and treats the rest of the terminal contraction as the fixed projective majorant. It neither selects nor estimates $\mathcal{C}_\star$. Thus the packet-attached restriction in the theorem statement for a *selected* aggregate-Ricci root is never invoked.

Multiplying the component estimates gives the right side of Equation (3.3.1w_{12r₈}). Restriction of a global strict assignment to the Haar-separated components is injective, so there is no intercomponent multinomial. Finally, the left side is already indexed by two-replica scalar records: every cross term between two one-replica exterior choices is one member of Ω_{ext} . The triangle inequality therefore sums the positive doubled majorants linearly; no orthogonality of one-replica endpoint vectors is asserted. $\square$

Lemma 3.36 (Causal restoration of the marked aggregate-Ricci time). *After the coefficient-nonzero Fourier branch has been discharged in the proof of Lemma 3.32, using Equations (3.3.1w_{12q_{12b}}) to (3.3.1w_{12q_{13e}}) for its spatial debit and the translation argument in Lemma 3.33 for its temporal sum, fix, for a fixed $L < \infty$, $c \geq 1$, and $m \leq cN$, a doubled ribbon and circuit state with total nonconstant slot count $r \leq L \log n$, an exterior record, and the complete finite status/reflection orbit in the remaining coefficient-zero part of the packetless cell*

$$(g, t_\partial, f_R; u, o, v) = (1, 1, 2; 0, 0, 1).$$

Let s be the aggregate-increment time, u the unique nonroot $\mathbb{V}$ time, and t the marked aggregate-Ricci root time. At the native selected-time cut, let C_u be the root-open endpoint field and put

$$\mathcal{E}_{\text{end}} := \sum_u \mathbb{E} \|C_u\|^2.$$

Write $M_r = A^{Cr}(r+1)^B$. After retaining and summing the complete marked-root position, both reflections, and every admissible equality face, the strict causal-interval, current/equality, and common-Ward parts satisfy

$$\begin{aligned}\mathbb{E}\|\mathcal{H}_{\text{strict}}\|^2 &\leq \frac{CM_r^C}{N}\mathcal{E}_{\text{end}}, \\ \mathbb{E}\|\mathcal{H}_{\text{current}}\|^2 &\leq \frac{CM_r^C m}{N^2}\mathcal{E}_{\text{end}}, \\ \mathbb{E}\|\mathcal{H}_{\text{Ward}}\|^2 &\leq \frac{CM_r^C m^2}{N^3}\mathcal{E}_{\text{end}}.\end{aligned}\tag{3.3.1w12r9}$$

No conditional centering of the marked root or of C_u is assumed, and the proof introduces no circuit base beyond those already used in the native cut.

Proof. We first restore the marked position algebraically. In one orientation, write R_s for the aggregate occurrence, V_u for the complete selected occurrence, and δ_t for the infinitesimal diagonal action supplied by the pivot. The chronological rule is $\partial_t T_v = \mathbf{1}_{\{v < t\}} \delta_t T_v$. Hence, when $s < u$, Leibniz' rule gives the exact decomposition

$$\partial_t \{R_s, V_u\} = \mathbf{1}_{\{s < t \leq u\}} \{\delta_t R_s, V_u\} + \mathbf{1}_{\{u < t\}} \delta_t \{R_s, V_u\}.\tag{3.3.1w12r10}$$

This is also the boundary identity in [Lemma 3.6](#) applied to the two displayed slots. The reflected orientation is its adjoint with the inequalities reversed. Thus the first term in [Equation \(3.3.1w12r10\)](#) has the single causal order $s < t \leq u$; the second is a common Ward action on the complete already-earlier block.

We next identify the native operator in the strict interval. Expand every gap by [Lemma 3.17](#). The definition of v in [Lemma 3.29](#), the exhaustion in [Lemma 3.34](#), and the opaque-component statement of [Lemma 3.35](#) put the selected V node and the reflected aggregate-Ricci root in the same residual temporal component. A Haar summand on their connecting chronological path would separate those two nodes. Consequently the root-facing active port on that path carries the residual transfer $\mathcal{K}_Q$, where

$$0 \preceq \mathcal{K}_Q \preceq (1 - \gamma_n)I, \quad \gamma_n = \frac{n+2}{2n(n-1)}\tag{3.3.1w12r11}$$

by [Equation \(3.2.10q4\)](#). This is a statement only about that distinguished chronological port; it does not assert $QG_{\Xi}Q = G_{\Xi}$ for the complete high-valent gate.

Here is the precise contextual wiring. Let $\mathcal{H}_{\text{act}}$ be the active root-facing curvature-letter carrier, and let $E_{\text{in}}, E_{\text{out}}$ contain every other selected-time rail. The proof of [Lemma 3.32](#) declares all those ports before estimating the complement and inserts the exact local map

$$G_{\Xi}(B) : \mathcal{H}_{\text{act}} \otimes E_{\text{in}} \longrightarrow W \otimes E_{\text{out}}, \quad G_{\Xi}(B) = N^{-1} \widehat{G}_{\Xi}(B), \quad \|\widehat{G}_{\Xi}(B)\| \leq C\|B\|_F\tag{3.3.1w12r12}$$

in the native chronological slot. The empty transfer immediately before its active input is $\mathcal{K}_Q \otimes I_{E_{\text{in}}}$; the other selected-time rails are static multiplicity. At a distinct reflected root, the literal output carrier is W , whose empty-step transfer is $\kappa_W I_W$ by [Equation \(3.3.1w12d\)](#). This does not identify $G_{\Xi}(B)$ with a smaller local root formula: it substitutes the proved generalized map into the already fixed typed slot.

Put $\ell = u - s$, $a = u - t + 1$, and $b = t - s - 1$. Then $a \geq 1$, $b \geq 0$, and $a + b = \ell$. There is no intervening utilized supernode because the surviving cell has no mixed hard child, has $o = 0$, and has $v = 1$. Multilinearity of the ported contraction therefore gives, before any norm,

$$\boxed{\mathcal{T}_{\ell}(B) = \frac{1}{N} \sum_{a=1}^{\ell} \kappa_W^{\ell-a} \widehat{G}_{\Xi}(B) (\mathcal{K}_Q^a \otimes I_{E_{\text{in}}}).}\tag{3.3.1w12r13}$$

The opposite Fourier mode may pair with the active mode to form a diagonally invariant tensor; it remains in E_{in} . It does not remove the decay in Equation (3.3.1w_{12r₁₁}), because the displayed chronological transfer acts on the active factor and as the identity on that multiplicity. At a coincident reflected root the literal- W factor is absent, leaving a smaller one-denominator word.

Define

$$k_\ell := \frac{C}{N} \sum_{a=1}^{\ell} \|\mathcal{K}_Q\|^a \kappa_W^{\ell-a}.$$

Using Equations (3.3.1w_{12d}) and (3.3.1w_{12r₁₁}),

$$\sum_{\ell \geq 1} k_\ell \leq \frac{C}{N\gamma_n(1 - \kappa_W)} = C \frac{n(n-1)}{(n+2)(n-2)} \leq \frac{6C}{5}. \quad (3.3.1w_{12r_{14}})$$

We include the probabilistic step because the endpoint field need not be centered. Suppose predictable bilinear maps $L_{s,u}$ satisfy

$$\|L_{s,u}(B, C)\| \leq M_r k_{u-s} \|B\|_F \|C\|.$$

For the transported aggregate increments in Equation (2.6.4), orthogonal invariance and the constant Frobenius norm of each coordinate increment give the conditional identities

$$\mathbb{E}(\Delta_s \mid \mathcal{F}_{s+1}^{\leftarrow}) = 0, \quad \mathbb{E}(\|\Delta_s\|_F^2 \mid \mathcal{F}_{s+1}^{\leftarrow}) = \frac{2}{n(n-2)} \leq \frac{C}{N}. \quad (3.3.1w_{12r_{15}})$$

Set $D_s = \sum_{u>s} L_{s,u}(\Delta_s, C_u)$. The native cut is made outside the two displayed bands in Equation (3.3.1w_{12r₁₃}). Sum the marked pivot and average the empty bands first; moving t without crossing s or u then changes only a and b . Thus C_u , the endpoint contractions, and the linear coefficient map $B \mapsto \mathcal{T}_{u-s}(B)$ are $\mathcal{F}_{s+1}^{\leftarrow}$ -measurable. In particular the D_s are reverse martingale differences. Conditioning, summing their squares, and applying scalar Young $\ell^1 * \ell^2 \rightarrow \ell^2$ gives

$$\begin{aligned} \mathbb{E} \left\| \sum_s D_s \right\|^2 &\leq \frac{CM_r^2}{N} \mathbb{E} \sum_s \left(\sum_{\ell \geq 1} k_\ell \|C_{s+\ell}\| \right)^2 \\ &\leq \frac{CM_r^2}{N} \|k\|_1^2 \sum_u \mathbb{E} \|C_u\|^2. \end{aligned} \quad (3.3.1w_{12r_{16}})$$

Equations (3.3.1w_{12r₁₃}) to (3.3.1w_{12r₁₆}) prove the first line of Equation (3.3.1w_{12r₉}); time reversal proves the reflected line. The endpoint $t = u \neq s$ is the zero-subgap endpoint of the same word and is bounded by a one-denominator subseries. If the residual bridge is the remaining rail, its mass is at most $C/(N\gamma_n)$; if the literal- W rail remains, its mass is at most $C/[N(1 - \kappa_W)]$. Both are bounded by the right side of Equation (3.3.1w_{12r₁₄}).

For the common Ward term in Equation (3.3.1w_{12r₁₀}), first sum all $t > \max\{s, u\}$ positions. By Lemma 3.19, internal actions cancel and the remaining action reaches one named exterior leg or finite Green gap. On a finite gap, the exact identity

$$[A, R_M(\mathcal{K})] = \sum_{j=1}^M \sum_{q=0}^{j-1} \mathcal{K}^q [A, \mathcal{K}] \mathcal{K}^{j-1-q} = AR_M(\mathcal{K}) - R_M(\mathcal{K})A \quad (3.3.1w_{12r_{17}})$$

sums the marked position and its internal action position into two boundary incidences. Keep the local half inside the plane-summed selected/root row and compose only the far half afterward.

The finite-kernel part of [Lemma 3.4](#) and the trace-ideal inequality $\|TK\|_1 \leq \|T\|_{\text{op}}\|K\|_1$ show that this composition changes only M_r . The native factor in [Equation \(3.3.1w_{12r₁₂}\)](#) is retained, so the resulting predictable two-time map obeys

$$\|\tilde{H}_{s,u}(B, C)\| \leq \frac{CM_r}{N} \|B\|_F \|C\|. \quad (3.3.1w_{12r_{18}})$$

There is no remaining t index. Applying the already proved elementary two-time estimate [Equation \(3.3.1w_{12q₁₅}\)](#) to [Equation \(3.3.1w_{12r₁₈}\)](#) gives the third line of [Equation \(3.3.1w_{12r₉}\)](#). This use is independent of the packet-bearing aggregate–Ricci renewal theorem.

It remains to classify the current faces. If $s = u$, the two indicators in the Leibniz rule coincide and there is no strict causal interval:

$$\partial_t \mathcal{W}_u = \mathbf{1}_{\{u < t\}} \delta_t \mathcal{W}_u. \quad (3.3.1w_{12r_{19}})$$

The family form of [Equation \(3.3.1w_{12q₃}\)](#), together with [Equation \(3.3.1w_{12q_{12a}}\)](#), bounds the native current gate by $CM_r/(Nn)$. The marked root and the fused current/front node remain in the same residual component, so their one active gap has mass at most

$$\sum_{q \geq 0} \|\mathcal{K}_Q\|^{q+1} \leq \gamma_n^{-1} \leq 2n.$$

After that gap is summed, the effective map has norm CM_r/N . Cauchy–Schwarz only in the remaining common translation therefore gives

$$\mathbb{E} \left\| \sum_u \mathcal{H}_u^{\text{current}}(C_u) \right\|^2 \leq m \sum_u \mathbb{E} \|\mathcal{H}_u^{\text{current}}(C_u)\|^2 \leq \frac{CM_r^2 m}{N^2} \mathcal{E}_{\text{end}},$$

which is the second line of [Equation \(3.3.1w_{12r₉}\)](#). The factor n^{-1} in the current-family gate is the size of that same current increment, so [Equation \(3.3.1w_{12r₁₅}\)](#) is not used a second time.

Finally, $s = t \neq u$ is absent from $D_{< t}$: its current deposit stays inside the marked root. When $s = u = t$, equal-time fusion leaves an aligned root or a previously removed hard block and no marked-root Green fiber. These cases, the strict endpoint $t = u \neq s$, the common-Ward face $s = u \neq t$, and their adjoints exhaust all equality patterns. The native N^{-1} in [Equation \(3.3.1w_{12r₁₂}\)](#) is used once; the endpoint Hilbert mass uses the predeclared circuit bases once; and the separate $O(n)$ root-tested square in [Equation \(3.3.1w_{12q_{13e}}\)](#) is not multiplied into this replacement estimate. This completes the resource audit and the proof. $\square$

Lemma 3.37 (Endpoint energy and chronological closure of the surviving cell). *Consider the complete restored contribution of the cell [Equation \(3.3.1w_{12q₅}\)](#), after all other low cells have been removed as in [Lemma 3.34](#). For every fixed L , $m \leq cN$, $c \geq 1$, and $2 \leq k \leq L \log n$, its squared L^2 norm is at most*

$$C_L n c^{k+1} (k+1)^B A_L^{2k}. \quad (3.3.1w_{12s})$$

The estimate includes both strict orders, the coincident root/front time, the common finite-Green simplex, all statuses and reflections, and the orthogonal-Weingarten sum. The coefficient-nonzero Fourier branch is already included through its separate estimate in the proof of [Lemma 3.32](#); the chronological argument below concerns the sole remaining coefficient-zero gate.

Proof. Put $M_r = A^{Cr}(r+1)^B$. First fix the doubled ribbon record, circuit state, status/reflection orbit, and every temporal record outside the special packetless aggregate–Ricci component. Inside

that component fix the causal order/equality type but leave the three positions (s, t, u) and their two adjacent empty bands unsummed. Let $\omega \in \Omega_{\text{ext}}$ denote the fixed temporal record on all other Haar-separated components and abbreviate its doubled positive majorant by $\mathbf{a}_\omega$.

For each selected-front position u , make the simultaneous native cut of [Lemma 3.32](#) outside the two root-facing bands in [Equation \(3.3.1w_{12r₁₃}\)](#). Let $K_{0,u}$ be the resulting once-normalized root-open complement from [Equation \(3.3.1w_{12q₁₃}\)](#), before those two bands are substituted. Define C_u by vectorizing $K_{0,u}$ and then applying the remaining root-leg permutation, orthogonal projections, and prescribed equivariant or one-half maps from the native matricization. Moving t within its fixed causal cell changes only the two displayed band lengths and hence does not change C_u . Every map used to define the endpoint has already been charged by its operator norm. Therefore [Equation \(3.3.1w_{12q_{13a}}\)](#) gives, uniformly in u ,

$$\mathbb{E}\|C_u\|^2 \leq A^{Cr}(r+1)^B \mathbf{a}_\omega. \quad (3.3.1w_{12s_1})$$

This is an S_2 estimate obtained by applying the S_1 and S_∞ tests to the same complete physical tangle. It neither splits the reflected pair nor introduces a second circuit base. Translating the selected front changes only its physical-time location; it creates no new atom, circuit, or positive presentation cost. Since there are at most m possible front times,

$$E_{\Gamma,\tau} := \sum_u \mathbb{E}\|C_u\|^2 \leq mA^{Cr}(r+1)^B \mathbf{a}_\omega. \quad (3.3.1w_{12s_2})$$

For clarity, the earlier aggregate-Ricci variable is not the unrelated $U_i(B, Z)$ of [Equation \(2.6.7\)](#). It is the reverse-martingale difference

$$\Delta_s = P_{s+1} B_{a_{I_s}} P_{s+1}^\top, \quad \mathbb{E}\|\Delta_s\|_F^2 = \frac{2}{n(n-2)} \leq \frac{C}{N}, \quad (3.3.1w_{12s_3})$$

whose partial sums are $\mathfrak{B}_r$ in [Equation \(2.6.5\)](#). The endpoint C_u is the endpoint field in [Lemma 3.36](#). By [Equation \(3.2.10j\)](#), $r \leq 2k+2 \leq (2L+2)\log n$ for $n \geq e$; thus, after enlarging the fixed logarithmic-degree constant once, the slot hypothesis of that lemma is satisfied. Apply it before restoring any exterior component. Together with [Equation \(3.3.1w_{12s₂}\)](#), it gives for the complete three-time contribution, including the strict causal intervals, all common-Ward positions, every equality face, and both reflections,

$$\begin{aligned} \mathbb{E}\|\mathcal{H}_\omega\|^2 &\leq CM_r^C \left(\frac{1}{N} + \frac{m}{N^2} + \frac{m^2}{N^3} \right) E_{\Gamma,\tau} \\ &\leq CM_r^C (c + c^2 + c^3) \mathbf{a}_\omega \leq CM_r^C c^3 \mathbf{a}_\omega. \end{aligned} \quad (3.3.1w_{12s_4})$$

The last inequality enlarges the numerical constant and uses $c \geq 1$. Thus the fixed- ω contribution is within the required $M_r^C c^3 \mathbf{a}_\omega$ budget. The family under consideration contains a nonroot monomer by the definition of $\mathcal{C}_{k,R}^{(0,+)}$. The doubled aggregate-Ricci root already uses four nonconstant half-slots, so [Equation \(3.2.10j\)](#) implies $k \geq 2$. Since $c \geq 1$, the displayed c^3 is admissible for this minimal cell.

It remains only to restore the records fixed in the first paragraph. Let a be the recurrence-dimer count still carried by the doubled record, and let q_{ext} be the number of ordinary component translations not already represented by the selected endpoint field. In the present $u = o = 0, v = 1$ cell, the unique $\mathbf{V}$ incidence is part of the fixed special component and no unrecorded port-open assignment targets that component. Therefore [Lemma 3.35](#) applies and gives

$$\sum_{\omega \in \Omega_{\text{ext}}} \mathbf{a}_\omega \leq A_L^r (r+1)^B \mathbf{a}_\star c^{q_{\text{ext}}}, \quad (3.3.1w_{12s_5a})$$

because $D_{\text{ext}} \leq 1$ and every ordinary factored endpoint satisfies $m\epsilon_\rho \leq c$.

The reference majorant $\mathfrak{a}_\star$ keeps the special component and the complete once-normalized spatial ribbon intact but excludes the fixed recurrence scalar and the ordinary endpoint resources. In this cell $g = t_\partial = 1$ and $f_R = 2$. Hence the Euler identity [Equation \(2.6.12\)](#), the once-normalized gluing [Equation \(2.6.14b\)](#), and [Definition 2.23](#) give

$$\mathfrak{a}_\star \leq A^r n^{2-g} = A^r n. \quad (3.3.1w_{12s_5})$$

The aggregate root uses four nonconstant half-slots and the unique nonroot V monomer uses two. Every ordinary translated component consumes its own disjoint pair of the remaining nonconstant half-slots, so $q_{\text{ext}} \leq r/2 - 3$. Together with the exact ledger [Equation \(3.2.10j\)](#) gives

$$a + q_{\text{ext}} \leq a + \left(\frac{r}{2} - 3\right) = k - 2. \quad (3.3.1w_{12s_6})$$

The previously factored dimer coefficient γ^a and the exterior endpoint translations are now restored, exactly once, and contribute at most $c^{a+q_{\text{ext}}}$. Consequently

$$c^3 c^{a+q_{\text{ext}}} \leq c^{k+1},$$

with no spare factor of c hidden in the restoration. Across the two replicas, the monomer–dimer tilings themselves have multiplicity at most 2^{2k} by [Lemma 3.16](#); this is absorbed by A_L^{2k} independently of r , including the all-dimer extreme. Their local-state and reflection choices have exponential mass in r . For defect $g = 1$, [Equation \(2.6.13\)](#) contributes at most $A^r(r+1)^B$ records, which is absorbed into $A_L^{2k}(k+1)^B$ because $r \leq 2k+2$. The special-component Green powers were summed only in the exact direct kernel [Equation \(3.3.1w_{12r_{13}}\)](#) and the exact Ward telescope [Equation \(3.3.1w_{12r_{17}}\)](#); they were not replaced by independently truncated rectangles. The exterior doubled records have already been summed linearly by [Lemma 3.35](#). Combining [Equations \(3.3.1w_{12s_4}\)](#) to [\(3.3.1w_{12s_6}\)](#) and only then summing the remaining finite spatial records proves [Equation \(3.3.1w_{12s}\)](#). $\square$

For reference, the marked derivative root is split before any packet or chronological estimate by the exact identity

$$[A_t, D_{<t}] = [A_t, S_{\mathcal{W}, <t}] + [A_t, X_{\mathfrak{B}_{t-1}^-}]. \quad (3.5.1)$$

The two summands are henceforth called the pure-Weyl and aggregate-Ricci branches, respectively. In particular, the next lemma is applied to the second summand with this split still inside the complete Christoffel–Darboux derivative-position sum.

Lemma 3.38 (Packetless aggregate-Ricci shell cancellation). *Form $\mathcal{C}_{k,R}$ by the exact split [Equation \(3.5.1\)](#), retaining the complete Christoffel–Darboux derivative-position sum, reflection orbit, weak chronological sum, and orthogonal-Weingarten sum. After equal-time fusion, fix a complete (p, q, r) atom only to classify its coefficient components. For its selected aggregate-Ricci root R , let $\mathcal{P}_R$ be the set of external multibox q -components assigned wholly to the fresh root color; immediately adjacent current-plane strings remain part of the root and are not in $\mathcal{P}_R$. Let $\mathcal{C}_{k,R}^{(0)}$ be the sum of all terms for which $\mathcal{P}_R = \emptyset$, with the full sums just listed restored before taking the L^2 norm. Remove the completely root-only recurrence-dimer subfamily of [Lemma 3.26](#), and write*

$$\mathcal{C}_{k,R}^{(0,+)} := \mathcal{C}_{k,R}^{(0)} - \mathcal{C}_{k,R}^{\text{dim}}.$$

Thus every term retained in $\mathcal{C}_{k,R}^{(0,+)}$ contains at least one nonroot monomer. Let $\mathcal{C}_{k,R}^{(0,\text{hard})}$ be its subfamily in which the Ricci residual component contains an ordinary hard resource in the sense of [Lemma 3.27](#), and put

$$\mathcal{C}_{k,R}^{(0,\text{soft})} := \mathcal{C}_{k,R}^{(0,+)} - \mathcal{C}_{k,R}^{(0,\text{hard})}.$$

Resolve the internal full-centered hard blocks of this last family as in [Lemma 3.29](#). Route every all-Weyl child and, by [Lemma 3.28](#), every nonzero mixed child back into $\mathcal{C}_{k,R}^{(0,\text{hard})}$; one-box mixed children vanish. Thus the residual soft family has $u = 0$. Let $\mathcal{C}_{k,R}^{(0,\text{low})}$ be the restored sum of exactly those resolved children for which

$$\Delta_0 = o + \frac{v}{2} < f_R.$$

Denote the restored sum of the complementary soft children with $\Delta_0 \geq f_R$ by $\mathcal{C}_{k,R}^{(0,\text{soft},\geq)}$; it is covered by [Lemma 3.29](#) and is not part of the low family. Resources in different Haar-separated components do not enter this classification.

For every fixed L there are $A_L, C_L < \infty$ and a numerical B such that, for $m \leq cN$, $c \geq 1$, and $k \leq L \log n$,

$$\|\mathcal{C}_{k,R}^{(0,\text{low})}\|_2^2 \leq C_L n c^{k+1} (k+1)^B A_L^{2k}. \quad (3.3.1w_{12k})$$

The assertion is only about this intact physical low-debit sum: the complete term has at least one nonroot monomer, while its Ricci component has no internal ordinary hard resource. The Ricci component may itself be all-dimer when the monomer lies behind a Haar separator; that exterior resource remains owned by its own component. It makes no termwise estimate after fixing a ribbon, status, Green coefficient, or Weingarten pairing.

Proof. The definition first removes the scalar-dimer family, then the complete internal-hard family, and finally the soft children with $\Delta_0 \geq f_R$. [Lemma 3.28](#) shows that every nonzero resolved packet child was included in the internal-hard family, so the remaining low branch has $u = 0$ exactly. [Lemma 3.34](#) reduces that complete restored branch to the unique surviving physical incidence, without taking a fixed-ribbon absolute value or changing the circuit normalization. If $k = 1$, that incidence is empty: the aggregate-Ricci root consumes four nonconstant half-slots and the required nonroot monomer consumes two more, whereas [Equation \(3.2.10j\)](#) allows only four. For $k \geq 2$, [Lemma 3.37](#) gives [Equation \(3.3.1w_{12k}\)](#), including the full reflection, status, weak-order, finite-Green, and orthogonal-Weingarten sums. The scalar-dimer, internal-hard, and soft-threshold families are disjoint by construction, so the displayed decomposition is exhaustive. This proves the lemma. $\square$

3.4 Global allocation and pure-Weyl weak-order renewal

3.4.1 Fixed components and token allocation

Lemma 3.39 (Fixed-component ported pure-Weyl renewal with retained debit). *Fix a spatial ribbon map Γ , one labelled residual component $\mathcal{C}$ with $r_{\mathcal{C}} \leq L \log n$ nonconstant slots, one physical anchor, and its complete packet/root channel. A hard ordinary supernode is one fused physical time containing either a nonsingleton packet resource or the time-indexed linear Jacobi child $-2N^{-1}W_i$. The coupled pure-Weyl root is a special hard symbol; the total number of special root symbols is bounded by an absolute constant B_0 . Aggregate Ricci is not present in this ported theorem and is treated separately in the aggregate-Ricci synthesis below.*

For every fixed L there are $A_L < \infty$ and a numerical exponent $B < \infty$ for which the estimate below holds.

Suppose $\mathcal{C}$ contains a hard resource and reserve its chronologically earliest resource-bearing hard

supernode as the endpoint, breaking a same-node tie by the fixed ribbon-slot order. Let

$$\begin{aligned} b_C &= \#\{\text{surplus ordinary hard supernodes}\}, \\ o_C &= \#\{\text{all } S^\pm \text{ singleton supernodes}\}, \\ v_C &= \#\{\text{all } V \text{ singleton supernodes}\}, \\ t_C &= b_C + o_C + v_C. \end{aligned} \tag{3.4.1x}$$

If $\mathbf{e}_C$ is the selected endpoint resource and $\mathbf{a}_{\Gamma,C}$ is the remaining normalized terminal amplitude, then, for $\delta = 1/16$,

$$\sum_{\substack{\text{weak orders, equality blocks, finite gaps,} \\ \text{and typed states on } C}} |\text{Contr}_{\Gamma,C}| \leq A_L^{r_C} (r_C + 1)^B \mathbf{e}_C \mathbf{a}_{\Gamma,C} n^{-\delta t_C}. \tag{3.4.1x_1}$$

The estimate holds at the application horizon $m = cN$ for each fixed $c \geq 1$, uniformly in every finite remaining-time constraint; its constant is independent of c , while the threshold in n may depend on c . Summing the one anchor costs at most m .

Proof. First fix the ordered supernodes, a star-state path, and all local typed choices. Every singleton letter is exactly 0, S^+ , S^- , or V , because the constant-to-constant block of $\mathbf{M}$ is zero. If it shares a physical time with a hard letter, the complete same-time rule fuses it into that hard supernode; otherwise it is counted in o_C or v_C .

Encode the residual gaps by one variable. With $R_z = (I - z\mathcal{K})^{-1}$, the terminal scalar has the form

$$\mathcal{F}_\omega(z) = \text{Contr}_\Gamma[\mathbf{B}_0 X_1 R_z Y_1^* \mathbf{B}_1 \cdots \mathbf{B}_{d-1} X_d R_z Y_d^* \mathbf{B}_d; \Xi], \tag{3.4.1x_2}$$

where $X_j, Y_j \in \{\mathcal{D}, \mathcal{G}, \mathcal{L}\}$, the B_j are complete same-time letters, and Ξ denotes the indivisible packet or root resources. This is a final-scalar identity: every temporary port is reclosed by [Proposition 2.22](#). If the total unused-time budget after the fixed anchor is M , the exact finite simplex is

$$\mathcal{X}_\omega^{(M)} = \sum_{q=0}^M [z^q] \mathcal{F}_\omega(z). \tag{3.4.1x_3}$$

Thus no rectangular gap domain has replaced the chronological simplex.

Along every maximal run of V -vertices choose alternate vertices. Their neighboring resolvent factors are disjoint, and at least $\lceil v_C/2 \rceil$ vertices are selected. Apply [Lemma 3.11](#) to each selected block. This includes a V -vertex immediately adjacent to a hard gate because the outer incidences in [Equation \(3.2.10q7\)](#) are typed. Each $S^\pm$ contributes $N^{-1/2}$. There are at most two $\mathcal{L}$ half-incidences in the two replicas, by [Lemma 3.19](#); all their $b_X b_Y$ factors therefore multiply to at most $(r_C + 1)^2$. The remaining prescribed gates have coefficient mass A^{r_C} . Hence

$$\sup_{|z|=1} |\mathcal{F}_\omega(z)| \leq A^{r_C} (r_C + 1)^2 \mathbf{e}_C \mathbf{a}_{\Gamma,C} N^{-b_C - o_C/2} n^{-v_C/4}. \tag{3.4.1x_4}$$

Here every ordinary nonroot hard time contributes one distinct N^{-1} : it is either a complete monochromatic packet resource or the explicit linear child $-2N^{-1}W_i$. All occurrences of one time/color were fused first, so several combs transporting one token are never charged separately. The scalar child $-\gamma N^{-1}I$ was removed before the temporal decomposition, as stated in [Lemma 3.8](#).

The L^1 norm of the scalar Dirichlet kernel bounds [Equation \(3.4.1x3\)](#) by $C \log(M + 2)$ times [Equation \(3.4.1x4\)](#). At $m = cN$, for each fixed c and all sufficiently large n this logarithm is at most $C \log(n + 2)$, with a numerical C . If $t_C > 0$, then $N \geq n^2/4$ and

$$\log(n + 2) N^{-b_C - o_C/2} n^{-v_C/4} \leq A^{r_C} n^{-t_C/8}. \tag{3.4.1x_5}$$

Indeed the power of n before the logarithm is at least $2b_C + o_C + v_C/4 \geq t_C/4$, and $\log(n+2) \leq Cn^{t_C/8}$. If $t_C = 0$, the component consists of its reserved ordinary packet/Jacobi endpoint, its bounded list of coupled special-root symbols, or both. The reserved endpoint estimate and the intact paired Riesz block, followed by [Proposition 2.22](#), give the same assertion with no dimension debit. Thus [Equation \(3.4.1x5\)](#), with right side one when $t = 0$, is the uniform pre-enumeration estimate in every case.

Let h_C be the total number of chronological supernodes and let $e_C \leq B_0$ count its special root symbols. The above classification is exhaustive, so literally

$$h_C \leq 1 + b_C + o_C + v_C + e_C \leq t_C + B_0 + 1. \quad (3.4.1x_6)$$

Assigning the r_C labelled slots to the ordered supernodes costs at most $h_C^{r_C}$ and already includes all weak equality patterns. Choose H_L so that $t/16 \geq L \log(t + B_0 + 1)$ for $t \geq H_L$. For such t , half of [Equation \(3.4.1x5\)](#) pays this assignment and half leaves $n^{-t/16}$. For $t < H_L$, the number of supernodes is bounded in terms of L , so its r_C th power is absorbed in $A_L^{r_C}$ while the debit is retained. The possible root placements, incidence types, and triples (b, o, v) add only exponential mass. This proves [Equation \(3.4.1x1\)](#); summing the fixed anchor has at most m terms. $\square$

Lemma 3.40 (Token-level rooted allocation). *Fix a fully colored circuit-anchored spatial state and a selected set of q resource-bearing component roots. Let B_{tok} be the number of surplus nonsingleton packet tokens and explicit linear Jacobi resources, including such resources fused into a selected root supernode. Let T_{nf} be the number of singleton supernodes not fused into a selected root. Then, for $r \leq L \log n$,*

$$\boxed{q^{B_{\text{tok}}+T_{\text{nf}}} N^{-B_{\text{tok}}} n^{-\delta T_{\text{nf}}} \leq A_L^r, \quad \delta = \frac{1}{16}.} \quad (3.4.1x_7)$$

Consequently the retained debits sum all assignments of these tokens and nonfused singleton supernodes to the selected roots with exponential cost. The assertion does not include port-open singleton occurrences whose weak-time node is a selected root supernode.

Proof. The occurrence set of every nonsingleton packet token is fixed by the coefficient pairing and declared circuit state. Distinct pairing-connected tokens retain distinct variance factors even when they have the same color. Thus every surplus packet token has one N^{-1} resource, including a token fused into a root equality block. The displayed coefficient of every linear Jacobi child supplies the same resource. [Lemma 3.39](#) retains $n^{-\delta}$ for each nonfused singleton supernode.

Assign the B_{tok} resources and T_{nf} singleton supernodes to the q canonical roots. This costs at most $q^{B_{\text{tok}}+T_{\text{nf}}}$. Since $q \leq r + B_0 \leq L \log n + B_0$ and $N = \binom{n}{2}$,

$$\left(\frac{q}{N}\right)^{B_{\text{tok}}} \left(\frac{q}{n^\delta}\right)^{T_{\text{nf}}} \leq A_L^r.$$

The assigned token occurrence sets and nonfused nodes recover the component sets. Canonical roots and their joint ordered anchor simplex introduce no additional component-order factorial. This proves the claim. A singleton occurrence fused into a selected root has neither of the two displayed debits and was expressly excluded. $\square$

Lemma 3.41 (Vertex-labelled positive-credit presentation). *Fix a circuit state τ , including a maximizing smoothing, all pair-support groups and prime tags, every declared port, and one base on each original $p \vee r$ circuit. For an occurrence v , let $\mathcal{R}_{\tau,v}$ be the finite set of typed nonfront roles*

in which that occurrence appears in the root, pass, fold, and signed-transport daughters. If D is a four-index curvature tensor, put

$$\mathfrak{e}_{\tau,v}(D) := \max_{M \in \mathcal{R}_{\tau,v}} \|M(D)\|_{\text{op}}.$$

The maximum over an empty role set is defined to be zero. Here the raw cups, caps, and circuit bases stay in the declared circuit wiring and are not absorbed into M . Every map which is absorbed into M is a wedge inclusion or projection, a normalized Brauer map, a signed permutation, a full transpose, or one of the finitely many Bianchi transfers. The first four kinds are contractions or isometries in the normalized coordinate Hilbert structures. Each normalized Bianchi transfer is a signed sum of at most a numerical number of coordinate-index isometries. Hence its operator norm is at most a numerical multiple of the four-index Hilbert–Schmidt norm, and consequently

$$\boxed{\mathfrak{e}_{\tau,v}(D) \leq C \|D\|_2}, \quad (3.4.1x7a)$$

where the norm on the right is the invariant four-index Hilbert–Schmidt norm and C is numerical.

Let P be a nonzero connected coefficient packet with $s \geq 2$ occurrences and $c = \mathfrak{c}^*(P)$. Give its vertices possibly different self-adjoint algebraic-curvature tensors D_v ; they need not be Ricci flat. Leave every declared gap and joint context as its formal multilinear slot, or equivalently factor out its prescribed operator/projective cost. Then $c \geq 2$, and there are two front occurrences r_P, s_P and a finite circuit-anchored presentation such that

$$\boxed{\|\zeta_{P,\tau}(\{D_v\})\|_{\mathcal{P}_\tau} \leq C^s n^{c-2} \|D_{r_P}\|_2 \|D_{s_P}\|_2 \prod_{v \notin \{r_P, s_P\}} \mathfrak{e}_{\tau,v}(D_v)}. \quad (3.4.1x7b)$$

Every coefficient occurrence, named gap, joint context, retained interface, and original circuit base occurs once. The root creates one two-front atom; every later prime transports that atom and creates no new front or base.

Proof. We first verify the packet proof with fully heterogeneous vertex labels. At canonical ordered roots $(0, 1)$ the two-box and three-box tensors, with vertex labels retained, are

$$\begin{aligned} Q_{abc,efg}^{01} &= \sum_{d_0, d_1} (D_0)_{ab,cd_0} (x_0^L)_{d_1 d_0} (D_1)_{d_1 e, fg}, \\ H_{abc,efg}^{01} &= \sum_{h, i, j} (D_2)_{fh, ci} (D_3)_{bh, je} (D_4)_{ij, ag}. \end{aligned} \quad (3.4.1x7c)$$

If $(T_D x)_{ijk} = \sum_\ell D_{ij, k\ell} x_\ell$, the first line is, up to the fixed signed leg permutation, $T_{D_0}(x_0^L)^* T_{D_1}^*$. Since $\|T_D\|_2^2 = 4\|D\|_2^2$, its atomic cost is at most

$$4\|D_0\|_2 \|x_0^L\|_{\text{op}} \|D_1\|_2. \quad (3.4.1x7d)$$

If x_0^L is a retained component interface or belongs to a joint attachment, Equation (3.4.1x7d) is read after one projective atom has been substituted in that slot. Before substitution the gap remains formal, and its cost is charged only once by Lemma 2.7. Ricci flatness only removes the trace summand from the range of T_D ; the factorization itself lives on the full three-strand space $H \otimes \Lambda^2 H$ and needs no positivity.

The residual rewrite uses Bianchi at one labelled gate only:

$$(D_4)_{ij, ag} = (D_4)_{ia, jg} - (D_4)_{ja, ig}. \quad (3.4.1x7e)$$

The three closures in the proof of Equation (2.3.9) are cyclic closures of this same identity, so they keep D_2, D_3, D_4 in their own slots. Hence the six root daughters and their coefficient mass remain unchanged. Signed transport moves the complete labelled tuple with its vertices. In pass mode, Equation (2.3.13) holds for every opaque Ξ ; applying Equation (3.4.1x7e) on the residual side therefore keeps the incoming cross atom and all five labelled gates once. Lemma 2.12 and Lemma 2.13 consequently serialize all later fixed tags without inspecting the range of the token.

The distance-hereditary fold proof is already gatewise: at each step it composes with the one actual full-wedge or Bianchi-transfer gate at that vertex. Share mutations are signed four-boundary isometries. Thus its proof also remains valid with vertex labels, with a nonfront occurrence charged in its actual role by $\mathfrak{k}_{\tau,v}$.

Now repeat the support-tree proof of Theorem 2.16. A K_5 or distance-hereditary root activates two carriers and supplies the two displayed fronts; the remaining $c - 2$ carriers cost n each. If every pair group is a singleton, two adjacent tree edges supply the two fronts on three carriers, the initial trace-ideal estimate costs one n , and the remaining $c - 3$ carriers cost n each. Both cases give n^{c-2} . Fixed tags are occurrence-disjoint, support groups partition the remaining boxes, and every pass or fold consumes its own labelled gates once. Lemma 2.7 postpones all raw bends and normalizes each original circuit only at its recorded base. This proves Equation (3.4.1x7b) and every once-only assertion.

It remains only to justify $c \geq 2$. A within-wedge column pair kills the term by skewness. Otherwise start with an allowed smoothing. If it has one circuit, first toggle the smoothing at one box. If some single toggle splits the Euler circuit, we are done. Otherwise the two visits at every box have opposite directions: two visits with the same direction would be separated by the alternative left-to-right reconnection. Record the two visits to each box as a chord on the Euler circle. Two interlacing chords can be toggled together to split the circle. If no two chords interlace, the diagram is laminar and an innermost chord has consecutive endpoints; the pairing edge between those visits then joins the two left darts or the two right darts of one box because the visits have opposite directions. This is the forbidden within-wedge case. Thus a nonzero packet with at least two boxes has an allowed smoothing with at least two circuits. This completes the proof. $\square$

For a coordinate Kac plane write the exact curvature decomposition

$$C_i = P_i - N^{-1}I_W = W_i^\circ + X_i, \quad X_i = X_{B_i}. \quad (3.4.1x7f)$$

Equation (2.6.4) and Equation (3.2.10r) give

$$\|X_i\|_2^2 = \frac{2}{n}, \quad \|W_i^\circ\|_2^2 = \frac{n-3}{n-1}, \quad \mathfrak{k}_{\tau,v}(X_i) \leq Cn^{-1/2}, \quad \mathfrak{k}_{\tau,v}(W_i^\circ) \leq C. \quad (3.4.1x7g)$$

The first inequality in the last pair is deliberately the norm of the actual typed matricization. We do not retype a bent Ricci occurrence as an unbent $W \rightarrow W$ gate and claim the sharper $2/n$ bound.

Corollary 3.42 (One typed Ricci debit per mixed packet). *Under the hypotheses of Lemma 3.41, label every vertex by either W_i° or X_i . If at least one vertex is labelled X_i , then*

$$\|\zeta_{P,\tau}\|_{\mathcal{P}_\tau} \leq C^{|P|} n^{c^*(P)-2} n^{-1/2}. \quad (3.4.1x7h)$$

This debit is pointwise in i and remains multiplicative under the simultaneous one-base-per-circuit contraction.

Proof. Choose one X_i occurrence. If it is one of the two fronts in Equation (3.4.1x7b), use $\|X_i\|_2 = \sqrt{2/n}$; otherwise use Equation (3.4.1x7g) in its actual nonfront role. Every other displayed norm is numerical. For the last assertion, fix all plane labels and one projective atom from every packet before closing a circuit. Lemma 2.7 gives

$$\left| \mathcal{G}_{\Gamma, \tau} \left(\widehat{\bigotimes}_P \zeta_{P, \tau}(i_P) \right) \right| \leq \prod_P \|\zeta_{P, \tau}(i_P)\|_{\mathcal{P}_\tau}. \quad (3.4.1x7i)$$

The values i_P may coincide. Distinct mixed packets still contain distinct actual X_i occurrences, and the raw snake identity uses only the single predeclared base on a circuit. Thus no independence, new variance token, or new base is used. $\square$

3.4.2 Root-fused allocation and global renewal

Theorem 3.43 (Root-fused open-singleton allocation, with packet-attached aggregate roots). *For every fixed L there are $A_L < \infty$ and a numerical B with the following property. Put $N = \binom{n}{2}$ and $r \leq L \log n$. Fix one reflected residual word with its r labelled nonconstant occurrences; one fully declared, once-normalized circuit-anchored spatial atom Γ with positive terminal projective majorant $\mathbf{a}_\Gamma$; all status, incidence, reversal, Jacobi, finite-Green, and nonroot chronological data; a selected root set $R = R_{\text{ord}} \sqcup R_{\text{sp}} = \{\rho_1, \dots, \rho_q\}$ with $|R_{\text{ord}}| \leq r$, $|R_{\text{sp}}| \leq B_0$, and hence $q \leq r + B_0$; and the set F of exactly those still-unrecorded port-open singleton occurrences whose weak-order node is one of the selected root supernodes. Here B_0 is numerical. The special set consists only of complete paired-marked blocks and complete aggregate-Ricci blocks with a distinguished literal- W coupon. At an aggregate-Ricci block, a same-color full-centered occurrence is classified only after the complete coefficient pairing has been fixed: an immediately adjacent string stays inside the root and is covered by Lemma 3.24; an external one-box q -packet vanishes; and an external multibox q -packet is the closed plane-blind scalar of Lemma 3.23. Corollary 3.22 is used only for its constant-zero and exposed-seed bookkeeping, not to bound an aligned $\mathbf{V}$ block before the root cancellation. Every other selected root is ordinary and contains a labelled nonconstant occurrence, giving the displayed bound on $|R_{\text{ord}}|$. If the special set contains an aggregate-Ricci root R , the assertion of this theorem is restricted to the packet-attached subfiber $\mathcal{P}_R(\phi) \neq \emptyset$; the packetless subfiber is outside its scope. If there is no aggregate-Ricci root, no such restriction is imposed.*

For each ordinary root, designate its one endpoint resource $\mathbf{e}_\rho$. If this is a packet token, let $\mathcal{E}_{\text{ord}}$ be the set of packets supplying those tokens; their carrier factors are left bare in $\mathbf{a}_\Gamma$, while the disjoint normalized N^{-1} tokens occur only in $\mathbf{e}_\rho$. These endpoint packets contain their selected hard root supernode, whereas a mobile packet consists only of occurrences in F whose root has not yet been selected. Hence $\mathcal{M} \cap \mathcal{E}_{\text{ord}} = \emptyset$. A linear Jacobi endpoint is treated in the same way, without adding a packet to $\mathcal{E}_{\text{ord}}$. When the special set contains the unique coupled pure-Weyl marked root, its nonmobile marked packet $P_\star$ instead keeps its normalized correlated-front weight

$$w_{P_\star} = n^{-2} \|W_i^\circ\|_2^2 \leq N^{-1} \quad (3.4.1x7j0)$$

inside $\mathbf{a}_\Gamma$; its endpoint $\mathbf{e}_\star$ contains only the remaining star-Casimir factor. For an aggregate-Ricci root R the endpoint is the joint coupon

$$\mathbf{e}_R = \frac{C}{n} \|B\|_F \|B'\|_F; \quad (3.4.1x7j1)$$

the trace-class cost of its crossing four-leg complement is already contained in the simultaneous circuit majorant $\mathbf{a}_\Gamma$. No factor belongs both to $\mathbf{a}_\Gamma$ and to an $\mathbf{e}_\rho$.

The majorant $\mathfrak{a}_\Gamma$ is the positive cost supplied by the fixed simultaneous presentation and [Lemma 2.7](#). It contains every fixed unspent coefficient factor. Put $\mathcal{B} = \mathcal{M} \sqcup \mathcal{E}_{\text{ord}}$, where $\mathcal{M}$ is the mobile packet set. For each $P \in \mathcal{B}$ it retains, up to a numerical factor $C^{|P|}$, the bare carrier factor $n^{c^*(P)}$. It contains neither a mobile allocation debit nor a separately displayed endpoint resource. Thus the sharper $n^{c^*(P)-2}$ two-front factor of [Lemma 3.41](#) is deliberately not booked as a second resource. In particular, $\mathfrak{a}_\Gamma$ is not the possibly cancelling numerical value of an open contraction. This choice is exactly Euler-compatible. If $d = d(\Gamma)$ is the coefficient-occurrence degree and $w_P \leq 1$ denotes every normalized weight retained by a packet outside $\mathcal{B}$, then [Equation \(2.6.12\)](#) and [Equation \(2.6.14b\)](#) give

$$\begin{aligned} \mathfrak{a}_\Gamma &\leq |\omega_\Gamma| \prod_{P \in \mathcal{B}} C^{|P|} n^{c^*(P)} \prod_{P \notin \mathcal{B}} C^{|P|} n^{c^*(P)} w_P \\ &\leq C^r n^{F-4d+\ell_\Gamma-gw_g+\sum_P c^*(P)} \prod_{P \notin \mathcal{B}} w_P = C^r n^{2-g} \prod_{P \notin \mathcal{B}} w_P. \end{aligned} \quad (3.4.1x_{7j})$$

Thus replacing the sharper mobile or factored-endpoint presentation by its bare carrier creates no supercritical power of n .

There is one conditioned refinement for an aggregate-Ricci special root. Let $\mathcal{P}_R(\phi)$ be the external multibox q -packets assigned wholly to R . Fix the complete (p, q, r) atom before opening packet fronts. In the unconditioned convention of [Equation \(3.4.1x_{7j}\)](#), these special-root packets have $w_P = 1$. Replacing each already included bare carrier $n^{c^*(P)}$ by its closed scalar in [Equation \(3.3.1w_{6b}\)](#) gives

$$\boxed{\mathfrak{a}_{\Gamma,\phi}^R \leq A^r n^{2-g} \prod_{P \in \mathcal{P}_R(\phi)} A^{|P|} n^{-c^*(P)}}. \quad (3.4.1x_{7k})$$

This is a replacement inside [Equation \(3.4.1x_{7j}\)](#), not a second copy of its packet carriers. If $\mathcal{P}_R(\phi) \neq \emptyset$, choose one $P \in \mathcal{P}_R(\phi)$ and let $f_R \in \{1, 2\}$ be the number of fundamental $p \vee r$ circuits met by the four root legs. Since every nonzero external multibox packet has $c^*(P) \geq 2$, the joint endpoint [Equation \(3.4.1x_{7j_1}\)](#) yields the sharp pre-anchor ledger

$$\mathfrak{a}_{\Gamma,\phi}^R \mathfrak{e}_R \leq A^r n^{1-g-c^*(P)} \|B\|_F \|B'\|_F \leq A^r n^{1-g-f_R} \|B\|_F \|B'\|_F. \quad (3.4.1x_{7l})$$

No row base, packet front, or second normalization is used in this comparison. If there is no external packet, the joint endpoint lemma still supplies its C/n operator coupon against the actual trace norm of the complement, but it supplies no extra n^{-f_R} circuit debit; that subfiber is not asserted in this theorem.

For an ordered distinct root-anchor vector $\mathbf{t} \in \Delta_q(m)$, let $\Phi_{\Gamma,\omega}(\mathbf{t})$ be the admissible maps $\phi : F \rightarrow R$, including their prescribed same-time/color and typed root attachments. Let $\mathcal{A}_{\Gamma,\omega}(\mathbf{t}, \phi)$ be the fully reclosed, once-normalized scalar, with every ϕ -dependent root block kept intact until the indicated sum is formed. If D_ω is the product of the already proved nonroot debits and $\mathfrak{e}_\rho$ is the selected endpoint resource before its time sum, then

$$\boxed{\sum_{\mathbf{t} \in \Delta_q(m)} \left| \sum_{\phi \in \Phi_{\Gamma,\omega}(\mathbf{t})} \mathcal{A}_{\Gamma,\omega}(\mathbf{t}, \phi) \right| \leq A_L^r (r+1)^B \mathfrak{a}_\Gamma D_\omega \prod_{\rho \in R} (m \mathfrak{e}_\rho)}. \quad (3.4.1x_8)$$

In fact the stronger estimate obtained by moving the absolute value inside the packetwise assignment sum also holds. In that stronger form, every aggregate-Ricci term (which here necessarily has $\mathcal{P}_R(\phi) \neq \emptyset$) is evaluated with $\mathfrak{a}_{\Gamma,\phi}^R$, and the factor $\mathfrak{a}_{\Gamma,\phi}^R \mathfrak{e}_R$ retains the bound [Equation \(3.4.1x_{7l}\)](#) through the anchor and record sums.

Proof. Fix the column pairing of Γ . Make a graph whose vertices are its coefficient boxes and whose edges join boxes when a column-pairing edge joins their four column darts. Adjoin a boundary vertex for any selected root which itself carries such darts. Haar integration is performed separately at each physical time, so both ends of every column-pairing edge have the same time/color. Hence color is constant on every connected column packet.

First condition on the occurrences assigned to the bounded set of special paired-marked or aggregate-Ricci roots. If their number is at most B_0 , all subset and target choices cost at most $(1 + B_0)^{|F|}$. A column packet split between this subset and its complement, or between incompatible special-root colors, is inadmissible and contributes zero. For an aggregate-Ricci root, retain only the packet-attached subfiber and fix the complete (p, q, r) atom before opening any packet front. Keep immediately adjacent strings inside the root and apply [Lemma 3.24](#). Every external one-box packet is zero. Every external multibox packet assigned wholly to this root is a closed scalar independent of the fresh plane by [Lemma 3.23](#); extract it before the root-plane sum. The remaining row and temporal wiring is one plane-independent joint trace-class complement, so [Lemma 3.25](#) gives [Equation \(3.4.1x_{7j1}\)](#). The exact replacement of its bare carrier is [Equation \(3.4.1x_{7k}\)](#), and [Equation \(3.4.1x_{7l}\)](#) pays every nonzero external-packet cell. Complete paired-marked blocks retain their previously proved local special-root coupon. Thus all special blocks stay intact.

Taking the triangle inequality over the special-root packet assignments does not erase this refinement. A connected column packet moves as a unit, has at most B_0 special targets, and cannot be split across the special set and its complement. Since the packet occurrence sets are disjoint and their total size is at most r , the product of all target choices and the factors $A^{|P|}$ in [Equation \(3.4.1x_{7k}\)](#) is at most A_0^r . We retain the displayed $n^{-c^*(P)}$ for every nonzero packet sent to an aggregate-Ricci root. This proves the asserted termwise conditioned version before the ordinary-root allocation begins.

Consider the remaining ordinary-root packets. If a packet contains a box whose physical time is already recorded, or meets an ordinary root boundary, its color and every assignment in it are forced. If it is not preassigned and meets F , then every box outside F would have supplied a recorded color, so the whole packet lies in F . All its occurrences must move together to one of at most q ordinary roots.

An unpreassigned one-box packet is zero before any other port is closed. Indeed the three perfect matchings of its four column darts give 0, $\text{Tr } C_i$, and $-\text{Tr } C_i$, respectively. For a packet P with at least two boxes, expand [Equation \(3.4.1x_{7f}\)](#) only at its vertices. Its all- W° child retains the one N^{-1} pure-Weyl packet resource proved by [Theorem 2.16](#), equations [Equations \(2.6.10\) to \(2.6.14\)](#), and [Proposition 2.21](#); explicitly, its normalized front weight is $\beta_i^2 = N^{-1} \|W_i^\circ\|_2^2 \leq N^{-1}$. If $c = c^*(P)$, its absolute packet estimate compares with the bare mobile factor in $\mathfrak{a}_\Gamma$ as

$$C^{|P|} n^{c-2} \|W_i^\circ\|_2^2 = C^{|P|} n^c [n^{-2} \|W_i^\circ\|_2^2] \leq C_0^{|P|} n^c N^{-1}. \quad (3.4.1x_{8b'})$$

Every other child is mixed and has the relative $n^{-1/2}$ debit of [Corollary 3.42](#); indeed $n^{c-2} n^{-1/2} = n^c n^{-5/2} \leq n^c n^{-1/2}$. For comparison, the unsplit full-centered packet obeys

$$C^{|P|} n^{c-2} \|C_i\|_2^2 \prod_{v \notin \{r_P, s_P\}} \mathfrak{k}_{\tau, v}(C_i) \leq C_0^{|P|} n^c, \quad (3.4.1x_{8b''})$$

and is therefore kept at unit relative cost; the favorable universal n^{-2} is discarded rather than spent as a variance coupon. The $2^{|P|}$ resolutions and the at most q ordinary-root choices therefore cost

$$C^{|P|} q \left(N^{-1} + n^{-1/2} \right) \quad (3.4.1x_{8c})$$

relative to $\mathfrak{a}_\Gamma$.

We record explicitly why equal physical colors do not invalidate this step. First expose all fixed- $\mathfrak{q}$ darts and column packets. For every finite mixed projective presentation, apply Equation (3.4.1x7i) atomwise with each intervening factor $D_\nu(i)$ charged by its declared operator/projective cost, sum the atoms, and only then apply Lemma 2.7 once to the complete simultaneous contraction. Thus the mixed debits multiply pointwise without creating a second circuit base. An all-Weyl packet keeps its correlated endpoint resource under an arbitrary bounded same-color environment: if $L(i), R(i)$ are its two front maps and $D(i)$ is the fully exposed intervening environment, let P_0, P_3 be the conditional-expectation projections onto the constant vectors in the two endpoint plane-star spaces. For the associated multiplication maps,

$$\|P_0 M_L\|^2 = \|\mathbb{E}_i L(i) L(i)^*\|, \quad \|M_R P_3\|^2 = \|\mathbb{E}_i R(i)^* R(i)\|.$$

Operator Cauchy–Schwarz on these same spaces therefore gives

$$\|P_0 M_L D M_R P_3\| \leq \|\mathbb{E}_i L(i) L(i)^*\|^{1/2} \|D\|_{L^\infty(\text{op})} \|\mathbb{E}_i R(i)^* R(i)\|^{1/2}. \quad (3.4.1x8d)$$

Thus mixed letters depending on the same plane stay inside one common environment and do not erase the pure token. Distinct all-Weyl column packets retain their distinct fronts by the fixed-column simultaneous spatial factorization of Proposition 2.21, not by probabilistic independence. No full-centered variance is assigned to a mixed component.

If b ordinary packets are mobile, multiplying Equation (3.4.1x8c) and using the single terminal contraction gives

$$C^r \left[q \left(N^{-1} + n^{-1/2} \right) \right]^b \mathfrak{a}_\Gamma D_\omega \prod_{\rho \in R} \mathfrak{e}_\rho. \quad (3.4.1x8e)$$

Since $q \leq r + B_0 \leq L \log n + B_0$, the bracket is uniformly bounded; its $b \leq r$ powers are absorbed into A_L^r . Forced packets have no membership multiplicity, zero packets vanish, and the special-root choices were already exponential. Lemma 3.40 pays every previously recorded surplus token and nonfused singleton assignment.

Finally

$$|\Delta_q(m)| \prod_{\rho \in R} \mathfrak{e}_\rho \leq \prod_{\rho \in R} (m \mathfrak{e}_\rho).$$

All status, incidence, reversal, Jacobi, and finite-Green choices have the already recorded cost $A_L^r(r+1)^B$. Combining this with Equation (3.4.1x8e) proves Equation (3.4.1x8), even after the triangle inequality is taken over the packet assignments.

For clarity, applying Lemma 3.41 to an unsplit C_i supplies only the unit relative cost already present in $\mathfrak{a}_\Gamma$; its two fronts are not a new variance coupon. Rebranding that existing topology as N^{-1} would double-spend the Euler/circuit normalization. The N^{-1} above is the independently proved all-Weyl resource, while $n^{-1/2}$ comes from an actual displayed X_i slot. Thus the two resources are disjoint in the fixed-column presentation. $\square$

Theorem 3.44 (Grouped ported pure-Weyl packet/root chronological renewal). *Fix a reflected two-replica monomer–dimer word with $r \leq L \log n$ nonconstant slots and one rooted spatial ribbon map Γ . Keep every same-time packet as one fused physical supernode, every packet token once, the coupled pure-Weyl root, and the complete finite-gap and weak chronological sums. Remove the scalar Jacobi child $-\gamma N^{-1} I$ before forming residual components. Fix a reduced temporal record ω which records all data listed in Theorem 3.43 except the ordered root anchors $\mathbf{t}$ and the root-fusion map ϕ . Put*

$$\mathcal{X}_{\Gamma, \omega}(\mathbf{t}) := \sum_{\phi \in \Phi_{\Gamma, \omega}(\mathbf{t})} \mathcal{A}_{\Gamma, \omega}(\mathbf{t}, \phi).$$

Then

$$\sum_{\mathbf{t} \in \Delta_{q(\omega)}(m)} |\mathcal{X}_{\Gamma, \omega}(\mathbf{t})| \leq A_L^r (r+1)^B \mathbf{a}_{\Gamma} D_{\omega} \prod_{\rho \in R(\omega)} (m \mathfrak{e}_{\rho}). \quad (3.4.1x_9)$$

Here $\mathbf{a}_{\Gamma}$ retains the positive, already normalized fixed-ribbon projective majorant and all unspent packet weights; the phrase “unspent” excludes the mobile all-Weyl and mixed debits already used in [Theorem 3.43](#). Summing the reduced records is paid by [Lemma 3.40](#) and the exponential local state count. Crucially, the absolute value is outside the complete ϕ -sum. No extra n , $\log n$, short-gap scale, Bell factor, matching factorial, or component multinomial occurs.

Proof. Expand every gap by [Lemma 3.17](#). By [Lemma 3.18](#), every passive-only residual component is zero before terminal gluing. Hence every nonzero component has a hard resource. For the fixed record ω , the fixed-component theorem has already summed all local weak orders and finite gaps and produced exactly D_{ω} ; every typed temporal gate is reclosed by [Proposition 2.22](#). The only data deliberately omitted from ω are the ordered anchors and the assignment of each port-open root-fused singleton occurrence to a selected root. [Theorem 3.43](#), applied without separating that assignment sum, is therefore exactly [Equation \(3.4.1x₉\)](#).

For completeness, before ω is fixed choose in every nonzero component its earliest resource-bearing hard supernode, with the fixed slot order breaking ties; the paired marked root is one root object. There are at most 2^{r+B_0} root-set choices. [Lemma 3.40](#) pays the assignments of every surplus packet/Jacobi token and every nonfused singleton supernode while retaining their debits. Incidence, state, and bounded special-symbol choices have exponential mass, and $\prod_C (r_C + 1)^B \leq 2^{Br}$. Thus summing the reduced records only enlarges A_L ; it does not move the absolute value inside the root-fusion fiber or reuse the fixed-ribbon amplitude. $\square$

3.5 Aggregate-Ricci synthesis and marked-shell closure

3.5.1 Aggregate-Ricci renewal

Theorem 3.45 (Operator-level aggregate-Ricci renewal). *Let $\mathcal{C}_{k,R}$ be the part of the marked shell obtained by the exact split [Equation \(3.5.1\)](#) inside the complete Christoffel–Darboux derivative-position sum. In the second branch do not decompose undifferentiated full-centered deposits $\mathcal{C}_{v_s}$ before equal-time fusion and fixed-column packetization. At the root-fused allocation step resolve only vertices of unpreassigned ordinary mobile fixed-column packets as in [Equation \(3.4.1x_{7f}\)](#); deposits assigned to a distinguished or special Ricci block remain intact. Retain each complete derivative/status orbit and its reflection before taking an absolute value. Then, for $m \leq cN$, $c \geq 1$, and $k \leq L \log n$,*

$$\|\mathcal{C}_{k,R}\|_2^2 \leq C_L n c^{k+1} (k+1)^B A_L^{2k}. \quad (3.5.2)$$

Here A_L, B are independent of c and n .

Proof. Define the complementary packet-bearing branch by the exact identity

$$\mathcal{C}_{k,R}^{(\geq 1)} := \mathcal{C}_{k,R} - \mathcal{C}_{k,R}^{(0)};$$

equivalently, it is the restored sum of the terms for which $\mathcal{P}_R \neq \emptyset$ in [Lemma 3.38](#). Thus the five summands below are defined before they are used and form a disjoint exhaustive split. Split

$$\mathcal{C}_{k,R} = \mathcal{C}_{k,R}^{\dim} + \mathcal{C}_{k,R}^{(0, \text{hard})} + \mathcal{C}_{k,R}^{(0, \text{soft}, \geq)} + \mathcal{C}_{k,R}^{(0, \text{low})} + \mathcal{C}_{k,R}^{(\geq 1)}$$

according to the classification in [Lemma 3.38](#). [Lemma 3.26](#) gives the required bound for the first summand, [Lemma 3.27](#) gives it for the second, and [Lemma 3.29](#) gives it for the third; the packetless lemma gives it for the fourth. The elementary inequality $\|x_1 + \dots + x_5\|_2^2 \leq 5 \sum_{j=1}^5 \|x_j\|_2^2$ reduces the theorem to the last summand. The remainder of this proof treats $\mathcal{C}_{k,R}^{(\geq 1)}$ and is unconditional.

Relative to the marked time t , the finite- N centering is exactly

$$D_m - (\gamma + 1)I = \sum_{s \neq t} \mathbf{C}_{v_s} + (\mathbf{P}_{v_t} - I), \quad \gamma = (m - 1)/N. \quad (3.5.3)$$

Thus [Equation \(3.5.1\)](#) is an exact split and every nonroot letter in the Ricci branch is a full centered deposit. The current letter lies in the two-dimensional algebra spanned by I and $\mathbf{C}_{v_t}$ and is kept inside the marked root.

Choose the Ricci root reflection-equivariantly: select the orbit of the earliest physical-time Ricci interface and distinguish its two reflected members together. Expose pivots from latest to earliest before fusing equal times. The Ward identity moves every fresh-plane action on an exterior factor to a named boundary incidence, so all flips, Brauer maps, and exterior operators in the remaining current-plane block are fixed with respect to that plane.

We now verify the physical joint-boundary hypothesis in [Lemma 3.25](#). Fix Γ , the complete orthogonal-Weingarten pair (p, q) , the trace successor pairing r , the status orbit, and every prescribed gate, before opening a coefficient packet front. The aggregate-Ricci passive boundary has no q -dart. Hence every external q -component assigned wholly to the Ricci root is a closed coefficient scalar. A one-box component vanishes, while a multibox component P is, by [Lemma 3.23](#), a scalar $z_{P,q}$ independent of the fresh root plane. Extract all such scalars from the normalized root-plane sum before choosing projective atoms at the remaining ports. An immediately adjacent current-plane string is not extracted: it stays inside the complete root source and is treated by [Lemma 3.24](#).

After this exact coefficient contraction, insert the normalized boundary maps and close every ancillary circuit leg except the four ordered physical passive rails in [Equation \(3.3.1w_{12a}\)](#). [Lemma 2.7](#), with one base on each original circuit, supplies one joint complement

$$K_\eta \in \mathcal{S}_1(\mathcal{R}^{\text{in}}, \mathcal{R}^{\text{out}}), \quad \|K_\eta\|_1 \leq \text{cost}(\eta). \quad (3.5.3a)$$

Reflection is already encoded by the exchange of the minus input and output rails. In particular, K_η is neither replaced by two single-replica kernels nor required to admit a replica-projective factorization. At distinct reflected root times apply [Lemma 3.25](#) to this one K_η and the finite or unit-circle Green rail. At coincident times apply its shared-plane version before attaching K_η .

If a marked action meets the last gap, the exact identity

$$[A, R_s] = \sum_{\ell=1}^s \sum_{q=0}^{\ell-1} \mathcal{K}^q[A, \mathcal{K}] \mathcal{K}^{\ell-1-q} = AR_s - R_s A \quad (3.5.4)$$

and [Lemma 3.19](#) move it to one boundary $\mathcal{L}$ half-incidence. Its local $d\rho(a_i)$ half remains inside the complete root/Riesz row while the fresh plane is summed; only the far formal port is substituted afterward. There is at most one such half in each replica, so [Lemma 3.4](#) changes the joint row square by at most $(r + 1)^2$ and introduces no gap length, logarithm, or power of n . Indeed, if the already plane-summed coefficient row is $M = (M_s)_s$ and the delayed far half is $T = XRY^*$, then $\widetilde{M} = MT$ and

$$\sum_s \widetilde{M}_s \widetilde{M}_s^* = MTT^*M^* \preceq \|T\|_{\text{op}}^2 MM^*, \quad \|T\|_{\text{op}} \leq b_X b_Y, \quad (3.5.4a)$$

where $b_{\mathcal{G}} = 1$, $b_{\mathcal{G}} = \sqrt{2}$, and $b_{\mathcal{L}} \leq r + 1$. Applying this calculation in both reflected rows and then using the joint Haagerup estimate proves the stated factor. If the delayed far port has already

become part of the terminal trace-class complement, keep it there instead: composition gives $\|TK\|_1 \leq \|T\|_{\text{op}}\|K\|_1$, and Equation (3.3.1w_{12c}) applies directly. Thus neither the operator-input nor the trace-ideal ancillary typing introduces a dimension factor. Only after this local composition do we apply the tensor and pinch estimates of Lemma 3.21; no assertion is made for arbitrary plane-dependent left and right environments.

Fuse equal physical times. Lemma 3.21 reduces each nonroot fused block to a passive singleton, a favorable scalar, or one hard A_L^r/N resource. Expand every remaining gap by Lemma 3.17. Lemma 3.18 kills every passive-only component termwise. On each surviving nonroot component select alternate internal V vertices and apply Lemma 3.11; every $S^\pm$ vertex costs $N^{-1/2}$. Exactly as in Equations (3.4.1x₄) to (3.4.1x₆), a component with b, o, v such nodes has, before its labelled slots are assigned, the factor

$$A_L^r(r+1)^2 N^{-b-o/2} n^{-v/4},$$

and the retained fractional debit pays all weak orders and equality assignments. This use of the whole-simplex estimate occurs only after the two-member Ricci reflection orbit has been formed. The Green gap carrying that orbit is not removed from the simplex: in the single generating function Equation (3.4.1x₂) it is the unit-circle block R_z , and Equation (3.3.1w_{12c}) supplies its joint $Cn^{-1}\|B\|_F\|B'\|_F\|K_\eta\|_1$ coupon uniformly for $|z| = 1$. Hence the supremum estimate Equation (3.4.1x₄) applies to the complete coupled simplex without a sequential Green elimination or a replica factorization. Root-time fusion has at most 2^r choices and is absorbed in A_L^r . If the two marked replica times coincide, use the shared-plane statement of Lemma 3.25; it gives the same C/n bound with the common plane left intact.

Current-plane deposits at a Ricci root obey an exact trichotomy. Immediately adjacent strings remain root-internal and use Lemma 3.24; an external one-box q -packet is zero; and an external multibox packet is the plane-blind closed scalar of Lemma 3.23. Thus the V block of an aligned string is never bounded by one before the root cancellation, and no termwise claim $\Psi(C_i Z) = 0$ is used. In the multibox case the fixed atom is estimated with the conditioned majorant $\mathbf{a}_{\Gamma, \phi}^R$ in Equation (3.4.1x_{7k}), and its product with the Ricci endpoint is bounded by Equation (3.4.1x_{7l}); the larger unconditioned $\mathbf{a}_\Gamma$ is not substituted back at this step. Ordinary component roots cost N^{-1} ; the Ricci component instead carries the intact $n^{-1}\|B\|_F\|B'\|_F$ coupon. Group the still-open singleton occurrences fused into selected root supernodes by their root assignment and apply Theorem 3.43 before taking an absolute value. Lemma 3.40 pays every nonfused singleton and surplus hard-token assignment. Restriction of a global time assignment to its Haar-separated components is injective, so no further intercomponent multinomial occurs.

The reverse matrix martingale obeys, by Hilbert-space Burkholder,

$$\left(\mathbb{E} \max_{u \leq cN} \|\mathfrak{B}_u\|_F^p \right)^{1/p} \leq C \left(\sqrt{cp} + \frac{p}{n} \right), \quad p \geq 2. \quad (3.5.5)$$

Indeed its predictable Frobenius square function is at most Cc and each increment is at most C/n . Thus the translated paired Ricci root costs at most $m(C/n)Cc \leq Cc^2n$.

Let a be the total recurrence-dimer count and q the number of nonzero residual components, including the Ricci component. The latter uses four nonconstant half-slots, and every other endpoint resource uses two disjoint half-slots. Hence $q \leq r/2 - 1$ and, by Equation (3.2.10j), $a + q \leq k$. Each ordinary component translation costs $m/N \leq c$ and each dimer costs $\gamma \leq c$. Therefore

$$c^2 n c^{a+q-1} \leq n c^{k+1}.$$

All monomer-dimer, two-term fusion, reflection/status, and incidence choices are exponential in k , while the two conserved derivative halves contribute only a fixed power of $k + 1$. This proves Equation (3.5.2). $\square$

3.5.2 Intact marked-shell closure

Theorem 3.46 (Intact marked-shell closure). *For every fixed L there are numerical A_{sh}, C, r such that*

$$\|\mathcal{C}_k\|_{L^2(\Omega; \mathfrak{so}(n))} \leq C\sqrt{n} c^{(k+1)/2} (k+1)^r A_{\text{sh}}^k, \quad 1 \leq k \leq L \log n. \quad (3.5.6)$$

In particular, the first shell has squared energy $O(n)$ for fixed c . This assertion is denoted $\text{MS}_{\text{intact}}$.

Proof. For $k = 1$, the assertion follows directly from [Proposition 3.5](#); in particular, no packetless-shell, weak-order, or ribbon estimate is invoked. Assume henceforth that $2 \leq k \leq L \log n$.

Make the exact split $\mathcal{C}_k = \mathcal{C}_{k,W} + \mathcal{C}_{k,R}$ from [Equation \(3.5.1\)](#). [Theorem 3.45](#) already gives [Equation \(3.5.6\)](#) for $\mathcal{C}_{k,R}$. It remains to treat the pure-Weyl branch. Expand the two replicas of $\langle \mathcal{C}_{k,W}, \mathcal{C}_{k,W} \rangle$ by [Equation \(3.2.7\)](#), but retain the complete sum of derivative positions in each replica. Next expand every Q_s by [Lemma 3.16](#). If a is the total number of dimers and ρ the total number of nonconstant slots, then

$$a + \rho/2 = k + 1. \quad (3.5.7)$$

There are at most 2^{2k} monomer–dimer choices. At this stage no coefficient packet, marked position, or reflected partner has been estimated separately.

Before residual components are formed, fuse every equal physical time and apply [Lemma 3.8](#). Remove the scalar child $-\gamma N^{-1}I$ from the temporal word: γ remains in the monomer–dimer coefficient and the extra N^{-1} is favorable but has no anchor. Only $-2N^{-1}W_i$ remains as a time-indexed Jacobi hard node.

For each fixed spatial ribbon map Γ , apply [Theorem 3.44](#) once. Its proof performs the exact Haar/residual expansion, annihilates every passive-only component, sums all weak orders and equality blocks of the surviving packet/root components, eliminates their finite gaps, and sums their varying labelled component sets and anchors. Thus no equality partition, intercomponent multinomial, or second Green summation is introduced here.

Every ordinary surviving component selects either its earliest nonsingleton packet token or its time-indexed linear Jacobi resource; in both cases its factored endpoint weight is $\mathfrak{e}_\rho = N^{-1}$; if it is a packet token, that packet’s carrier is bare in $\mathfrak{a}_\Gamma$. All other packet tokens are kept for terminal closure. The unique marked component instead keeps the coupled reflected root. Its correlated-front weight $w_{P_\star} \leq N^{-1}$ remains in $\mathfrak{a}_\Gamma$ and is not part of the endpoint resource. [Lemma 3.7](#) expresses the remaining derivative action in the $\mathscr{D}, \mathscr{G}, \mathscr{L}$ channels. The adjoint Casimir on a degree- $O(\rho)$ boundary is at most $C(\rho+1)^2 n$, so the disjoint marked endpoint factor is

$$\mathfrak{e}_\star \leq C(\rho+1)^2 \frac{n}{N}. \quad (3.5.7a)$$

No passive-only endpoint survives [Lemma 3.18](#).

Every residual component has one free physical translation, with at most $m = cN$ choices. Thus each ordinary endpoint resource and its translation cost at most c , while the marked component costs at most $C(\rho+1)^2 cn$. If q is the number of residual components, their distinct endpoint resources consume disjoint pairs of nonconstant half-slots, so $q \leq \rho/2$. [Equation \(3.5.7\)](#) and $\gamma \leq c$ give the exact ledger

$$\gamma^a c^q \leq c^{a+q} \leq c^{k+1}. \quad (3.5.8)$$

The dirty root and its at most two $\mathscr{L}$ half-incidences were kept inside the paired finite-Green form by [Lemma 3.19](#) and [Lemma 3.39](#); no standalone row norm was taken. Every nonscalar repeated-label child remains in its intact packet and is covered by [Theorem 2.24](#). The scalar Jacobi child has already been removed, and the linear child is already one of the ordinary endpoint or surplus resources counted by the chronological theorem.

We next sum the static ribbon maps. Let $\mathcal{B} = \mathcal{M} \cup \mathcal{E}_{\text{ord}}$ be the disjoint bare packet set declared in [Theorem 3.43](#). There is no independent plane-label sum over $P \in \mathcal{M}$: each such label was assigned to an already selected root, and its at most q choices and its root anchor were already summed in [Equation \(3.4.1x₈\)](#). The normalized token of every $P \in \mathcal{E}_{\text{ord}}$ is likewise present only in its factored endpoint $\mathfrak{e}_\rho$. Therefore [Equation \(3.4.1x_{7j}\)](#), or equivalently [Equation \(2.6.12\)](#) and [Equation \(2.6.14b\)](#), gives the exact mixed packet ledger

$$\begin{aligned} |\omega_\Gamma| \prod_{P \in \mathcal{B}} n^{c^*(P)} \prod_{P \notin \mathcal{B}} n^{c^*(P)-2} \|C_P\|_2^2 \|C_P\|_{\text{op}}^{m_P-2} \\ \leq C^\rho n^{2-g} \prod_{P \notin \mathcal{B}} \left(n^{-2} \|C_P\|_2^2 \|C_P\|_{\text{op}}^{m_P-2} \right). \end{aligned} \quad (3.5.8a)$$

The marked packet is outside $\mathcal{B}$, because its physical time/color is recorded at the coupled root. Its factor in the last product is $w_{P_\star} \leq N^{-1}$ from [Equation \(3.4.1x_{7j0}\)](#). All other displayed normalized weights are at most one. Consequently

$$\mathfrak{a}_\Gamma \leq C^\rho n^{-g}. \quad (3.5.8b)$$

This is the single global-root normalization; it is not also placed in $\mathfrak{e}_\star$. Combining [Equation \(3.5.7a\)](#), the N^{-1} ordinary endpoints, and all anchor sums gives the requested disjoint identity

$$\begin{aligned} \mathfrak{a}_\Gamma \prod_{\rho \in R} (m\mathfrak{e}_\rho) &\leq C^\rho n^{-g} \left(C(\rho+1)^2 \frac{mn}{N} \right) \left(\frac{m}{N} \right)^{q-1} \\ &\leq C^\rho (\rho+1)^2 n c^q n^{-g}. \end{aligned} \quad (3.5.8c)$$

Thus the bare factors introduce neither $n^{2|\mathcal{B}|}$ nor an extra leading N , and every endpoint and packet-front weight is used once. After the leading nc^q factor is separated, a fixed defect g supplies $A^\rho n^{-g}$, and [Lemma 2.20](#) gives at most $A^\rho \rho^{Cg}$ maps. Since $\rho = O_L(\log n)$,

$$\sum_{g \geq 0} \left(\frac{\rho^C}{n} \right)^g$$

is bounded for all sufficiently large n ; the finitely many remaining n are absorbed by enlarging the numerical constant. The choices of incidence type, root exit, reflection, and ordered separator boundaries are all finite per slot and contribute $A^\rho (\rho+1)^{O(1)}$.

Collecting the preceding estimates, for a numerical r_0 and a constant A_{sh} depending only on the fixed degree range L ,

$$\mathbb{E} \|\mathcal{C}_{k,W}\|^2 \leq C n c^{k+1} (k+1)^{2r_0} A_{\text{sh}}^{2k}.$$

The same form holds for $\mathcal{C}_{k,R}$ by [Theorem 3.45](#). Hence $\|X+Y\|_2^2 \leq 2\|X\|_2^2 + 2\|Y\|_2^2$ proves the displayed estimate for $\mathcal{C}_k$. The constants are independent of $c \geq 1$; for each fixed c only the threshold in n changes in the martingale good event. Taking square roots proves [Equation \(3.5.6\)](#) for $k \geq 2$; together with [Proposition 3.5](#) this proves the theorem and its final assertion. $\square$

3.6 Inverse expansion and score energy

By [Theorem 3.46](#), on the regular set, [Equation \(3.1.7\)](#), $M = 2D$, $j_t = \sqrt{2}v_t$, and [Equation \(3.2.9\)](#) give

$$b = \frac{1}{\sqrt{2}} \sum_{t=1}^m \partial_{\theta_t}(D^{-1}v_t). \quad (3.6.1)$$

The generating function of U_k therefore gives the exact common-frame inverse and score series throughout its convergence domain:

$$D^{-1} = \frac{1}{\gamma} \sum_{k \geq 0} (-\gamma^{-1})^k Q_k(D), \quad b = \frac{1}{\sqrt{2}\gamma} \sum_{k \geq 1} (-\gamma^{-1})^k \mathcal{C}_k. \quad (3.6.2)$$

Choose $K_n = \lceil L_1 \log n \rceil$. If $c > A_{\text{sh}}^2$, then

$$\|b_{\leq K_n}\|_2 \leq C \frac{\sqrt{n}}{c}. \quad (3.6.3)$$

The choices are noncircular. Equation (3.1.10) permits fixed numbers $0 < \alpha < \beta < 2$ and $c_0 < \infty$ such that, for every $c \geq c_0$, the inner band and a slightly larger outer band lie in $[\alpha c, \beta c]$. The characteristic roots in the proof of Lemma 3.47 depend continuously on $(c^{-1}, x/c)$; compactness of $[0, c_0^{-1}] \times [\alpha, \beta]$ and the strict Jury inequalities give one $q_0 < 1$ valid for all these bands and all sufficiently large n . Choose $L_1 |\log q_0| > 3/2$, invoke Theorem 2.24 only in that now-fixed logarithmic degree range to obtain A_{sh} , and finally enlarge c so that $c > A_{\text{sh}}^2$. This enlargement preserves the same q_0 .

Choose once and for all

$$0 < a'_c < a_c < b_c < b'_c < 2c + 1. \quad (3.6.3a)$$

For all sufficiently large n , $b'_c < 2(\gamma + 1)$. For a scalar $x > 0$, the absolute convergence criterion for Equation (3.6.2) is exactly $0 < x < 2(\gamma + 1)$: substitute $z = -\gamma^{-1/2}$ and $y = (x - \gamma - 1)/(2\sqrt{\gamma})$ in $\sum_{k \geq 0} U_k(y)z^k = (1 - 2yz + z^2)^{-1}$. Thus Equation (3.6.2) converges in operator norm on the compact band $a'_c I \preceq D \preceq b'_c I$. The following elementary estimate controls both the tail and its Fréchet derivative.

Lemma 3.47 (Finite- N inverse tail). *Let $\gamma > 1$ and let $[a, b] \subseteq (0, 2(\gamma + 1))$. Put*

$$S_k(x) = (-\gamma^{-1})^k Q_k(x), \quad R_K(D) = D^{-1} - \frac{1}{\gamma} \sum_{k=0}^K S_k(D).$$

There are $C < \infty$ and $q < 1$, depending only on (γ, a, b) , such that every self-adjoint D with spectrum in $[a, b]$ satisfies

$$\|R_K(D)\| \leq Cq^{K+1}, \quad \|dR_K(D)[E]\| \leq C(K+1)q^K \|E\|. \quad (3.6.4)$$

Proof. The recurrence Equation (3.2.6) gives

$$\sum_{k \geq 0} S_k(x)z^k = \left(1 + \frac{x - \gamma - 1}{\gamma}z + \frac{1}{\gamma}z^2\right)^{-1}. \quad (3.6.4a)$$

Thus $S_k(x)$ has characteristic polynomial

$$r^2 + \frac{x - \gamma - 1}{\gamma}r + \frac{1}{\gamma}.$$

The quadratic Jury criterion places both roots strictly inside the unit disc precisely when

$$\gamma > 1, \quad x > 0, \quad x < 2(\gamma + 1).$$

Compactness of $[a, b]$ gives uniform constants $C < \infty$ and $q < 1$ such that $|S_k(x)| \leq Cq^k$. The spectral theorem proves the first estimate in Equation (3.6.4).

Multiplying Equation (3.2.7) by $(-\gamma^{-1})^k$ gives

$$dS_k(D)[E] = -\frac{1}{\gamma} \sum_{j=0}^{k-1} S_{k-1-j}(D)ES_j(D). \quad (3.6.4b)$$

Consequently

$$\|dS_k(D)[E]\| \leq \frac{C^2}{\gamma} kq^{k-1}\|E\|.$$

Summing over $k > K$ proves the second estimate in Equation (3.6.4). $\square$

Write R_{K_n} for this tail on $[a'_c, b'_c]$, and let $q = q(c) < 1$ be supplied by Lemma 3.47. At a pivot,

$$\partial_{\theta_t} D = [A_t, D_{<t}], \quad \|D_{<t}\| \leq \|D\|, \quad (3.6.5)$$

where the last inequality is positivity on the band. The score tail is exactly

$$b - b_{\leq K_n} = \frac{1}{\sqrt{2}} \sum_{t=1}^m dR_{K_n}(D)[[A_t, D_{<t}]]v_t. \quad (3.6.5a)$$

Since $\|A_t\| \leq C$, $\|v_t\| = 1$, and $m = cN$, Lemma 3.47 gives

$$\|b - b_{\leq K_n}\| \leq C_c N(K_n + 1)q^{K_n}. \quad (3.6.6)$$

Choose $L_1|\log q| > 3/2$. Since $N \asymp n^2$, the right side is $o(\sqrt{n}/c)$. The inverse and b need not exist on singular index words. If w is any cutoff supported in this compact convergence band, the valid conclusion is the cutoff-local estimate

$$\left(\mathbb{E}[w\|b\|^2]\right)^{1/2} \leq \|b_{\leq K_n}\|_2 + \sup_{\text{supp } w} \|b - b_{\leq K_n}\| \leq C \frac{\sqrt{n}}{c} + o(\sqrt{n}/c). \quad (3.6.7)$$

4 Spectral surgery, endpoint calculus, and total variation

We give the endpoint integration-by-parts argument in full. Fix an index word and regard its endpoint as a map $F : \mathbb{T}^m \rightarrow \text{SO}(n)$. Its right-trivialized Jacobian is

$$J_{\theta}u = \sum_{t=1}^m u_t j_t, \quad M_{\theta} = J_{\theta}J_{\theta}^* = \sum_t j_t j_t^* = 2D_m. \quad (4.1.1)$$

The angle torus has normalized Lebesgue measure. The calculation below is then averaged over the finite set of index words.

Lemma 4.1 (Weighted submersion integration by parts). *Let ψ be Lipschitz with $0 \leq \psi \leq 1$, supported where $M_{\theta} \succeq aI$ for some $a > 0$, put $w = \psi^2$ and $p = \mathbb{E}w > 0$. The pushforward of $w d\mathbb{P}/p$ by F has a Haar density f_w . If b is the pathwise vector in Equation (3.1.7), then*

$$I_H(f_w) := \int_{\text{SO}(n)} \|\nabla \log f_w\|^2 f_w dH \leq \frac{2}{p} \mathbb{E}[w\|b\|^2] + \frac{8}{ap} \mathbb{E}\|\nabla_{\theta}\psi\|^2. \quad (4.1.2)$$

Proof. For $x \in \mathfrak{so}(n)$, define its minimum-norm horizontal lift

$$h_x(\theta) = J_\theta^* M_\theta^{-1} x, \quad (h_x)_t = \langle j_t, M_\theta^{-1} x \rangle. \quad (4.1.3)$$

Then $J_\theta h_x = x$, so for every smooth φ on the group,

$$d(\varphi \circ F)_\theta[h_x] = (R_x \varphi)(F(\theta)), \quad (4.1.4)$$

where R_x is the right-invariant derivative. Since $\partial_{\theta_t} j_t = 0$ and $\partial_{\theta_t} M = [A_t, M_{<t}]$,

$$\begin{aligned} \operatorname{div}_\theta h_x &= \sum_t \partial_{\theta_t} \langle j_t, M^{-1} x \rangle \\ &= - \left\langle M^{-1} \sum_t [A_t, M_{<t}] M^{-1} j_t, x \right\rangle = \langle b, x \rangle. \end{aligned} \quad (4.1.5)$$

Thus Equation (3.1.7) is obtained from the endpoint map itself; it is not imported from a coupling theorem.

The coarea theorem supplies a density because J is surjective on a neighbourhood of $\operatorname{supp} \psi$. On that neighbourhood define h_x by Equation (4.1.3). The Sobolev vector field $\psi^2 h_x$, extended by zero, is globally defined on the torus; approximation by smooth compactly supported fields gives, without a boundary term,

$$\mathbb{E}[\psi^2 (R_x \varphi)(F)] = -\mathbb{E} \left[\varphi(F) \{ \psi^2 \langle b, x \rangle + 2\psi \langle \nabla_\theta \psi, h_x \rangle \} \right]. \quad (4.1.6)$$

Let $\widehat{\mathbb{P}}$ be the tilted path law $d\widehat{\mathbb{P}} = \psi^2 d\mathbb{P}/p$. Haar invariance and Equation (4.1.6) show in the weak sense that the right Haar score of f_w is the conditional expectation, under $\widehat{\mathbb{P}}$ given F , of

$$q_\psi = b + 2\psi^{-1} M^{-1} J \nabla_\theta \psi \quad \text{on } \{\psi > 0\}. \quad (4.1.7)$$

Conditional Jensen and $(M^{-1}J)(M^{-1}J)^* = M^{-1} \preceq a^{-1}I$ give

$$\begin{aligned} I_H(f_w) &\leq \mathbb{E}_{\widehat{\mathbb{P}}} \|q_\psi\|^2 \\ &\leq \frac{2}{p} \mathbb{E}[\psi^2 \|b\|^2] + \frac{8}{ap} \mathbb{E} \|\nabla_\theta \psi\|^2. \end{aligned}$$

The quotient in Equation (4.1.7) is needed only on $\{\psi > 0\}$, and its tilted energy is the finite second term above. This proves Equation (4.1.2). $\square$

Choose deterministic nested bands

$$\mathcal{B}_{\text{in}} = \{a_c I \preceq D_m \preceq b_c I\}, \quad \mathcal{B}_{\text{out}} = \{a'_c I \preceq D_m \preceq b'_c I\}, \quad (4.1.8)$$

with the inner band from Proposition 3.3 and the outer endpoints from Equation (3.6.3a). In particular the entire outer band lies in the operator-norm convergence domain of Equation (3.6.2). Choose one-dimensional piecewise-linear functions $\eta_-, \eta_+ \in [0, 1]$ such that $\eta_-(x) = 0$ for $x \leq a'_c$, $\eta_-(x) = 1$ for $x \geq a_c$, $\eta_+(x) = 1$ for $x \leq b_c$, and $\eta_+(x) = 0$ for $x \geq b'_c$. Define the concrete Lipschitz spectral cutoff

$$\psi(D_m) = \eta_-(\lambda_{\min}(D_m)) \eta_+(\lambda_{\max}(D_m)).$$

It equals one on $\mathcal{B}_{\text{in}}$ and zero off $\mathcal{B}_{\text{out}}$. Put

$$w = \psi^2, \quad p = \mathbb{E} w \geq 1 - \delta_n. \quad (4.1.9)$$

Lemma 4.2 (Cutoff gradient). *For this cutoff,*

$$\mathbb{E}\|\nabla_{\theta}\psi\|^2 \leq C_c n \delta_n. \quad (4.1.10)$$

Proof. For $\tau > 0$ define the smooth spectral soft maximum and its density matrix by

$$\Lambda_{\tau}(D) = \tau \log \operatorname{Tr} e^{D/\tau}, \quad P_{\tau}(D) = \frac{e^{D/\tau}}{\operatorname{Tr} e^{D/\tau}}. \quad (4.1.11)$$

Its derivative in the direction E is $\operatorname{Tr}(P_{\tau}E)$. We first prove a bound uniform over every density matrix $P = \sum_r \alpha_r x_r x_r^*$. Jensen's inequality, skew-adjointness of A_t , and $D_{<t} \preceq D_m \preceq b'_c I$ give

$$\begin{aligned} |\operatorname{Tr} P[A_t, D_{<t}]|^2 &\leq \sum_r \alpha_r |\langle x_r, [A_t, D_{<t}] x_r \rangle|^2 \\ &\leq 4(b'_c)^2 \sum_r \alpha_r \|A_t x_r\|^2. \end{aligned} \quad (4.1.12)$$

For $T_x : y \mapsto [y, x]$, $M = 2D_m \preceq 2b'_c I$ and the adjoint Casimir identity give

$$\begin{aligned} \sum_t \|A_t x\|^2 &= \sum_t \|[j_t, x]\|^2 = \operatorname{Tr}(T_x M T_x^*) \\ &\leq 2b'_c \operatorname{Tr}(T_x T_x^*) = 2b'_c(n-2)\|x\|^2, \end{aligned} \quad (4.1.13)$$

and therefore

$$\sum_t |\operatorname{Tr} P[A_t, D_{<t}]|^2 \leq 8(b'_c)^3(n-2). \quad (4.1.13a)$$

Apply [Equation \(4.1.13a\)](#) first to $P_{\tau}(D_m)$ and then to $P_{\tau}(-D_m)$. Since $\Lambda_{\tau}(D) \rightarrow \lambda_{\max}(D)$ and $-\Lambda_{\tau}(-D) \rightarrow \lambda_{\min}(D)$ uniformly as $\tau \downarrow 0$, weak lower semicontinuity of the Sobolev energy proves

$$\|\nabla_{\theta}\lambda_{\max}(D_m)\|^2 + \|\nabla_{\theta}\lambda_{\min}(D_m)\|^2 \leq 16(b'_c)^3(n-2) \quad (4.1.13b)$$

almost everywhere on $\mathcal{B}_{\text{out}}$.

Let $\ell_{\pm}$ be the Lipschitz constants of $\eta_{\pm}$. The product rule and $0 \leq \eta_{\pm} \leq 1$ show that

$$\|\nabla_{\theta}\psi\|^2 \leq 32(b'_c)^3(\ell_-^2 + \ell_+^2)(n-2)\mathbf{1}_{\mathcal{B}_{\text{in}}^c}. \quad (4.1.13c)$$

The derivative also vanishes off $\mathcal{B}_{\text{out}}$. Taking expectations and using $\Pr(\mathcal{B}_{\text{in}}^c) \leq \delta_n$ proves [Equation \(4.1.10\)](#). $\square$

On the support of w , $M = 2D_m \succeq 2a'_c I$. [Lemma 4.1](#) with $a = 2a'_c$, the cutoff-local score estimate [Equation \(3.6.7\)](#), and [Lemma 4.2](#) give

$$I_H(f_w) \leq C \frac{n}{c^2} + C_c n \delta_n + o(n/c^2). \quad (4.1.14)$$

For completeness, the final functional-inequality step is as follows; we use the standard Bakry–Émery normalization of the Haar log-Sobolev inequality [\[BGL14\]](#). For the Hilbert–Schmidt metric, the Killing form of $\mathfrak{so}(n)$ gives

$$\operatorname{Ric} = \frac{n-2}{4}g. \quad (4.1.15)$$

On a compact manifold with $\operatorname{Ric} \geq \kappa_{\operatorname{Ric}}g$, the Bochner identity implies $I_H(P_t f) \leq e^{-2\kappa_{\operatorname{Ric}}t} I_H(f)$ for the heat semigroup, while $d \operatorname{Ent}_H(P_t f)/dt = -I_H(P_t f)$. Integrating from zero to infinity gives

$$\operatorname{Ent}_H(f) \leq \frac{1}{2\kappa_{\operatorname{Ric}}} I_H(f). \quad (4.1.16)$$

With $\kappa_{\text{Ric}} = (n - 2)/4$, Pinsker's inequality therefore yields

$$\|f_w H - H\|_{\text{TV}} \leq \sqrt{\frac{I_H(f_w)}{n - 2}} \leq \frac{C}{c} + C_c \sqrt{\delta_n} + o(1). \quad (4.1.17)$$

Because $0 \leq w \leq 1$, the original endpoint law is the positive mixture

$$\mu_m = p\mu_w + (1 - p)\mu_{\text{bad}}.$$

Thus

$$\|\mu_m - H\|_{\text{TV}} \leq \frac{C}{c} + C_c \sqrt{\delta_n} + \delta_n + o(1). \quad (4.1.18)$$

Given $\varepsilon > 0$, choose the fixed $c = c(\varepsilon)$ so that the first term is below $\varepsilon/2$, then take n large. Left equivariance makes the estimate uniform over the starting rotation, and total variation contracts under further steps. This proves the upper bound in [Theorem 1.1](#).

For the lower bound, condition on a coordinate word of length t . Its endpoint is the image of a smooth map $\mathbb{T}^t \rightarrow \text{SO}(n)$. If $t < N$, the area formula (equivalently, Sard's theorem) makes this image Haar-null. There are only N^t words, so their mixture is Haar-singular and has total-variation distance one. Hence $t_{\text{mix}}^{(n)}(\varepsilon) \geq N$.

5 Dependency audit

Stage	Direct input	Location
Circuit-anchored physical root/pass bridge	Exact Bianchi root, typed 2 3 pass, twenty transports, and one base per original circuit	Theorem 2.4
One-resource all-order comb lift	Physical bridge, actual-pass serialization, and atomic circuit contraction	Theorem 2.14
Positive-credit packet presentation and ribbon estimate	Comb lift, maximizing smoothing, Euler defect, and simultaneous gluing	Theorem 2.16 and Theorem 2.24
Constant covariance band	Ribbon moments and the quantitative pseudo-Haar input	Proposition 3.3
Local temporal calculus	Direct first-shell estimate, Kac Riesz blocks, chronological exposure, passive renewal, Haar-residual splitting, and typed root estimates	Proposition 3.5 and Sections 3.2 to 3.3
Root-fused allocation	Fixed-component renewal, vertex-labelled packets, and the one-base circuit majorant	Theorem 3.43
Packetless aggregate-Ricci shell	Root-local Ricci estimates, low-cell exhaustion, the normalized invariant-column insertion, causal restoration of the marked root, and endpoint-energy closure	Lemmas 3.36 to 3.38
Global pure-Weyl and aggregate-Ricci renewal	Root-fused allocation, packetless Ricci shell, and the complete weak-order sums	Theorem 3.44 and Theorem 3.45

Stage	Direct input	Location
Intact marked-shell estimate	Direct first-shell estimate for $k = 1$; ribbon estimate and both renewal branches for $k \geq 2$	Proposition 3.5 and Theorem 3.46
Score and total-variation upper bound	Covariance band, marked shells, inverse tail, cutoff calculus, and weighted submersion integration by parts	Section 3.6 and Section 4
Geometric N lower bound	Area formula for a t -torus in the N -dimensional group	Section 4

Use of AI

OpenAI Codex using GPT-5.6 Sol was the primary AI system used in the discovery, audit, and preparation of this proof. Claude Opus 5 was later used by a collaborator for two audit passes. The scope, approximate duration, division of work, and current human-verification status are described in [Section 1.5](#). The author reviewed the manuscript and sources, made all final decisions, and remains responsible for every claim and any remaining errors.

References

- [BGL14] D. Bakry, I. Gentil, and M. Ledoux. *Analysis and Geometry of Markov Diffusion Operators*. Vol. 348. Grundlehren der mathematischen Wissenschaften. Cham: Springer, 2014. DOI: [10.1007/978-3-319-00227-9](#).
- [BM86] H.-J. Bandelt and H. M. Mulder. “Distance-Hereditary Graphs”. In: *Journal of Combinatorial Theory, Series B* 41.2 (1986), pp. 182–208. DOI: [10.1016/0095-8956\(86\)90043-2](#).
- [CCL03] E. A. Carlen, M. C. Carvalho, and M. Loss. “Determination of the Spectral Gap for Kac’s Master Equation and Related Stochastic Evolutions”. In: *Acta Mathematica* 191.1 (2003), pp. 1–54. DOI: [10.1007/BF02392695](#). arXiv: [math-ph/0109003](#).
- [CM09] B. Collins and S. Matsumoto. “On Some Properties of Orthogonal Weingarten Functions”. In: *Journal of Mathematical Physics* 50.11 (2009), p. 113516. DOI: [10.1063/1.3251304](#). arXiv: [0903.5143](#).
- [CM17] B. Collins and S. Matsumoto. “Weingarten Calculus via Orthogonality Relations: New Applications”. In: *ALEA, Latin American Journal of Probability and Mathematical Statistics* 14.1 (2017), pp. 631–656. DOI: [10.30757/ALEA.v14-31](#). arXiv: [1701.04493](#).
- [CS06] B. Collins and P. Śniady. “Integration with Respect to the Haar Measure on Unitary, Orthogonal and Symplectic Group”. In: *Communications in Mathematical Physics* 264.3 (2006), pp. 773–795. DOI: [10.1007/s00220-006-1554-3](#). arXiv: [math-ph/0402073](#).
- [DS00] P. Diaconis and L. Saloff-Coste. “Bounds for Kac’s Master Equation”. In: *Communications in Mathematical Physics* 209.3 (2000), pp. 729–755. DOI: [10.1007/s002200050036](#).
- [Has70] W. K. Hastings. “Monte Carlo Sampling Methods Using Markov Chains and Their Applications”. In: *Biometrika* 57.1 (1970), pp. 97–109. DOI: [10.1093/biomet/57.1.97](#).
- [HJ17] B. Hough and Y. Jiang. “Cut-off Phenomenon in the Uniform Plane Kac Walk”. In: *The Annals of Probability* 45.4 (2017), pp. 2248–2308. DOI: [10.1214/16-AOP1111](#). arXiv: [1211.2031](#).

- [Jan01] É. Janvresse. “Spectral Gap for Kac’s Model of Boltzmann Equation”. In: *The Annals of Probability* 29.1 (2001), pp. 288–304. DOI: [10.1214/aop/1008956330](https://doi.org/10.1214/aop/1008956330).
- [Jia12] Y. Jiang. “Total Variation Bound for Kac’s Random Walk”. In: *The Annals of Applied Probability* 22.4 (2012), pp. 1712–1727. DOI: [10.1214/11-AAP810](https://doi.org/10.1214/11-AAP810). arXiv: [0905.1539](https://arxiv.org/abs/0905.1539).
- [Jia17] Y. Jiang. “Kac’s Random Walk on the Special Orthogonal Group Mixes in Polynomial Time”. In: *Proceedings of the American Mathematical Society* 145.10 (2017), pp. 4533–4541. DOI: [10.1090/proc/13598](https://doi.org/10.1090/proc/13598).
- [Kac56] M. Kac. “Foundations of Kinetic Theory”. In: *Proceedings of the Third Berkeley Symposium on Mathematical Statistics and Probability, Volume 3: Contributions to Astronomy and Physics*. Berkeley: University of California Press, 1956, pp. 171–197. URL: <https://projecteuclid.org/euclid.bsmsp/1200501656>.
- [Mas03] D. K. Maslen. “The Eigenvalues of Kac’s Master Equation”. In: *Mathematische Zeitschrift* 243.2 (2003), pp. 291–331. DOI: [10.1007/s00209-002-0466-y](https://doi.org/10.1007/s00209-002-0466-y).
- [Oli09] R. I. Oliveira. “On the Convergence to Equilibrium of Kac’s Random Walk on Matrices”. In: *The Annals of Applied Probability* 19.3 (2009), pp. 1200–1231. DOI: [10.1214/08-AAP550](https://doi.org/10.1214/08-AAP550). arXiv: [0705.2253](https://arxiv.org/abs/0705.2253).
- [PS07] I. Pak and S. Sidenko. “Convergence of Kac’s Random Walk”. Preprint. 2007. URL: <https://www.math.ucla.edu/~5C%7Epak/papers/psid3.pdf>.
- [PS16] N. S. Pillai and A. Smith. *On the Mixing Time of Kac’s Walk and Other High-Dimensional Gibbs Samplers with Constraints*. 2016. arXiv: [1605.08122](https://arxiv.org/abs/1605.08122). URL: <https://arxiv.org/abs/1605.08122>.
- [PS17] N. S. Pillai and A. Smith. “Kac’s Walk on n -Sphere Mixes in $n \log n$ Steps”. In: *The Annals of Applied Probability* 27.1 (2017), pp. 631–650. DOI: [10.1214/16-AAP1214](https://doi.org/10.1214/16-AAP1214). arXiv: [1507.08554](https://arxiv.org/abs/1507.08554).
- [PS26a] N. S. Pillai and A. Smith. *Kac’s Walk on Rotation Matrices Mixes in $n^2 \log n$ Steps*. 2026. arXiv: [2604.23828](https://arxiv.org/abs/2604.23828). URL: <https://arxiv.org/abs/2604.23828>.
- [PS26b] N. S. Pillai and A. Smith. *Mixing on k Columns of the Transvection Walk*. 2026. arXiv: [2605.27212](https://arxiv.org/abs/2605.27212). URL: <https://arxiv.org/abs/2605.27212>.
- [PSV26] N. S. Pillai, A. Smith, and V. Vaikuntanathan. *On the Pseudo-Mixing of Kac’s Walk*. 2026. arXiv: [2608.17374](https://arxiv.org/abs/2608.17374). URL: <https://arxiv.org/abs/2608.17374>.
- [Ros94] J. S. Rosenthal. “Random Rotations: Characters and Random Walks on $SO(N)$ ”. In: *The Annals of Probability* 22.1 (1994), pp. 398–423. DOI: [10.1214/aop/1176988864](https://doi.org/10.1214/aop/1176988864).
- [Tra09] L. Traldi. *Binary Nullity, Euler Circuits and Interlace Polynomials*. 2009. arXiv: [0903.4405](https://arxiv.org/abs/0903.4405). URL: <https://arxiv.org/abs/0903.4405>.
- [Tro11] J. A. Tropp. “Freedman’s Inequality for Matrix Martingales”. In: *Electronic Communications in Probability* 16.25 (2011), pp. 262–270. DOI: [10.1214/ECP.v16-1624](https://doi.org/10.1214/ECP.v16-1624). arXiv: [1101.3039](https://arxiv.org/abs/1101.3039).